



Mapping Rice Area in the Cauvery Delta Zone of Tamil Nadu Using Sentinel 1A Synthetic Aperture Radar (SAR) Data

M. Ajay Prakash^{a+++*}, S. Pazhanivelan^{b#},
D. Muthumanickam^{a†}, K. P. Rangunath^{b‡}
and A. P. Sivamurugan^{b^}

^a Department of Remote Sensing and GIS, Tamil Nadu Agricultural University, Coimbatore, India.

^b Centre for Water and Geo-Spatial Studies (CWGS), TNAU, Coimbatore, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i102630

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

<https://www.sdiarticle5.com/review-history/104540>

Short Research Article

Received: 06/06/2023

Accepted: 11/08/2023

Published: 12/08/2023

ABSTRACT

Since remote sensing based crop inventory provides accurate and timely information as compared to the conventional survey methods of estimating area, Multi-temporal Sentinel 1A Synthetic Aperture Radar data was used for the estimation of rice area during Samba season 2022 in the Cauvery delta zone comprising Thanjavur, Thiruvaru, Mayiladuthurai, and Nagapatnam districts of

⁺⁺ PG Scholar;

[#] Director (CWGS);

[†] Professor & Head;

[‡] Associate Professor (SS&AC);

[^] Associate Professor (Agronomy);

*Corresponding author: E-mail: aj23723@gmail.com;

Tamil Nadu. SAR data was preferred over optical satellite data due to excess cloud cover during cropping the major season in Tamil Nadu. Temporal back-scatter (dB) signature of rice crop was generated from the multi-temporal processed SAR data utilizing the modules of a fully automated MAPscape software aiding the discriminating of the crop from others. The signatures revealed that the dB levels to be the lowest during agronomic floods, reached the highest during maximum tillering stage and started declining thereafter. Multi-temporal feature extraction module of Mapscape was used to estimate rice area and validated for accuracy using ground truth data collected during survey. A total of 3.05 lakh ha of rice area was estimated with an overall accuracy of 90.8 % and 0.82 kappa coefficient. Largest area of 1.12 lakh ha was recorded in Thanjavur followed by Thiruvavur and Mayiladuthurai with 0.95 and 0.51 lakh ha respectively.

Keywords: Rice area mapping; sentinel-1a; synthetic aperture radar; cauvery delta zone.

1. INTRODUCTION

Rice is the most significant food crop in the world, feeding half of all people. For 20% of their daily calories, more than 3.5 billion people rely on rice. Rice is cultivated throughout the globe, it supports the economy in many nations and accounts for more than 150 million hectares of rice fields or 11% of the world's agriculture. The consumption of rice has rapidly increased due to high population growth and shifting consumer tastes, and the output of rice is predicted to reach 555 million metric tonnes in 2035 [1-3]. The globe rice cultivation, from the wet regions to the harsh deserts, in a variety of climatic conditions. In South, South-East, and East Asia, rice takes up an exceptionally large proportion of the overall planted area. India is one of the largest producers and consumers of rice in the world. Rice is the staple food for a significant portion of the Indian population, and it plays a crucial role in the country's agriculture and food security. The total area under rice cultivation in India varies from year to year depending on factors such as weather conditions, government policies, and market demand. However, it is generally around 43-45 million hectares with an annual rice production of around 110-120 million metric tons. Rice cultivation in Tamil Nadu typically varies from 2.5 to 3.0 million hectares contributing around 7 to 9 million metric tons of rice annually to India's overall production. India produced 12.436 million metric tonnes of rice on 4.56 million hectares of rice fields in 2020–2021, with a productivity of 2717 kg/ha. In 2022-2023, Kharif's output target is 112000 tonnes, while Rabi's target is 18500 tonnes.

Based on rainfall, soil characteristics, irrigation potential, and cropping patterns, Tamil Nadu has seven different agro-climatic zones. Tamil Nadu is the country's tenth-biggest and seventh-most populated state (6%). It has a total geographical

area of 130.33 Lha, which is 4% of the total land area of the country. The availability of water resources per capita is 750 cubic metres per year, compared to the national average of 2,200 cubic metres. The state has 81.18 lakh operational land holdings, operating 64.88 Lha. Small and marginal holdings account for 92% of overall holdings, with 61% of the total sown area under cultivation. In Tamil Nadu, Rice productivity in the 2020–2021 growing season in Kharfi and Rabi was 3483 and 4436 metric tonnes (Total). Tamil Nadu had a rice coverage of 2024.33/Hectare and 192.94/hectare, respectively [4].

The information about the paddy area and the crop growth conditions were helpful topolicymakers, government, farm managers, and farmers in formulating policies and targeting interventions. Land use map of Cauvery delta zone was prepared using satellite data and it revealed the crop lands were classified in this zone. The distribution of land use, (124927 ha, 54.3%), crop (1069 ha, 0.46%), crop (17461 ha, 7.59%) and aquaculture (721 ha, 0.31%) within forest, marsh vegetation and deciduous forest area were 251 ha (0.11%) and 6782 ha (2.95%), respectively. This classification was help to find crop area distribution of total area at geographical area which is used by the government agency and use for policy making in farmers community [5].

The issues in consolidation of crop area and production through conventional methods warrant new innovative technologies usage. A cutting-edge approach of remote sensing allows real-time crop monitoring, mapping, crop inventory, and yield estimation [2]. Precise crop identification can produce precise figures on crop area spread, planting structure and regional distribution, as well as provide important input traits for crop yield prediction. Hence crop

identification and delineation is important for building a national food policy and economic strategy. Remote sensing data allow fast, accurate, detailed, and objective crop labelling, crop tracking, area estimation, and yield estimation. Remote sensing has been used in a wide range of earth observation activities due to the benefits of high time resolution, wide coverage, and low cost, and thus offers a useful tool for crop recognition and planting area tracking on a large scale [6]. Hence, the acquisition of data over vast geographic areas, remote sensing offers a cheaper alternative [7]. Crop inventory by remote sensing provides rapid and exact information, while traditional methods of area estimation are costly, time-consuming, less dependable, and less precise [8]. Based on biophysical characteristics of crops and/or soils, remote sensing technology has the potential to revolutionise the detection and characterisation of agricultural production [9].

The introduction of modern satellites, as well as the open availability of data, enhances the quality and accuracy of agricultural data. When compared to optical satellite data, synthetic aperture radar (SAR) has been extensively used in crop inventory because of its ability to penetrate clouds and be time-independent [2,10]. Sentinel-1A data is freely accessible to users, and the satellite was launched in 2014 by the European Space Agency for Earth Observation. It is usually utilize microwave data for area estimation rather than optical data since cloud cover is an inescapable phenomenon in the tropical region especially during the monsoon season. This SAR sensor can collect data day or night, regardless of cloud cover, by providing its own light source. Because of how effectively SARs react to these structural variations, these sensors can reliably identify the kind of crop being grown and have shown to be sensitive to a number of crop biophysical characteristics, including leaf area index, biomass, and canopy height. The benefits of combining SAR data from several frequencies (L-, C-, and X-band) for crop categorization were established by [11]. Every crop has a distinct backscattering coefficient, which has been used to identify various crops. It has a C-band Synthetic Aperture Radar that delivers data in dual-polarisation of both VV (Vertical-Vertical) and VH (Vertical-Horizontal), with data accessible at twelve-day intervals regardless of time or weather [6].

The temporal signature of the rice crop generated by SAR backscatter and its

relationship to the crop stages serve as the foundation for rice identification. Several studies has attempted to study the use of SAR data in estimation and monitoring of crop viz., rice, maize, cotton, mango, and banana [8, 12-19]. With this background, this study attempts to estimate the area and distribution of paddy in the Cauvery Delta region of Tamil Nadu during the Samba season 2022–23.

2. MATERIALS AND METHODS

2.1 Study Area

The Cauvery Delta Zone (CDZ) is situated in Eastern Tamil Nadu surrounded by the Palk Straight in the South, the Bay of Bengal in the East, Perambalur and Ariyalur districts to the West and Cuddalore district to the North. Rice is the primary crop in this area which is grown either in a single season or double season based on the availability of water and other climatic conditions. During Samba season (2022–23), this research was undertaken in the Cauvery Delta region comprising the districts of Thanjavur, Thiruvarur, Nagapattinam, and Mayiladuthurai (Fig.1).

2.2 Ground Truth Data

In the study area, an extensive survey was made and around 300 ground truth points were gathered during Samba, 2022 growing season comprising 200 rice points and 100 non-rice points. During the survey, rice and non-rice points were gathered with information on location, type of land use, age of crop and other ancillary data to aid in precise estimation of crop area. These points were utilized for training and validation of rice area estimates generated from the pre-processed satellite data in Mapscape software.

2.3 Satellite Data

SAR has the benefit of working at wavelengths that are not impeded by cloud cover or a lack of light, and it can capture data over a location at any time of day or night, in any weather. Sentinel-1A's C band sensor can offer consistent wide-area surveillance. Sentinel 1A Level 1 ground range (GRD) data with a resolution of 20m from August through January, 2022-23 were downloaded from the website (<https://sci-hub.copernicus.eu/dhus/>) and utilised in this investigation. Sentinel 1A VH polarisation data

with 12-day temporal resolution were utilised to estimate the paddy area.

2.4 Pre-processing of Sentinel 1A

SAR GRD multi-temporal data were processed using a fully automated chain processing methodology established by Holecz et al. [8] to provide terrain-geocoded σ° values in the MAPscape-RICE software. Basic SAR data processing comprises radiometric calibration, normalisation, strip mosaicking, co-registration, time-series speckle filtering, terrain geocoding, and anisotropic non-linear diffusion (ANLD) filtering. Atmospheric attenuation is eliminated (Nelson et al., 2014). On processing, this software translates temporal images to backscattering coefficients (σ°) based on terrain (Fig.2).

2.5 Classification Using Parameters

Multi-temporal feature viz., maximum, minimum, mean, and range of degrees were extracted from the temporal characteristics of satellite data for the ground truth points. Then, the (i) minima and (ii) maxima of those mean σ° values across fields; the (iii) maxima of the minimum σ° values across fields; the (iv) minima of the maximum σ° value across fields; and the (v) minimum and (vi)

maximum of the range of σ° values [8] were computed for the ground truth points. The parameters $t_{\text{minlength}}$, $t_{\text{maxlength}}$ and $t_2 * t_1$ are easier to estimate. $T_{\text{minlength}}$ restricts the number of days between a start-of-season detection and the subsequent highest σ° value in the temporal signature. Since X-band σ° saturates before rice flowering, this value can be set to 40–50 days. $T_{\text{maxlength}}$ restricts the duration between two σ° minima in the series and 120 days is a suitable cutoff that would be representative of an intensive triple rice system. $t_2 - t_1$ is the maximum period of agronomic flooding at the start of the season, which is set to be a high value of 40 to 50 to capture the longest land preparation phases.

2.6 Accuracy Assessment

The Error matrix and Kappa statistics were used to assess classification accuracy. A comparison of the paddy area map is used to determine accuracy and versus data from the ground. There are two types of validation points: paddy points and non-paddy points. Non-paddy points include all land cover groups other than paddy. The error matrix is used to determine accuracy measurements such overall accuracy, producer accuracy, user accuracy, and kappa value [20].




Fig. 1. Study area of Cauvery Delta region




Fig. 2. Rule-based rice detection algorithm from multi-temporal Sentinel 1A

3. RESULTS

The multi-temporal 12 days interval Sentinel 1A SAR images from August, 2022 and January, 2023 were downloaded, processed and stacked by transforming into terrain-geocoded σ° images. The SAR satellite data gathered throughout the cropping season was fed into the automated processing chain, and analysis was performed using training pixels collected via ground-truthing to develop a temporal back scattering signature (σ°) for the rice crop in the research region. The temporal characteristics recovered from VH polarisation were used to produce the dB curves for paddy in monitoring fields. Paddy's dB value varies from -24 to -12 dB. Using time series Sentinel 1A SAR images, a paddy area map comprising four Cauvery delta districts, Thanjavur, Thiruvarur, Nagapattinam, and Mayiladuthurai, was estimated. For the study region, district-level paddy area data were computed, and a summary of district-wise paddy area is shown in Table 1. In the research field, during the samba season 2022–23, 305388.58 hectares of rice area was estimated from multi-temporal Sentinel 1A SAR data using a parameterized classifier combining multi-temporal characteristics. Thanjavur had the

largest area of roughly 112534.24 hectares, followed by Thiruvarur, Mayiladuthurai, and Nagapattinam, with 95670.62 ha, 51248.20 ha, and 45935.52 ha, respectively.

Table 1. District wise rice area in Cauvery delta region during samba season 2022-23

S. No	District	Rice Area (ha)
1.	Mayiladuthurai	51248.20
2.	Nagapattinam	45935.52
3.	Thiruvarur	95670.62
4.	Thanjavur	112534.24
Total		305388.58

The ground truth points collected during the survey were used to assess the accuracy of the estimated rice area. There were 200 rice points and 100 non-rice points among the 300 ground truth points collected. A confusion matrix was built to validate the rice area computation and to assess classification accuracy. During samba season 2022, 180 of the 200 ground truth rice points obtained were correctly classified as rice crop. As a result, the overall accuracy of the rice area map was 90.8 %, and the kappa co-efficient was 0.82, indicating that the technique was accurate (Table 2).




Fig.3. Rice area in Cauvery Delta Zone during Samba 2022-23

Table 2. Confusion Matrix for accuracy assessment

		Predicted class from the map		
		Rice	Non-Rice	Accuracy
Actual class from	Rice	95	8	92.2%
	Non-Rice	3	14	82.4%
	Reliability	96.9%	63.6%	90.8%
Average accuracy				87.3%
Average reliability				80.3%
Overall accuracy				90.8%
Kappa index				0.82

Table 3. Block wise rice area during samba season 2022-23

S. No	District	Block	Area in Ha
1.	Mayiladuthurai	Kollidam	7867.16
		Sirkali	6962.44
		Mayiladuthurai	11851.20
		Sembanarkoil	12630.80
		Kuthalam	11936.60
2.	Nagapattinam	Thirumarugal	9167.28
		Nagappattinam	4465.72
		Kivelu	7398.12
		Keelaiyur	6494.56
		Talanayar	7894.84
		Vedharanyam	10515.00
3.	Thiruvarur	Kudavasal	9975.560
		Nannilam	9895.4
		Kordacheri	7458.20
		Thiruvarur	4932.36
		Needamangalam	13651.70
		Mannargudi	11215.10
		Kottur	15477.00
		Muthupettai	11567.40
		Thiruthuraipoondi	11497.90
		4.	Thanjavur
Kumbakonam	6214.04		
Thiruvaidaimaruthur	8370.00		
Papanasam	3428.20		
Ammappettai	14496.50		
Thanjavur	12378.60		
Thiruvaiyaru	5447.60		
Budalur	9312.04		
Orathanadu	17635.10		
Thiruvanam	8641.72		
Madukkur	3223.08		
Pattukottai	4908.80		
Sethubavachatram	4441.48		
Peravurani	5573.72		

Block wise statistics were collected to understand the pattern of rice area distribution throughout the research region's blocks. The Orathanadu block of Thanjavur district recorded the largest area with 17635.10 ha, followed by Kottur block of Thiruvarur district, Sembanarkoil block of Mayiladuthurai district and Vedharanyam

block of Nagapattinam doistrict with 15477.00 ha, 12630.80 ha, and 10515.00 ha, respectively. The Madukkur block in Thanjavur District reported the lowest rice area of 3223.08 ha, followed by Nagappattinam block, Thiruvarur block, and Sirkali block, with 4465.72 ha, 4932.36 ha and 6962.44 ha respectively (Table 3).

4. DISCUSSION

The study was conducted by Tamilmounika, R., et al., using Sentinel-2A data for estimation of rice area. During (2020-2021) Thanjavur recorded the highest area under paddy of about 122684 ha and followed by Thiruvarur, Mayiladuthurai and Nagapattinam with an area of 119379 ha, 57015 ha and 46250 ha, respectively. A total paddy area of 345328 ha. Similar to this Kannan, Sugavaneshwaran, et al. [18] used Sentinel-2A, calculated rice area around 1,41,639 ha, 1,25,497 ha, and 1,17,703 ha were rice-growing areas in Thanjavur, Thiruvarur, and Nagapattinam districts, respectively. The accuracy of the rice area map was assessed through the confusion matrix using the ground truth points to classify rice and non-rice pixels [21-26]. Rice points were classified with an accuracy of 95.3 per cent, while non-rice points were classified with an accuracy of 92.0 per cent. Considering the efficiency of the methodology of mapping rice area with SAR data, the overall accuracy of the rice area map was 94.5 per cent [27-30].

5. CONCLUSION

The Cauvery Delta Zone (CDZ) of Tamil Nadu rice area estimation approach was found to be successful in detection, delineation, and estimation, as shown by the high classification accuracy of 90.8% and a kappa value of 0.82. Due to its weather and time independence, SAR satellite data is perfect for immediate area mapping. In order to create policies and focus interventions, policymakers, the government, farm managers, and farmers used information regarding the paddy area and crop growth conditions. The results derived will be of much use in policy decisions due to its high accuracy estimates.

COMPETING INTERESTS

Authors have declared that they have no known competing financial interests or non-financial interests or personal relationships that could have appeared to influence the work reported in this paper.

ACKNOWLEDGEMENT

The authors would like to express their gratitude to the Director Centre for Water and Geospatial Studies, advisory committee members, Staff members and Colleagues Department of

Remote Sensing and GIS, Tamil Nadu Agricultural University, for their unwavering support and suggestions throughout the research process.

REFERENCES

1. Khush GS. "What it will take to feed 5.0 billion rice consumers in 2030". *Plant molecular Biology*. 2005; 59(1):1-6.
2. Zhang H, Q Li, X Du, and M Zhang. Estimate rice acreage in Hunan province using the China Environment Satellite data". In proceedings of 2013 IEEE International Geoscience and Remote Sensing Symposium-IGARSS. 2013;3254-3257.
3. Son D, Lee J, Qiao S, Ghaffari R, Kim J, Lee JE, Song C, Kim SJ, Lee DJ, Jun SW, Yang S. Multifunctional wearable devices for diagnosis and therapy of movement disorders. *Nature nanotechnology*. 2014 May;9(5):397-404.
4. *Indiastat*; 2021. Available: <https://www.indiastat.com/table/agriculture/area-under-cultivation-foodgrains-india-1950-1951-/7417>.
5. Pazhanivelan S, Kannan P, Christy Nirmala Mary P, Subramanian E, Jeyaraman S, Nelson A, Yadav M. Rice crop monitoring and yield estimation through COSMO Skymed and TerraSAR-X: A SAR-based experience in India. *The International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences*. 2015;40:85-92.
6. Raman MG, Kaliaperumal R, Pazhanivelan S. Rice Area Estimation in Tiruvarur District of Tamil Nadu using VV Polarized Sentinel 1A SAR Data. *Indian Journal of Natural Sciences*. 2017;8(44): 12782-12793.
7. De beurs K, Townsend P. Estimating the effect of gypsy moth defoliation using MODIS. *Remote Sens Environ*. 2008;112:3983-90.
8. Holecz F, Barbieri M, Collivignarelli F, Gatti L, Nelson A, Setiyono TD, Pazhanivelan S. An operational remote sensing based service for rice production estimation at national scale. In *Proceedings of the living planet symposium*; 2013.
9. Liaghat S, Balasundram SK. (A review: The role of remote sensing in precision agriculture. *American Journal of*

- Agricultural and Biological Sciences. 2010; 5(1):50-55.
10. Forkuor G, Conrad C, Thiel M, Ullmann T, Zoungrana E. Integration of optical and Synthetic Aperture Radar imagery for improving crop mapping in Northwestern Benin, West Africa. *Remote Sensing*. 2014;6(7):6472-6499.
 11. Shang J, McNairn H, Champagne C, Jiao X. Contribution of multi-frequency, multi-sensor, and multi-temporal radar data to operational annual crop mapping. *IEEE Transactions on Geoscience and Remote Sensing*. 2009;3(1):III378-III381.
 12. Setiyono TD, Quicho ED, Holecz FH, Khan NI, Romuga G, Maunahan A, Mabalay MRO. Rice yield estimation using synthetic aperture radar (SAR) and the ORYZA crop growth model: development and application of the system in South and South-east Asian countries. *International Journal of Remote Sensing*. 2019;40(21): 8093-8124.
 13. Sudarmanian NS, Pazhanivelan S, Panneerselvam S. Estimation of methane emission from paddy fields using sar and modis satellite data. *Journal of Agrometeorology*. 2019;21(1):102-109.
 14. Ramalingam K, Ramathilagam AB, Murugesan P. Area Estimation of Cotton and Maize Crops in Perambalur District of Tamil Nadu Using Multi Date SENTINEL-1A SAR Data & Optical Data. *The International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences*. 2019;42:137-140.
 15. Venkatesan M, Pazhanivelan S, Sudarmanian NS. Multi-temporal feature extraction for precise maize area mapping using time-series Sentinel 1A SAR data. *The International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences*. 2019;42:169-173.
 16. Kaliaperumal R, GR M. Mapping mango area using multi-temporal feature extraction from Sentinel 1A SAR data in Dharmapuri, Krishnagiri and Salem districts of Tamil Nadu. *Madras Agricultural Journal*. 2019;106.
 17. Karthikkumar A, Pazhanivelan S, Jagadeeswaran R, Ragunath KP, Kumaraperumal R. Generating Banana area map using VV and VH Polarized Radar Satellite Image. *Madras Agricultural Journal*. 2019;106.
 18. Kannan, Sugavaneshwaran, et al. Rice Area Estimation using Sentinel 1A SAR Data in Cauvery Delta Region. *Int. J. Curr. Microbiol. App. Sci*. 2021;10(2):848-853.
 19. Tamilmounika R, Pazhanivelan S, Ragunath KP, Sivamurugan AP, Sudarmanian NS, Kumaraperumal R, Thirumeninathan S. Paddy area estimation in Cauvery Delta Region Using Synthetic Aperture Radar. *International journal of Environment, Ecology and Conservation*. 2022;S517-S522.
 20. Congalton RG. A review of assessing the accuracy of classifications of remotely sensed data. *Remote sensing of environment*. 1991;37(1):35-46.
 21. Stroppiana D, Boschetti M, Azar R, Barbieri M, Collivignarelli F, Gatti L, Holecz F. In-season early mapping of rice area and flooding dynamics from optical and SAR satellite data. *European Journal of Remote Sensing*. 2019;52(1): 206-220.
 22. Zhang Y, Yan W, Yang B, Yang T, Liu X. Estimation of rice yield from a C-band radar remote sensing image by integrating a physical scattering model and an optimization algorithm. *Precision Agriculture*. 2020;21:245-263.
 23. Raviz J, Mabalay MR, Laborte A, Nelson A, Holecz F, Quilang EJ, Doverté M. (Mapping rice areas in Mindanao using the first images from Sentinel-1A: The PRISM Project experience. In 36th Asian Conference on Remote Sensing (ACRS), Manila. 2015;19-23.
 24. Nelson A, Setiyono T, Rala AB, Quicho ED, Raviz JV, Abonete PJ, Ninh NH. Towards an operational SAR-based rice monitoring system in Asia: Examples from 13 demonstration sites across Asia in the RIICE project. *Remote Sensing*. 2014;6(11):10773-10812.
 25. Nihar MA, Ahamed JM, Pazhanivelan S, Kumaraperumal R, Raj KG. Estimation of cotton and maize crop area in Perambalur district of Tamil Nadu using multi-date sentinel-1A SAR data. *The International Archives of Photogrammetry, Remote Sensing and Spatial Information Sciences*. 2019;42:67-71.
 26. Satheesh S, Pazhanivelan S, Kumaraperumal R, Radhamani S, Ragunath KP, Sudarmanian, NS. Area Assessment for Rice Crop in Thiruvavur District Assimilating Sentinel 1A Satellite Data. *International Journal of Environment*

- and Climate Change. 2022;12(10):1270-1278.
27. Doraiswamy, Paul C, et al. Application of MODIS derived parameters for regional crop yield assessment. Remote sensing of environment. 2005;97(2):192-202.
 28. Kingra PK, Debjyoti Majumder, Som Pal Singh. Application of remote sensing and GIS in agriculture and natural resource management under changing climatic conditions." Agricultural Research Journal. 2016;53(3).
 29. Kannan P, Sankar M, Prabhavathi M. Remote sensing and GIS for land use mapping and crop planning in Cauvery delta region of Tamil Nadu, India. Indian Journal of Soil Conservation. 2010; 38(3):199-204.
 30. Stroppiana D, Boschetti M, Azar R, Barbieri M, Collivignarelli F, Gatti L, Fontanelli G, Busetto L, Holecz F. In-season early mapping of rice area and flooding dynamics from optical and SAR satellite data. European Journal of Remote Sensing. 2019;52(1):206-220.

© 2023 Prakash et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/104540>