



Assessment of Integrated Pest Management Modules against Fall Army Worm and Its Economic Impact in Maize

V. Lakshmi Narayanamma ^a, V. Ratnakar ^{b*}, M. Ram Prasad ^c,
B. Shiva ^a, R. Vishwatej ^a, G. Veeranna ^d and R. Uma Reddy ^e

^a Krishi Vigyan Kendra, Professor Jayashankar Telangana State Agricultural University, Kothagudem, India.

^b Agricultural College, Professor Jayashankar Telangana State Agricultural University, Jagtial, India.

^c Agricultural College, Professor Jayashankar Telangana State Agricultural University, Aswaraopet, India.

^d DAATTC, Professor Jayashankar Telangana State Agricultural University, Warangal, India.

^e Regional Agriculture Research Station, Professor Jayashankar Telangana State Agricultural University, Warangal, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/V13i102949

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/102204>

Original Research Article

Received: 20/04/2023

Accepted: 23/06/2023

Published: 11/09/2023

ABSTRACT

Aim: The present study was undertaken to assess the IPM modules and their economic impact against Fall Armyworm in maize through on farm trail (OFT).

Place and Duration of Study: The present study was carried out in the farmers' fields of Bhadradi Kothagudem district of Telangana during Rabi 2019-20, 2020-21 and 2021-22.

Methodology: The main objective of on farm trail is to test and evaluate the findings of the research stations at the farmer's field and to refine and modify the technologies, if required for better adoption by farmers. The KVK, Bhadradi Kothagudem conducted on farm trail for the

*Corresponding author: E-mail: ratnakar.ento@gmail.com;

management of fall armyworm in maize during *Rabi* 2019-20, 2020-21 and 2021-22, as part of technical programme of Krishi Vigyan Kendra, Bhadradi Kothagudem in KVK operational area of adopted villages.

Results: In technology assessment plot (On farm trail) the treatments followed were avoiding staggered sowing of maize, installation of pheromone traps @ 8-10 per acre, clean cultivation, balanced application of fertilizers, erection of bird perches @10/acre, spraying of Azadiractin (1500ppm) to repel the egg laying and need based whorl application of Emamectin benzoate @ 0.5g/l and *Metarrhizium anisopliae* @ 5g/l of water, if more than 8 adult moths were trapped in the pheromone trap for three consecutive days. It was compared with the farmers practice plots i.e application of Carbofuran 3G granules @ 3kg/acre and indiscriminate usage of fertilizers and insecticides. The experiment was conducted in five locations. The benefit cost ratio (BC Ratio) was higher in technology assessed plots with 2.78, 2.63 and 2.75 whereas BC ratio was comparatively lower in farmers practice plots with 2.43, 2.21 and 2.25 in corresponding *Rabi* 2019-20, 2020-21 and 2021-22 respectively.

Conclusion: Technology assessed package proved its effectiveness among maize farmers with increased net returns, lower per cent of FAW incidence and high yield and BC ratio compared to the farmers practiced during *Rabi* 2019-20, 2020-21 and 2021-22, respectively. Hence, this on farm trial proved to be feasible for demonstration in wide scale to farming community of Bhadradi Kothagudem district.

Keywords: *Fall army worm; maize; invasive; polyphagous; pheromone; emamectin benzoate; Metarrhizium anisopliae.*

1. INTRODUCTION

In India, maize is the third most important cereal crop after rice and wheat, both in terms of area and production, registering maximum growth rate among food crops [1,2]. Though maize is emerging as an important industrial crop in India, its productivity (3.1 tonne/ha) is much lower than the world average (5.62 t/ha) [3,4], because of major biotic stresses. Among them, FAW is the major constraint limiting the production. It emerged as the most destructive pest of maize in India since its report in May, 2018 and spread to almost all the maize growing Indian states except Jammu and Kashmir. In 2019, Karnataka State had the highest area affected with FAW (2,11,300 ha) followed by Telangana (24,288 ha), Maharashtra (5,144 ha) and low infestation in Tamil Nadu, Madhya Pradesh, West Bengal and Arunachal Pradesh (< 500 ha) [3]

Fall armyworm (*Spodoptera frugiperda* (J. E. Smith))(Lepidoptera: Noctuidae), a native pest of North and South America. It is likely to be a real threat to Indian agriculture in near future as it is currently observed to be causing damage only on maize in India and likely to spread to other crops due to its Polyphagous nature. It can spread quickly across large geographic areas as adult moths can move up to 500 km. It does not undergo diapause in any stage of life cycle hence, can persist throughout the year on many crops and weeds.

The caterpillars of FAW feed mainly on leaves, munching on their edges and making holes, giving them a ragged and torn appearance. If the larvae are numerous, they can completely defoliate the plants. When they have exhausted their food source, they invade neighbouring vegetation - which may often be another crop - in large numbers sites close to the ground. By far most of the damage, however, is caused by the last instar larva, which consumes more plant biomass than all of the other younger larval instars put together. This factor is often responsible for people thinking and assessing that the attacks by fall armyworm are sudden and occurred at once. In corn, the caterpillars feed deep in the whorl, often concealed under their own frass. While the leaves may be heavily damaged, the plant usually compensates for the loss. If the growing point is destroyed, however, "dead heart" symptoms will appear. As the crop grows tasselling stage, emerging tassel often pushes the feeding larva out of the whorl. At this stage, some larvae may start feeding on the side leaves while some larvae may shift to the developing cob. The larvae shifted to developing cob frequently feed on the immature kernels inside the ear. This generally causes greater injury than the leaf damage. Further, the extent of loss depends on the stage of cob development at the time of attack and the extent of feeding by the larva on the cob. When the larvae are very numerous they defoliate the preferred plants, acquire an "armyworm" habit

and disperse in large numbers, consuming nearly all vegetation in their path. Due to cannibalistic behaviour of FAW, the larvae as they grow in size are forced to disperse to neighbouring plants from the plant where eggs are laid. Hence, only one or very few larvae can be found per plant, despite adults laying egg masses comprising hundreds of eggs [5].

Bhadradi Kothagudem district falls under Central Telangana Zone of Telangana State with tropical climate typically receives about 147.76 millimeters (5.82 inches) of precipitation with 135.35 rainy days annually. The reduction in maize growing areaduring 2019-20 issolely attributed to the failure of the maize crop due to infestation of Fall Army Worm (FAW). Farmers' resort to spray various insecticides within short intervals resulted in resistance, secondary pest outbreak and pest resurgence along with destruction of natural enemies and environmental pollution.The farmers invested on an average of Rs.10,000-12,000/acre on pesticides for the managementof this dreaded pest. This resulted in severe damage to the natural resources as well as natural enemies and reduction in net income of the farmers. Considering the above situation and to overcome this menace, KVK, Bhadradi Kothagudem conducted an on farm trial (OFT)“Assessment of integrated pest management modulesagainst fall armyworm and its economic impact in Maize” in a systematic manner on farmers' fields to show the worth of a technology and its feasibility in Bhadradi Kothagudem by convincing farmers about potentialities of improved management practices of maize for further adoption.

2. MATERIALS AND METHODS

The present study was undertaken at five farmers fields with two plots per location (0.2ha/plot) in Bhadradi Kothagudem district of Telangana state, India in medium black soils with cropping patterns of maize followed by summer pulses with two treatments viz.,technology assessment (treatment) and check (farmers practice). The experiment was conducted consecutively for three years i.e. during the *Rabi* season (October–February) of 2019-20, 2020-21 and 2021-22. The experimental design, site and farmer selection, the layout of demonstration, farmer's participation, etc., were followed as suggested by Choudhary [6].

In technology assessment plot (On farm trail) the treatments followed were avoiding staggered sowing of maize, installation of pheromone traps @ 8-10 per acre, clean cultivation, balanced application of fertilizers, erection of bird perches @10/acre, spraying of Azadiractin (1500ppm) to repel the egg laying and need based whorl application of Emamectin benzoate @ 0.5g/l and *Metarrhizium anisopliae* @ 5g/l of water, if more than 8 adult moths were trapped in the pheromone trap for three consecutive days. It was compared with the farmers practice plots *i.e* application of Carbofuran 3G granules @ 3kg/acre and indiscriminate usage of fertilizers and insecticides. The farmers don't have the sufficient knowledge on IPM components to manage the FAW in maize crop.

The pest incidence was calculated by randomly selecting 20 plants and observing the damage symptoms visually. The yield parameters were collected from both technology assessment plot and farmers practiced plot by random crop cutting method and analyzed by using simple statistical tools. Using the yield parameters extension gap, technology gap, yield gap, technology index was calculated as procedure suggested by Rajashekhar et al. [7] and Samui et al. [8]. To find out the economic impact of treatments effected due to FAW incidence and yield the benefit cost ratio was calculated.

Extension gap (q/ha) = Demonstrations yield –Yield under existing farmer's practice

Technology gap (q/ha) = Potential Yield – Demo Yield

Additional return = Demonstration return – farmer's practice return

$$Yield\ gap\ (\%) = \frac{Extension\ gap}{Yield\ under\ farmers\ practice} \times 100$$

$$Technology\ gap\ (\%) = \frac{Technology\ gap}{Potential\ yield} \times 100$$

$$Technology\ index\ (\%) = \frac{Potential\ yield - Demonstration\ yield}{Potential\ yield} \times 100$$

3. RESULTS AND DISCUSSION

The results of the present study state that the incidence of fall armyworm was lower in the technology assessed plot compared to the

farmers practice. The reason is attributed to the regular monitoring of the fall armyworm by erection of pheromone traps followed by prophylactic spraying of Azadiractin 1500ppm which act as a strong oviposition deterrent & repellent against FAW adults. Further, if more than 8 adult moths were trapped in the pheromone trap for three consecutive days, the technology assessed plot was sprayed with Emamectin benzoate @ 0.5g and *Metarrhizium anisopliae* @ 5g/l of water which further reduced the FAW incidence up to 18-40 per cent compared to farmer practice (Fig. 1).

The highest yield of 71.0, 65.0 and 68.8 (Q/ha) was recorded in the technology assessment plot, while lower yield of 63.50, 58.75 and 59.20 (Q/ha) was recorded in the farmers practice plots

during *Rabi* 2019-20, 2020-21 and 2021-22, respectively. The per cent increase in yield in technology assessment plot when compared with the farmers practiced plot was 11.81, 10.64 and 16.22, respectively (Table 2). Subsequently, higher net returns were recorded in the technology assessment plots with Rs.91,000/-, Rs.89,000 and Rs.96,320/ha and lower net returns of Rs.76,675/-, Rs.77,500 and Rs. 82,880/-, respectively were recorded in the farmers practice plots during 2019-20, 2020-21 and 2021-22, respectively.

The benefit cost ratio (BC Ratio) was higher in technology assessment plots with 2.78, 2.63 and 2.75 in *Rabi* 2019-20, 2020-21 and 2021-22, respectively whereas benefit cost ratio was lower comparatively in farmers practiced plots with 2.43, 2.21 and 2.25 (Table 3).




Fig. 1. Results on incidence of fall armyworm on maize

Table 1. Difference between technological intervention through OFT in maize

Technology intervention	Farmers practice	Gap
Avoiding staggered sowing of maize	Not Followed	Full Gap
Installation of pheromone traps @ 8-10 per acre	No Pheromone traps used	Full Gap
Clean cultivation, balanced application of fertilizers, erection of bird perches @10/acre,	Not Followed	Full Gap
Spraying of Azadiractin (1500ppm)	Not Followed	Full Gap
Application of Emamectin benzoate @ 0.5g and <i>Metarrhizium anisopliae</i> @ 5g/l of water	Indiscriminate use of insecticides	Partial Gap

Table 2. Economic impact of experiment

Year	Yield (Q/ha)		% Increase in yield	Net returns Rs. /ha		B:C Ratio	
	Technology assessment	Farmers practiced		Technology assessment	Farmers practiced	Technology assessment	Farmers practiced
2019-20	71.0	63.5	11.81	91,000	76,675	2.78:1	2.43:1
2020-21	65.0	58.75	10.63	89,000	77,500	2.63:1	2.21:1
2021-22	68.8	59.2	12.9	96320	82880	2.75:1	2.25:1



Fig. 2. OFT on management against fall army worm

Table 3. Grain yield and gap analysis grain yield and gap analysis

Year	Yield gap (%)	Extension gap (Q/ha)	Technology gap (Q/ha)	Technology index (%)	Additional returns (Rs.)
2019-20	11.81	7.5	4	5.3	14,325
2020-21	10.64	6.25	10	13.3	11,500
2021-22	16.22	9.6	6.2	8.3	13,440

The results are in concurrence with the previous publication where, Rajashekhar et al. [9] observed FAW incidence before and after application of chemicals, showed Azadiractin 1500 ppm acted as the best oviposition deterrent, due to which most of the eggs failed to hatch and after whorl application of Emamectin benzoate @ 0.5 g/lit the incidence FAW was reduced up to 47-63 % compared to farmer practice which is similar to the present study. Dhaka et al. [10], Mistry et al. [11] and Bhati et al. [12] where maize yields were improved along with net returns. A similar study was reported in tribal areas of Andhra Pradesh where maize yields were increased from 62.1 q to 85.3 q/ha [13]. Reddy et al [14] reported that benefit cost ratio was significantly higher in the recommended technology (2.51) compared to farmers practice (2.12) which is similar to the current experiment.

4. CONCLUSION

The fall armyworm (*Spodoptera frugiperda*) has been one of the most devastating pests worldwide [15], capable of causing a 100% yield loss due to its unforeseen occurrence from the

seedling to the cob formation stage [16]. Based on the results obtained in the present study it can be concluded that the yield gap between farmer practice and technology assessment plot was perceptibly higher. There is urgent need to make stronger extension services for educating the cultivators in the implementation of improved technology. However, the yield level under local practice was lower which could be further improved by adopting recommended integrated pest management technologies. The OFT intervention is highly effective among maize farmers which increased their net returns. With the results obtained in technology assessment plot it is recommended to take further into wide demonstration which may record the same trend of positive results in Bhadradi Kothagudem district.

ACKNOWLEDGEMENT

Authors are sincerely grateful to thank to ICAR-ATARI & PJTSAU, Rajendranagar, Hyderabad for the financial support. We also thank innovative maize farmers of the Bhadradi Kothagudem district, Telangana for their help in the field layout, experiment planning, data

collection and smooth execution of the investigation in three consecutive years.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Goergen G, Kumar PL, Sankung SB, Togola A, Tamò M. First report of outbreaks of the fall armyworm *Spodoptera frugiperda* (J E Smith) (Lepidoptera, Noctuidae), a new alien invasive pest in West and Central Africa. PLoS One. 2016;11:e0165632.
2. Rwomushana I, Bateman M, Beale T, Beseh P, Cameron K, Chiluba M, Clotley V, Davis T, Day R, Early R, Godwin J, Gonzalez-Moreno P, Kansime M, Kenis M, Makale F, Mugambi I, Murphy S, Nunda W, Phiri N, Pratt C, Tambo J. Fall armyworm: Impacts and implication for Africa. Evidence Note Update. CAB International; 2018.
3. ICAR–National Bureau of Agricultural Insect Resources Newsletter. 2018;X(4).
4. Rakshit S, Chikkappa GK. Perspective of maize scenario in India: way forward. Maize J. 2018;7(2):49–55.
5. Sparks AN. A review of the biology of the fall armyworm. Florida Entomologist. 1979;62:82-87.
6. Choudhary BN. Krishi Vigyan Kendra-A guide for KVK managers. Publication, Division of Agricultural Extension, ICAR. 1999:73-78.
7. Rajashekhar M, Prabhakar Reddy T, Chandrashekara KM, Rajashekar B, Jagan Mohan Reddy M, Ramakrishna K et al. Evaluation of integrated pest management module for pink bollworm, *Pectinophora gossypiella* (Saunders) and its economic analysis under farmer's field conditions. International Journal of Pest Management. 2022;68(3):1-9.
8. Samui SK, Maitra S, Roy DK, Mondal AK, Saha D. Evaluation of front line demonstration on groundnut (*Arachis hypoggaea*L.) in Sundarbans. Journal of the Indian Society of Coastal Agricultural Research. 2000;18(2):180-183.
9. Rajashekhar M, Prabhakar RT, Rajashekar B, Jagan MRM, Ramakrishna K, Shankar A, Afifa J. Evaluation of IPM modules against fall armyworm management on maize entomology innovation and entrepreneurship; 2022.
10. Dhaka BL, Meena BS, Suwalka RL. Popularization of improved maize production technology through frontline demonstrations in South-Eastern Rajasthan. J Agric Sci. 2010;1(1):39-42. DOI: 10.1080/09766898.2010.11884652
11. Mistry JJ, Viho KJ, Patel GJ. Impact of front line demonstrations on maize. Gujarat J Extension Educ. 2015;26(1):18-20.
12. Bhati BS, Soni RL, Rathore RS. Impact of front line demonstration on maize yield improvement in tribal belt of rajasthan. Trends Biosci. 2017;10(21): 4192-4.
13. Swaroopa VJ, Mounica D, Pavani U, Sree D. Popularization of maize production technology through front line demonstration in tribal areas of East Godavari. J Krishi Vigyan. 2016;4(2):80-2.
14. Reddy BKK, Jyothi VS, Sadhineni M, Johnson M, Swamy GN. Evaluation of IPM modules against fall armyworm in Maize through frontline demonstration and its economic impact. International Journal of Environment and Climate Change. 2023; 13(1):24-29.
15. Sagar GC, Aastha B, Laxman K.. An introduction of fall armyworm (*Spodoptera frugiperda*) with management strategies: A review paper. Nippon Journal of Environmental Science; 1(4):1010 Available:https://doi.org/10.46266/njes.1010
16. Kumar RM, Gadratagi BG, Paramesh V, Kumar P, Madivalar Y, Narayanappa N, Ullah F. Sustainable management of invasive fall army worm, *Spodoptera frugiperda*. Agronomy. 2022;12(9):2150. Available:https://doi.org/10.3390/agronom y12092150

© 2023 Narayanamma et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/102204>