



# **Assessment of Heavy Metal Concentrations in Pumpkin Grown along River Benue, Makurdi Local Government Area of Benue State, Nigeria**

**O. A. Onekutu<sup>1</sup>, H. O. A. Oluma<sup>1</sup> and G. Dajoh<sup>1\*</sup>**

<sup>1</sup>*Department of Biological Sciences, University of Agriculture, Makurdi, Nigeria.*

## **Authors' contributions**

*This work was carried out in collaboration between all authors. Author GD designed the study, performed laboratory analysis, the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors OAO and HOAO supervised and provided the guide managed the analyses and approved the study and also managed the literature searches. All authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.9734/PSIJ/2018/39036

### Editor(s):

- (1) Alexander Tolstoguzov, Department of Physics, Universidade Nova de Lisboa, Portugal.  
(2) Chao-Qiang Geng, Professor, National Center for Theoretical Science, National Tsing Hua University Hsinchu, Taiwan.  
(3) Thomas F. George, Chancellor/Professor Department of Chemistry and Physics, University of Missouri-St. Louis, Boulevard St. Louis, USA.

### Reviewers:

- (1) Falusi Bamidele Ayodeji, Federal College of Education, Nigeria.  
(2) Şana Sungur, Mustafa Kemal University, Turkey.  
Complete Peer review History: <http://www.sciencedomain.org/review-history/24138>

**Original Research Article**

**Received 11<sup>th</sup> November 2017**  
**Accepted 20<sup>th</sup> January 2018**  
**Published 13<sup>th</sup> April 2018**

## **ABSTRACT**

This study aimed at detecting and assessing the concentrations of heavy metals in Pumpkin irrigated using surface and groundwater sources grown along the banks of River Benue, Makurdi, Nigeria. Eight heavy metals were detected to include cadmium, chromium, cobalt, copper, iron, manganese, lead and zinc. The mean concentration of heavy metals in pumpkin irrigated with surface water obtained from Fiidi, Wurukum, new garage and Wadata areas of Makurdi was determined using triplicate values obtained from heavy metal analysis. The results show the mean concentrations of heavy metals to include Cd ( $2.07 \pm 0.1$ ), Cr ( $2.07 \pm 0.1$ ), Co ( $0.803 \pm 0.01$ ), Cu ( $3.14 \pm 0.1$ ), Mn ( $2.51 \pm 0.02$ ), Pb ( $1.05 \pm 0.05$ ) and Zn ( $9.03 \pm 0.03$ ). Fiidi had Cd ( $1.02 \pm 0.03$ ), Cr ( $2.24 \pm 0.04$ ), Co ( $0.417 \pm 0.02$ ), Cu ( $2.61 \pm 0.01$ ), Fe ( $1.03 \pm 0.02$ ), Pb ( $1.22 \pm 0.02$ ) and Zn ( $6.81 \pm 0.001$ ). Wurukum had mean concentration of Cd Cr, Co, Cu, Fe, Mn, Pb, and Zn to be 0.920  $\pm$

\*Corresponding author: E-mail: [ashinya78@gmail.com](mailto:ashinya78@gmail.com);

0.03,  $1.61 \pm 0.02$ ,  $0.410 \pm 0.02$ ,  $2.42 \pm 0.02$ ,  $1.02 \pm 0.03$ ,  $1.22 \pm 0.02$ ,  $1.41 \pm 0.01$  and  $8.54 \pm 0.04$  for the heavy metals respectively. On the other hand, pumpkin irrigated with surface water around New garage had a mean concentration of  $0.710 \pm 0.03$ ,  $2.13 \pm 0.03$ ,  $0.410 \pm 0.01$ ,  $2.90 \pm 0.01$ ,  $1.11 \pm 0.02$ ,  $2.03 \pm 0.06$ ,  $0.527 \pm 0.03$ ,  $3.11 \pm 0.01$ ,  $1.22 \pm 0.07$ ,  $1.81 \pm 0.02$ ,  $1.22 \pm 0.03$  and  $6.83 \pm 0.06$  for Cd, Cr, Co, Cu, Fe, Mn, Pb and Zinc respectively. The mean concentrations of heavy metals in pumpkin irrigated with groundwater were Manganese was highest in Zua ( $4.64 \pm 0.03$ ) and cobalt ( $0.410 \pm 0.02$ ) was lowest, Fiidi had Zinc ( $9.01 \pm 0.01$ ) as the highest whereas cobalt had  $0.517 \pm 0.02$  as the lowest mean concentration. The new garage had  $7.85 \pm 0.06$  and  $0.517 \pm 0.02$  whereas Wadata had  $8.24 \pm 0.2$  and  $0.4 \pm 0.02$  for zinc and cobalt as the highest and lowest mean concentration respectively. The presence of the heavy metals in pumpkin planted in the locations studied did not vary significantly since all the heavy metals detected were present in all the samples studied at  $P > 0.05$ .

*Keywords: Heavy metals; pumpkin; irrigation.*

## 1. INTRODUCTION

Irrigation is the controlled application of water for agricultural purposes through man-made systems to supply water requirements not satisfied by rainfall. Crop irrigation is vital throughout the world in order to provide the world's ever-growing populations with enough food. An irrigation activity that is gradually gaining recognition is the one being practiced under the urban and peri-urban agriculture (UPA). The system involves the use of stream water to irrigate lands at the banks with the objective of producing fruits and vegetables for the consumption of city dwellers [1].

The heavy metals pollution of Rivers effects on the irrigation water quality along the river, as well as crops irrigated with it. The irrigated crops may take-up these heavy metals and consequently, introduce them into food chain, resulting to gradual accumulation in the humans, and thereafter, present health hazard.

Studies of [2] and [3] have shown that intake of trace metals from dietary sources may represent a significant exposure pathway for human populations. However, dietary exposure to trace metals is highly variable. For example, [4] has observed that for Cd, the principal exposure route for the general population is through uptake by food plants. Where metal concentrations in crops exceed the limits, it may be possible to use this produce in animal feeds in order to minimize the effect upon the human diet. However, animals fed on a metal-enriched diet may have elevated concentrations of these metals in their tissues and milk. Reilly [3] noted that regular consumption of metal-enriched animal products may lead to adverse health effects in humans.

Furthermore, [5] have observed that the greatest degree of metal accumulation occurs in offal, such as livers and kidneys.

Heavy metals such as Cd, Zn, Pb, Cr, Fe and Cu are commonly found on contaminated sites [6]. Sources of heavy metal contamination to the environment have been identified by various researchers to include; exhaust pipes, wastewater effluents and oil exploration activities. The implication of gas explosive on human health is all related to the exposure of those hazardous air pollutant emitted during incomplete combustion, acid rain effect and subsequent acidification of the soil [7]. Bioaccumulation of these heavy metals has been identified mainly in surface water, soil and in plants [8].

The supply of clean and uncontaminated water is a great challenge facing developing nations. In Nigeria, pollution is a major threat to both surface and underground water bodies [6,9]. Green vegetables such as pumpkin leaf form a substantial portion of the daily human diet of the southern part of Nigeria. These are grown on all types of available land such as banks of polluted rivers, alongside road and areas enclosing wastewater ponds. These plants absorb some of these chemicals which are toxic to human health [10] Industrial emissions of contaminant to the atmosphere which is finally deposited on soil may cause some pollution problem. The contaminant concentration in soil mainly depends on the adsorption properties of soil matter [11]. The high toxic and persistent nature of heavy metals in the environment has made them priority pollutants. The present study quantified the heavy metals; cadmium, chromium, cobalt, copper, iron, lead manganese and zinc (Cd, Cr, Co, Cu, Fe, Pb, Mn, Zn) in pumpkin

irrigated with ground and surface water from different locations in Makurdi the Benue State capital of Nigeria.

## 2. METHODOLOGY

### 2.1 Sample

#### 2.1.1 Pretreatment of samples for analysis of metals in vegetable samples

The plant Samples were thoroughly washed rinsed with tap water and then with double distilled water to remove any attached soil particles. It was then be cut into small portions and placed in a large crucible where they will be oven dried at 60°C overnight. The dried plant was then be grounded into fine particles using a clean mortar and pestle.

The triacid method of digestion was employed. The acids used were Nitric acid (HNO<sub>3</sub>), Perchloric acid (HClO<sub>4</sub>) and concentrated Sulphuric acid (H<sub>2</sub>SO<sub>4</sub>) in the ratio of 68:8:2 respectively.

0.2 g of the powdered crop sample was weighed into a 100 ml glass beaker. 30 ml of the acid mix was added and the content swirled and placed on a hot plate. The beaker was heated until the brown fumes of the Nitric acid go off, and the heating continued until the content of the beaker is reduced to about 5 ml. The content was allowed to cool and little amount of distilled water were added and the beaker swirled again. The content was then be poured into a 50 ml volumetric flask and made up to the required mark. The digest obtained was used for the determination of the metals [12].

#### 2.1.2 Determination of heavy metals

The measurement of heavy metal concentration was carried out with an Atomic Absorption Spectrophotometer (AAS). All concentrations were determined using the absorbance made air-acetylene flame. Eight working solutions were prepared from the stock solution for each of the metals (Cr, Cu, Cd, Zn, Co, Fe, Pb, and Mn) by successive serial dilution and each of the standard solutions were aspirated into the flame of AAS and the absorbance recorded in each case. A plot of the concentration against the corresponding absorbance gave the calibration

curve of each of the metals. The samples were aspirated into the flame and the absorbance obtained [13].


## 3. RESULTS

Fig. 1 shows the occurrence of heavy metals in pumpkin plants for irrigated with surface water obtained from Fiidi, Wurukum, New garage and Wadata areas of Makurdi. The result show the mean concentrations of heavy metals to include (in mg/kg) Cd (2.07 ± 0.1), Cr (2.07 ± 0.1), Co (0.803 ± 0.01), Cu (3.14 ± 0.1), Mn (2.51 ± 0.02), Pb (1.05 ± 0.05) and Zn (9.03 ± 0.03). Fiidi had Cd (1.02 ± 0.03), Cr (2.24 ± 0.04), Co (0.417 ± 0.02), Cu (2.61 ± 0.01), Fe (1.03 ± 0.02), Pb (1.22 ± 0.02) and Zn (6.81 ± 0.001). wurukum had mean concentrations of Cd Cr, Co, Cu, Fe, Mn, Pb, and Zn to be 0.920 ± 0.03, 1.61 ± 0.02, 0.410 ± 0.02, 2.42 ± 0.02, 1.02 ± 0.03, 1.22 ± 0.02, 1.41 ± 0.01 and 8.54 ± 0.04 for the heavy metals respectively. On the other hand, pumpkin irrigated with surface water around New garage had a mean concentration of 0.710 ± 0.03, 2.13 ± 0.03, 0.410 ± 0.01, 2.90 ± 0.01, 1.11 ± 0.02, 2.03 ± 0.06, 0.527 ± 0.03, 3.11 ± 0.01, 1.22 ± 0.07, 1.81 ± 0.02, 1.22 ± 0.03 and 6.83 ± 0.06 for Cd, Cr, Co, Cu, Fe, Mn, Pb and Zinc respectively.

Fig. 2 shows the mean concentration of heavy metals in pumpkin irrigated using ground water in the locations studied. The mean concentration of manganese was highest in Zua (4.64 ± 0.03) and cobalt (0.410 ± 0.02), Fiidi had Zinc (9.01 ± 0.01) and cobalt 0.517 ± 0.02 as the heavy metals with highest and convert mean concentration respectively. New garage had 7.85 ± 0.06 and 0.517 ± 0.02 whereas Wadata had 8.24 ± 0.2 and 0.4 ± 0.02 for zinc and cobalt respectively.


## 4. DISCUSSION

Anthropogenic activities, land use, lithogenic contribution by surface runoff erosion, deposition of eroded materials, temperature etc can artificially change organic matter content as well as a heavy metal concentration in the soil. The heavy metals studied in pumpkin with the exception of chromium had values higher than FAO/WHO recommended permissible levels (Pb 0.3, Cd 0.2, Fe 425, Zn 99.40, Cu 73.0, Cr 0.05, Mn 1.0 (in mg/kg).



**Fig. 1. Heavy metal concentration in pumpkin irrigated with surface water**

Key: PZS= Pumpkin crop-Zua-surface water irrigated  
 PFS= Pumpkin crop-Fiidi-Surface water irrigated  
 PWS= Pumpkin crop-Wurukum-surface water irrigated  
 PNS=Pumpkin crop-new garage-Surface water irrigated  
 PWAS= Pumpkin crop-Wadata-surface water irrigated



**Fig. 2. Heavy metal concentration in pumpkin irrigated with ground water**

**Table 1. Concentration of Heavy metals in pumpkin irrigated with ground water in Wadata (in mg/kg)**

	Cd	Cr	Co	Cu	Fe	Mn	Pb	Zn
PWAG1	0.800	2.00	0.400	2.00	1.20	2.80	0.900	8.30
PWAG2	0.800	2.20	0.400	2.00	1.23	2.82	0.910	8.43
PWAG3	0.830	2.30	0.430	2.20	1.20	2.83	0.920	8.00
Mean	0.810	2.17	0.41	2.07	1.21	2.82	0.91	8.24
SD	0.02	0.2	0.02	0.1	0.02	0.02	0.01	0.2
RSD	2.1	7.1	4.2	5.6	1.4	0.5	1.1	2.7

**Table 2. Concentration of heavy metals in pumpkin irrigated with ground water in new Garrage (mg / kg)**

	<b>Cd</b>	<b>Cr</b>	<b>Co</b>	<b>Cu</b>	<b>Fe</b>	<b>Mn</b>	<b>Pb</b>	<b>Zn</b>
PNG1	1.00	2.20	0.500	1.90	1.40	3.20	1.30	7.80
PNG2	1.00	2.20	0.520	1.90	1.43	3.20	1.04	7.91
PNG3	1.08	2.30	0.530	1.86	1.45	3.32	1.46	7.83
Mean	1.03	2.23	0.517	1.89	1.43	3.24	1.27	7.85
SD	0.05	0.06	0.02	0.02	0.03	0.07	0.21	0.06
RSD	4.5	2.6	3.0	1.2	1.8	2.1	16.7	0.7

**Table 3. Concentration of heavy metals in pumpkin irrigated with ground water in Fiidi (mg/kg)**

	<b>Cd</b>	<b>Cr</b>	<b>Co</b>	<b>Cu</b>	<b>Fe</b>	<b>Mn</b>	<b>Pb</b>	<b>Zn</b>
PFG1	0.800	2.00	0.400	2.20	1.50	3.50	1.70	7.20
PFG2	0.820	2.00	0.430	2.20	1.50	3.56	1.72	7.40
PFG3	0.810	2.08	0.440	2.23	1.50	3.54	1.73	7.32
Mean	0.810	2.03	0.423	2.21	1.50	3.53	1.72	7.31
SD	0.01	0.05	0.02	0.02	0.0	0.03	0.02	0.1
RSD	1.2	2.3	4.9	0.8	0.0	0.9	0.9	1.4

**Table 4. Concentration of heavy metals in pumpkin irrigated with ground water in Zua (in mg/kg)**

	<b>Cd</b>	<b>Cr</b>	<b>Co</b>	<b>Cu</b>	<b>Fe</b>	<b>Mn</b>	<b>Pb</b>	<b>Zn</b>
PZG1	1.80	1.80	0.400	2.30	1.50	4.60	1.50	2.30
PZG2	1.82	1.80	0.400	2.32	1.53	4.65	1.53	2.31
PZG3	1.80	1.83	0.430	2.30	1.54	4.66	1.55	2.33
Mean	1.81	1.81	0.410	2.31	1.52	4.64	1.53	2.31
SD	0.01	0.02	0.02	0.01	0.02	0.03	0.03	0.02
RSD	0.6	1.0	4.2	0.5	1.4	0.7	1.6	0.7

**Table 5. Concentration of heavy metals in pumpkin irrigated with ground water in Wurukum (mg/kg)**

	<b>Cd</b>	<b>Cr</b>	<b>Co</b>	<b>Cu</b>	<b>Fe</b>	<b>Mn</b>	<b>Pb</b>	<b>Zn</b>
PWG1	1.20	2.20	0.500	2.10	1.40	3.00	1.10	9.00
PWG2	1.23	2.24	0.520	2.13	1.41	3.09	1.10	9.00
PWG3	1.20	2.21	0.530	2.10	1.44	3.07	1.00	9.02
Mean	1.21	2.22	0.517	2.11	1.42	3.05	1.07	9.01
SD	0.02	0.02	0.02	0.02	0.02	0.05	0.06	0.01
RSD	1.4	0.9	3.0	0.8	1.5	1.5	5.4	0.1

Key: PZG= Pumpkin crop-Zua- ground water irrigated  
 PFG= Pumpkin crop-Fiidi- ground water irrigated  
 PWG= Pumpkin crop-Wurukum- ground water irrigated  
 PNG= Pumpkin crop-new garage-ground water irrigated  
 PWAG= Pumpkin crop- Wadata- ground water irrigated

The bioaccumulation of these metals into the pumpkin leaf may be due to non - formation of complexes with the organic matters in the soil thereby making it possible for these metals to leach from the soil into the plant. [11,14]. Akhionbare and Akhionbare, [15] reported that increase in total zinc content of the soil could lower fluted pumpkin lead accumulation. [16] also reported that the soil, root and stem play a role in

the process of heavy metal transport. These the level of Pb, Cr and Zn reported in the pumpkin leaf from these locations were in agreement with those reported in the study of [16] for pumpkin and Amaranths leaf. Comparing the average heavy metal contents obtained in vegetable to the maximum levels permitted by FAO/WHO, we observed that people eating vegetable grown at these locations, were consuming it at

concentration levels that are potentially hazardous to their health.

The results of the present study showed high concentration levels of the heavy metals in pumpkin samples irrigated using ground water appreciably above FAO threshold limit though with the exclusion of lead and iron which was below the threshold. The high concentration of the heavy metals in the crops reflected the concentration in ground water source used for irrigation [17] and Connor [18] opined that heavy metals can be transferred from soil to other components of the ecosystem such as underground water or crops. Studies of [18] observed normal levels of Fe, Cu, Zn, Mn, Ni and Cd in edible portions of vegetable crops irrigated groundwater sources with the exception of Cr and Pb which exceeded the FAO/ WHO limits of 0.5 mg/kg. High accumulation of Pb in vegetables irrigated with heavy metal contaminated groundwater sources was previously reported in Nebal [19].

The results of analysis of heavy metals in pumpkin, spinach and tomato samples irrigated using surface water appreciably above FAO threshold limit though with the exclusion of lead and iron which were below the threshold. Previous research findings of [20] in four wild plants namely *Xanthium strumarium*, *Acacia modesta*, *Dodmea viscosa* and *Tamarix aphyda* showed that cadmium, zinc, iron, copper, nickel, chromium and lead were present. Plants samples were analyzed separately for their root, stem and leaves. Flame absorption spectrometer was used for analyzing the samples and concentrations of cadmium, chromium, iron and lead in water were recorded above the permissible limits set by WHO while zinc and copper were recorded below the permissible limits and no concentration of nickel was recorded in water samples.

Chiroma et al. [13] in their study on heavy metal contamination of vegetables and soils irrigated with sewage water in Yola, Nigeria reported high concentration of heavy metals (especially Fe, Zn, Cu, Mg, Mn and Pb) contamination in the soils irrigated with sewage water. Their study also found out that there is accumulation of these metals in different parts of plant cultivated on the soil. The study also shows that the heavy metal concentration vary in different parts of the plant with Fe shown to accumulate in roots and leaves while Zn accumulates in roots and translocates gradually to the leaves and Mn and Mg show

greater accumulations in unwashed leaves. [21] carried out a study in Rajasthan India to assess the level of different heavy metals such as iron, manganese, copper and zinc, in vegetables irrigated with water from different sources. The results indicated a substantial build-up of heavy metals in vegetables irrigated with wastewater.

The heavy metals detected in both surface and ground water sources were cadmium (Cd), Chromium (Cr), cobalt (Co), Copper (Cu), iron (Fe), Manganese (Mn), lead (Pb) and Zinc (Zn). The results revealed high level concentration of the detected heavy metals in both sources above FAO limits for irrigation water except for Lead and Iron. Previous studies of [22] revealed the presence of cadmium (Cd), chromium (Cr), Copper (Cu), lead (Pb) Nickel (Ni) and zinc (zn) in ground water used for irrigation in Tabriz city in Iran. The average and even maximum concentrations of heavy metals in ground water was reportedly less than the toxicity threshold limits of agricultural water, thus signifying the effectiveness of the efforts of authorities in reducing the rate and level of heavy metal contamination, hence posed no health risk as compared to the findings of our study.

As an interim measure, the source of pumpkin in the affected areas studied should be regularly checked so as to avoid food and vegetables irrigated with polluted water since the linkage between crops, heavy metals content and heavy metals in irrigation water has been established in previous studies.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Binns JA, Maconachie RA, Tanko AI. Water, land and health in urban and peri urban food production: The case of Kano, Nigeria; 2003. Available:[www.cityfarmer.org/WaterLandHealthKano](http://www.cityfarmer.org/WaterLandHealthKano)
2. Baes CF, Sharp RD, Sjoreen AL, Shor RW. Review and analysis of parameters for assessing transport of environmentally released radionuclides through agriculture. US. Department of Commerce, National

- Technical Information Service, Virginia; 1991.
3. Reilly C. Metal contamination of food. (Second edition. Elsevier Applied Science London and New York); 1991.
  4. Lopez - Artiguez M, Soria ML, Camean A, Repetto M. Cadmium in the diet of the local population of Seville (Spain). *Bulletin of Environmental Contamination and Toxicology*. 1993;50:417-424.
  5. Beresford NA, Mayes RW, Crout NMJ, Maceachern PJ, Dodd BA, Bernett CL, Lamb CS. Transfer of cadmium and mercury to sheep tissues. *Environmental Science and Technology*. 1999;33:2395-2402.
  6. Oathman OC. Heavy metal in green vegetables and soils from vegetable gardens. *Tanzanian Journal of Science*. 27;37-48. Press Inc., Boca Raton, Florida. 2001;80(83):104-106.
  7. Akhionbare SMO. Factors in migration of heavy metals in the Otamiri river system. *International Journal of Science and Nature*. 2011;2(4):856-860.
  8. Akoto O, Bruce TN, Darko G. Heavy metals pollution profiles in stream serving the Owabi reservoir. *African Journal of Environmental Science and Technology*. 2008;2(2):354-359.
  9. Osabolien E, Otutu OY, Oluya BO. Concentration of heavy metals in soil, tubers and leaves of cassava plants grown around some oil spill and gas flaring Zones in delta state, Nigeria. *Applied Science Research Journal*. 2013;1(1):1-6.
  10. Baranowska KS, Wlochowicz A, Szczepanik K. Determination of heavy metal contents in sample of medicinal herbs. *Polish Journal of Environmental Studies*. 2002;11(5):467-471.
  11. Deka J, Sarma HP. Heavy metal contamination in soil in an industrial zone and its relation with some soil properties. *Archives of Applied Science Research*. 2012;4(2):831-836.
  12. Salawu K, Barau MM, Mohammed D, Mikailu DA, Abdullahi BH, Uroko RI. Determination of some selected heavy metals in spinach and irrigated water from Samaru Area within Gusau Metropolis in Zamfara State, Nigeria. *Journal of Toxicology and Environmental Health Sciences*. 2015;7(8)76-80. DOI:0.5897/JTEHS2015.0339 Article Number: 2E6F3D355501
  13. Chiroma TM, Hymore, FK, Ebawele RO. Heavy metal contamination of vegetables and soils irrigated with sewage water in Yola. *NJER*. 2003;2(3):25-31. 55:251-260.
  14. Kabata-Pendias A, Henryk P. Trace elements in soils and plants CRC; 1984.
  15. Akhionbare SMO, Akhionbare WN. Effect of impoundment on the temperature oxygen regime of 1 kpoba River, Nigeria, *International Research Journal In Engineering Science and Technology (IREJEST)*. 2004;1(1):46-58.
  16. Oshman OC. Heavy metals in green vegetables and soils from vegetable gardens in Dar Es Salaam, Tanzania. *Tanzania Journal of Science*. 2001;27:37-48.
  17. Kumar NJI, Soni H, Kumar RNJ. Characterization of heavy metals in vegetables using inductive coupled plasma analyzer (ICPA). *Appl. Sci. Environ. Manage*. 2007;11:75.
  18. Conor R. Metal contamination of food: Its significance for food quality and human health: 3rd Edition. Black-well Science Publishing. 2002;222.
  19. AL-Jaboobi Muamar, Abdelmajid Zouahri, M'hamed Tijane, Abdellah El Housni, Zakaria Mennane, Hasna Yachou, Mohammed Bouksaim. Assessment of the impact of wastewater use on soil properties. *J. Mater. Environ. Sci*. 2014; 5(3):961-966.
  20. Sharma RK, Agrawal M, Marshall FM. Heavy metals contamination of soil and vegetables in suburban areas of Varanasi, India. *Ecotoxicology and Environmental Safety*. 2007;66:258-266.
  21. Arora M, Kiranb B, Rania S, Rania A, Kaura B, Mittala N. Heavy metal accumulation in vegetables irrigated with water from different sources. *Food Chemist. J*. 2008;111:811-815.
  22. Taghipour H, Mosaferi M, Armanfar F, Gaemmagam SJ. Heavy metals pollution in the soils of suburban areas in big cities: A case study; 2013.

**APPENDICES**

	<b>Cd</b>	<b>Cr</b>	<b>Co</b>	<b>Cu</b>	<b>Fe</b>	<b>Mn</b>	<b>Pb</b>	<b>Zn</b>
PZS1	2.00	2.00	0.800	3.10	1.00	2.50	1.00	9.00
PZS2	2.02	2.12	0.810	3.20	1.11	2.50	1.09	9.05
PZS3	2.20	2.10	0.800	3.13	1.10	2.54	1.07	9.03
Mean	2.07	2.07	0.803	3.14	1.07	2.51	1.05	9.03
SD	0.1	0.1	0.01	0.1	0.1	0.02	0.05	0.03
RSD	5.3	3.1	0.7	1.6	5.7	0.9	4.5	0.3

	<b>Cd</b>	<b>Cr</b>	<b>Co</b>	<b>Cu</b>	<b>Fe</b>	<b>Mn</b>	<b>Pb</b>	<b>Zn</b>
PFS1	1.00	2.20	0.400	2.60	1.00	2.30	1.20	6.80
PFS2	1.06	2.27	0.430	2.62	1.00	2.34	1.23	6.82
PFS3	1.00	2.26	0.420	2.61	1.08	2.30	1.22	6.80
Mean	1.02	2.24	0.417	2.61	1.03	2.31	1.22	6.81
SD	0.03	0.04	0.02	0.01	0.05	0.02	0.02	0.01
RSD	3.4	1.7	3.7	0.4	4.5	1.0	1.3	0.2

	<b>Cd</b>	<b>Cr</b>	<b>Co</b>	<b>Cu</b>	<b>Fe</b>	<b>Mn</b>	<b>Pb</b>	<b>Zn</b>
PWS1	0.900	1.60	0.400	2.40	1.00	1.20	1.40	8.50
PWS2	0.900	1.63	0.400	2.42	1.00	1.21	1.42	8.57
PWS3	0.960	1.61	0.430	2.44	1.05	1.24	1.40	8.56
Mean	0.920	1.61	0.410	2.42	1.02	1.22	1.41	8.54
SD	0.03	0.02	0.02	0.02	0.03	0.02	0.01	0.04
RSD	3.8	0.9	4.2	0.8	2.8	1.7	0.8	0.4

	<b>Cd</b>	<b>Cr</b>	<b>Co</b>	<b>Cu</b>	<b>Fe</b>	<b>Mn</b>	<b>Pb</b>	<b>Zn</b>
PNS1	0.700	2.10	0.400	2.90	1.10	2.00	1.10	9.10
PNS2	0.690	2.13	0.420	2.91	1.10	2.00	1.10	9.08
PNS3	0.740	2.15	0.410	2.90	1.13	2.10	1.12	9.30
Mean	0.710	2.13	0.410	2.90	1.11	2.03	1.11	9.16
SD	0.03	0.03	0.01	0.01	0.02	0.06	0.01	0.1
RSD	3.7	1.2	2.4	0.2	1.6	2.8	1.0	1.3

	<b>Cd</b>	<b>Cr</b>	<b>Co</b>	<b>Cu</b>	<b>Fe</b>	<b>Mn</b>	<b>Pb</b>	<b>Zn</b>
PWAS1	0.800	2.00	0.500	3.10	1.20	1.80	1.20	6.90
PWAS2	0.860	2.10	0.550	3.12	1.30	1.84	1.20	6.79
PWAS3	0.810	2.12	0.530	3.10	1.16	1.80	1.26	6.80
Mean	0.823	2.07	0.527	3.11	1.22	1.81	1.22	6.83
SD	0.03	0.06	0.03	0.01	0.07	0.02	0.03	0.06
RSD	3.9	3.1	4.8	0.4	5.9	1.3	2.8	0.9

	<b>Cd</b>	<b>Cr</b>	<b>Co</b>	<b>Cu</b>	<b>Fe</b>	<b>Mn</b>	<b>Pb</b>	<b>Zn</b>
PZG1	1.80	1.80	0.400	2.30	1.50	4.60	1.50	2.30
PZG2	1.82	1.80	0.400	2.32	1.53	4.65	1.53	2.31
PZG3	1.80	1.83	0.430	2.30	1.54	4.66	1.55	2.33
Mean	1.81	1.81	0.410	2.31	1.52	4.64	1.53	2.31
SD	0.01	0.02	0.02	0.01	0.02	0.03	0.03	0.02
RSD	0.6	1.0	4.2	0.5	1.4	0.7	1.6	0.7

	<b>Cd</b>	<b>Cr</b>	<b>Co</b>	<b>Cu</b>	<b>Fe</b>	<b>Mn</b>	<b>Pb</b>	<b>Zn</b>
PFG1	0.800	2.00	0.400	2.20	1.50	3.50	1.70	7.20
PFG2	0.820	2.00	0.430	2.20	1.50	3.56	1.72	7.40
PFG3	0.810	2.08	0.440	2.23	1.50	3.54	1.73	7.32
Mean	0.810	2.03	0.423	2.21	1.50	3.53	1.72	7.31
SD	0.01	0.05	0.02	0.02	0.0	0.03	0.02	0.1
RSD	1.2	2.3	4.9	0.8	0.0	0.9	0.9	1.4



	<b>Cd</b>	<b>Cr</b>	<b>Co</b>	<b>Cu</b>	<b>Fe</b>	<b>Mn</b>	<b>Pb</b>	<b>Zn</b>
PWG1	1.20	2.20	0.500	2.10	1.40	3.00	1.10	9.00
PWG2	1.23	2.24	0.520	2.13	1.41	3.09	1.10	9.00
PWG3	1.20	2.21	0.530	2.10	1.44	3.07	1.00	9.02
Mean	1.21	2.22	0.517	2.11	1.42	3.05	1.07	9.01
SD	0.02	0.02	0.02	0.02	0.02	0.05	0.06	0.01
RSD	1.4	0.9	3.0	0.8	1.5	1.5	5.4	0.1

	<b>Cd</b>	<b>Cr</b>	<b>Co</b>	<b>Cu</b>	<b>Fe</b>	<b>Mn</b>	<b>Pb</b>	<b>Zn</b>
PNG1	1.00	2.20	0.500	1.90	1.40	3.20	1.30	7.80
PNG2	1.00	2.20	0.520	1.90	1.43	3.20	1.04	7.91
PNG3	1.08	2.30	0.530	1.86	1.45	3.32	1.46	7.83
Mean	1.03	2.23	0.517	1.89	1.43	3.24	1.27	7.85
SD	0.05	0.06	0.02	0.02	0.03	0.07	0.21	0.06
RSD	4.5	2.6	3.0	1.2	1.8	2.1	16.7	0.7

	<b>Cd</b>	<b>Cr</b>	<b>Co</b>	<b>Cu</b>	<b>Fe</b>	<b>Mn</b>	<b>Pb</b>	<b>Zn</b>
PWAG1	0.800	2.00	0.400	2.00	1.20	2.80	0.900	8.30
PWAG2	0.800	2.20	0.400	2.00	1.23	2.82	0.910	8.43
PWAG3	0.830	2.30	0.430	2.20	1.20	2.83	0.920	8.00
Mean	0.810	2.17	0.41	2.07	1.21	2.82	0.91	8.24
SD	0.02	0.2	0.02	0.1	0.02	0.02	0.01	0.2
RSD	2.1	7.1	4.2	5.6	1.4	0.5	1.1	2.7

© 2018 Onekutu et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
 The peer review history for this paper can be accessed here:  
<http://www.sciencedomain.org/review-history/24138>