



Antimicrobial Susceptibility Profile of Coliforms from Bovine Mastitis Cases among Pastoral Herds in Parts of Kaduna State, Nigeria: Curbing the Environmental Health Risk

D. Makolo^{1,2*}, A. B. Suleiman¹, O. S. Olonitola¹, M. Bello³, M. I. Alfa⁴, I. Ahmadu¹ and F. O. Awulu^{1,5}

¹*Department of Microbiology, Faculty of Life Sciences, Ahmadu Bello University, Zaria, Nigeria.*

²*School of Preliminary Studies, Kogi State Polytechnic, Lokoja, Nigeria.*

³*Department of Veterinary Public Health and Preventive Medicine, Faculty of Veterinary Medicine, Ahmadu Bello University, Zaria, Nigeria.*

⁴*Department of Civil Engineering, Faculty of Engineering, University of Jos, Nigeria.*

⁵*Department of Science Laboratory Technology, Kogi State Polytechnic, Lokoja, Nigeria.*

Authors' contributions

This work was carried out in collaboration between all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJARR/2019/v3i230084

Editor(s):

(1) Dr. Muge K. Davran, Associate Professor, Department of Agricultural Economics, Faculty of Agriculture, University of Cukurova, Turkey.

Reviewers:

(1) Michael Hässig, University of Zurich, Switzerland.

(2) A. Salama Ahmed Osman, Kafrelsheikh University, Egypt.

(3) Odetoyin Babatunde, Obafemi Awolowo University, Nigeria.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/46287>

Original Research Article

Received 29 October 2018

Accepted 17 January 2019

Published 15 February 2019

ABSTRACT

Consumption of raw milk from pastoral bovines have been identified as a major source of public and environmental health risk in developing countries. Antimicrobial resistance is a global health challenge threatening the lives of humans and animals. The indiscriminate use of antimicrobial agents among the pastoralists on commercial animals, especially for non-therapeutic purposes has been linked to the development of resistant strains of potentially pathogenic bacteria which are being transferred from animals to humans. This study investigated the antimicrobial susceptibility profile of coliform bacteria isolated from mastitis milk of pastoral herds. Out of 147 milk samples

*Corresponding author: Email: makolodaniel@gmail.com;

collected and screened for subclinical mastitis, 29 (19.7%) were positive. Out of the 29 mastitis positive samples, 13 (8.8%) were positive for coliforms (6 *E. coli* and 7 *K. pneumoniae*). All the coliform isolates showed 100% resistance to Penicillin and Tetracycline, and were all 100% susceptible to Imipenem. High multidrug resistance was expressed by all the isolates to Penicillin, Tetracycline and Erythromycin. All the isolates (100%) had Multiple Antibiotic Resistance Index (MARI) of 0.2 and above which is an indication of gross abuse of antibiotics within the studied population. However, antibiotics still effective against the coliform species tested were Imipenem (100%), Ciprofloxacin (92.3%), Gentamycin (92.3%), Chloramphenicol (84.6%), Amoxicillin/Clavulanic acid (84.6%) and Sulfamethoxazole/Trimethoprim (69.2%).

Keywords: Coliforms; antimicrobial; susceptibility; mastitis; bovines; pastoral herds.

1. INTRODUCTION

Mastitis is a disease of economic concern in dairy herds involving the swelling of the mammary gland which often results in changes in the physical, chemical and bacteriological characteristics of milk amongst other symptoms [1,2,3,4]. The occurrence of this disease is an outcome of interplay between three factors: infectious agents, host resistance, and environmental factors [5].

Mastitis is arguably the most widespread infectious diseases in dairy cattle. There is a consensus of opinions that it is the most destructive from an economic point of view [6,7,8,9,10,11]. The problem of mastitis is not a localised one as it adversely affects animal health, production of milk and the economics of milk production on a global scale. The challenge of mastitis cuts across both the developed and the developing economies of the world resulting in great economic losses [12].

The two major forms of the disease are the clinical and subclinical mastitis [13]. The clinical form is often obvious and can be detected easily. It is often characterised by changes in the composition and appearance of milk, reduction in the quantity of milk produced as well as the manifestation of such signs as pain, swelling and redness, with or without heat in infected mammary quarters [13]. Detection of the subclinical form of mastitis on the other hand is more difficult since the signs are not usually obvious [13].

Furthermore, Somatic Cell Count (SCC) has been accepted as the best index to use to predict udder infection in bovines, and has been used extensively as an indicator since the 1960s [13,14]. Under field conditions, determination of somatic cell count in milk is usually done using the California Mastitis Test (CMT); In fact, CMT

scores are directly related to average SCC [14]. CMT has found wide application principally because it is affordable and its results are very useful in the selection of the quarters for subsequent bacteriological examination [13].

Indiscriminate use and continuous abuse of antibiotics among the pastoralists for both therapeutic treatment of infections) and non-therapeutic (growth promoters) purposes on dairy animals has resulted to the increasing emergence of resistant strains of pathogenic bacteria, which is a great threat to human and animal health [11]. Hence, this study was embarked on to investigate the antimicrobial susceptibility profile of coliform isolates from mastitis milk samples of dairy cows among the pastoral herds in parts of Kaduna State, Nigeria.

2. MATERIALS AND METHODS

2.1 Study Area

The study was carried out in Giwa, Igabi, Chikun, Soba, Zaria, Sabongari and Birnin Gwari Local Government Areas (LGA) of Kaduna State, Nigeria (Fig. 1). These are seven out of the 23 LGAs in the state. The state lies between latitude 9.00° and 11.52° North and longitude 6.08° and 8.83° East and is 608 m above sea level. The number of LGAs studied was limited by the serious security challenge in the Northern part of Nigeria. The study area has distinct wet and dry seasons within the Guinea Savannah and part of the Sudan Savannah in Nigeria. Agriculture is the main occupation of Kaduna State with about 80% of the people actively engaged in farming. Another major occupation of the people is animal rearing and poultry farming. The animals reared include cattle, sheep, goats and pigs [15]. Pastoralism, Agro-pastoralism and intensive dairy farming are the predominant dairy production systems in Kaduna State. The pastoralists move around with their herds in

search of fresh pasture land or grazing areas. Agro-pastoralism is practiced by farmers who grow food crops and keep livestock, while the intensive dairy farmers use part or all of their land to grow fodder crops for their dairy cattle [16].




Fig. 1. The map of Nigeria and Kaduna State showing the study LGAs Adapted from the Administrative Map of Nigeria [17,18]

2.2 Study Design

A cross-sectional study was carried out among 147 lactating bovines from 30 herds spread across seven Nomadic settlements within seven LGAs in Kaduna state between May, 2017 and July 2018 using quantitative methods of data collection.

2.3 Inclusion and Exclusion Criteria

The study population constitutes all the lactating bovines of the Nigerian indigenous breeds within the study area. All farmers/pastoralist who declined consent as well as regions within the state that have been identified as volatile security spots were excluded. The animals were selected from herdsmen settlements in parts of Kaduna State, Nigeria. More so, only lactating bovines that are not currently on treatment were included, while those currently undergoing any form of treatment were excluded.

2.4 Sample Size Determination and Sampling Technique

The sample size was calculated using the formula of Sarmukaddam and Gerald [19] expressed by Eq. 1. Mbuk et al. [11] recorded a prevalence of 10.3% for bovine coliform mastitis in Kaduna state, Nigeria which was used for sample size estimation.

$$n = \frac{Z^2 p(1-p)}{L^2} \quad (1)$$

Where:

n = is the number of samples

Z = is the standard normal distribution at 95% confidence interval = 1.96

p = is the prevalence of previous study = 10.3% = 0.103

L = is the allowable error, which is taken at 5% = 0.05

Therefore, sample size,

$$n = \frac{1.96^2 \times 0.103 \times (1 - 0.103)}{0.05^2} = 142$$

A sample size of 142 was estimated at 5% level of significance. This was approximated to 147 for ease of proportionate distribution among the settlements.

A multi-stage sampling technique was used in this study. The seven LGAs were purposively

selected out of 23 LGAs in Kaduna state, being the LGAs with fewer security risks and were accessible at the time of this study. This was followed by the purposive selection of a settlement from each of the seven LGAs (total of seven settlements) based on the availability of lactating bovines that are not currently on treatment, willingness of the farmers/pastoralists to participate in the study, and accessibility of the location in order to easily transport samples collected to the laboratory for further analysis. Finally, 147 bovines were randomly and proportionately selected from all herds within the seven settlements. Bovine listing and enumeration was done to a total of 50, 30, 39, 27, 55, 40 and 68 for Settlements A, B, C, D, E, F and G, respectively out of which 24, 15, 19, 12, 26, 19 and 32 were respectively selected. A herd of bovines whose owner consented was sampled and in the event that he or she declined, the next contiguous herd of bovines was sampled. Computer generated list of random numbers from Minitab 14.2 statistical software was used to select the bovines for each of the settlements.

2.5 Sample Collection and Screening for Subclinical Mastitis

Strict aseptic procedures were followed to prevent contamination with microorganisms present on the skin of udder and teats, hands of samplers and barn environment according to the methods of National Mastitis Council Guidelines described by Middleton et al. [20]. Prior to milk sample collection, udders and teats were cleaned using a disposable paper towel immersed in 70% ethyl alcohol and dried to avoid presence of faecal contamination in the milk as it could interfere with the interpretation of CMT result. Foremilk (first jets) was discharged to reduce the contamination of teat canal. Sterile universal bottles with tight fitting cups were used. The bottles were labelled appropriately with permanent marker before sampling. To reduce contamination of teat ends during sample collection, the near teats were sampled first and then followed by the far ones. About 8ml of raw milk was aseptically collected from each bovine (2 ml from each quarter). The California Mastitis Test (CMT) Reagent was used according to the manufacturer's instructions on the field to identify samples with subclinical mastitis. The recommended 2 ml of milk samples was collected directly from each quarter of the udder and mixed together. This formed a composite from which 2 ml of the composite milk sample was then added to 2 ml of CMT reagent on the

test paddle and mixed gently to observe reaction. The result was graded as described by various authors [21,22]. All samples that tested positive for subclinical Mastitis were properly labelled and immediately transported to the Bacteriological Analysis Laboratory of the Department of Microbiology, Faculty of Life Science, Ahmadu Bello University, Zaria in an ice box for processing.

2.6 Bacteriological Analysis of CMT Positive Milk Samples

2.6.1 Inoculation of raw milk samples

The CMT positive milk samples were inoculated on MacConkey agar (Oxoid, England) by streak method as described by Mekonnin et al. [23]. A loop full of milk sample was streaked on the agar plates aseptically using quadrant method for each sample. The plates were incubated at 37°C and examined after 24 hours for growth.

2.6.2 Primary isolation of coliform bacteria

Bacteriological analysis was focused only on the identification and isolation of Coliform bacteria. Hence, pink to red distinct colonies resulting from the utilization of lactose on MacConkey agar were presumptively considered as Coliform bacteria. The suspected isolates were sub-cultured to get pure isolates. The pure isolates were cultured on Eosin Methylene Blue Agar (EMB) which is selective and differential for Coliform bacteria. Isolates that showed metallic green sheen on EMB were presumptively considered as *E. coli*, while those with other coloured appearance such as mucoid pink were considered to be other Coliform bacteria. The suspected Coliform isolates were stored in Nutrient Agar slant for further characterization and identification using the conventional biochemical tests and Microgen A+B ID Kits (UK).

2.6.3 Biochemical characterization

All suspected coliform bacterial isolates that stained red with Gram reaction were subjected to Conventional biochemical tests. The tests conducted were: Indole, Methyl Red, Voges-Proskauer and Citrate Utilization (IMViC). The suspected coliform bacterial isolates from the tests were identified up to species level using Microgen A+B Kit (UK) in accordance with the manufacturer's instructions.

2.6.4 Antibiotic susceptibility testing

Antibiotic susceptibility testing was conducted for all the isolated Coliform species using Kirby-Bauer disk diffusion method according to the criteria of the Clinical and Laboratory Standard Institute [24]. Direct colony suspension of the isolates was adjusted to a turbidity equivalent to a 0.5 McFarland standard and was aseptically inoculated on Mueller-Hinton agar (Oxoid, UK) using spread plate technique. The antibiotic impregnated disks (Oxoid, UK) were aseptically fixed on the solid agar surface 15mm apart using a dispenser (Oxoid, UK). The plates were incubated at 37°C for 24 hours.

Commercially available antibiotics (Oxoid, UK) recommended as drugs of choice against *enterobacteriaceae* and those frequently used in the treatment of human and animal infections within the study area were selected. Thus, a total of ten antibiotics were used. The antibiotic disks used with their various concentrations were: Amoxicillin-Clavulanic acid (30 µg), Imipenem (10 µg), Ciprofloxacin (5 µg), Gentamicin (30 µg), Chloramphenicol (30 µg), Trimethoprim/Sulphamethoxazole (25 µg), Erythromycin (15 µg), Penicillin (10 µg), Streptomycin (30 µg) and Tetracycline (30 µg).

Furthermore, the diameters of the zones of inhibition around the disks were measured to the nearest millimeter using caliper. The isolates were classified as susceptible, intermediate and resistant according to the interpretive standards of Clinical and Laboratory Standard Institute [24]. Moreover, isolates that showed resistance to two or more classes of antibiotics were considered as multidrug resistant [25,26,27,28].

3. RESULTS AND DISCUSSION

3.1 Prevalence of Subclinical Mastitis and Coliforms in the Study Area

Out of 147 milk samples from pastoral herds, 29 (19.7%) were positive for subclinical Mastitis out of which only 13 (8.8%) species of coliforms were isolated (six *E. coli* and seven *K. pneumoniae*). This implies that the prevalence of coliform mastitis in the study area (Parts of Kaduna state) is 8.8%. Samples from Birnin-gwari LGA harboured the highest number of coliforms 4 (2.7%) while no coliform bacteria were isolated from samples collected from Soba Local Government Area (Table 1). *K. pneumoniae* and *E. coli* were the species

associated with mastitis milk (Table 2 and Table 3).

3.2 Antimicrobial Susceptibility

The coliform isolates showed 100% resistance to Penicillin and Tetracycline, and were all 100% susceptible to Imipenem. Although this findings agree with previous reports that coliforms are 100% resistant to Penicillin [11,29], the 100% resistance recorded for Tetracycline is an indication of gross abuse of the drug through self-medication. This may not be unconnected to its availability and affordability. High multidrug resistance was expressed by all the isolates to Penicillin, Tetracycline and Erythromycin. All the isolates (100%) had MARI of 0.2 and above which is an indication of gross abuse of antibiotics within the studied population which was further buttressed by the 100% resistance displayed against penicillin and tetracycline. However, antibiotics still effective against the coliform species tested were Imipenem (100%), Ciprofloxacin (92.3%), Gentamycin (92.3%),

Chloramphenicol (84.6%), Amoxicillin/Clavulanic acid (84.6%) and Sulfamethoxazole/Trimethoprim (69.2%) (Table 4).

More so, all the isolates tested exhibited six resistance patterns (A-F) according to their resistance to different antimicrobial groups (Table 5).

All the isolates tested were considered multiple drug resistant (MDR) as they showed resistance to more than two classes of antibiotics tested. However, *Escherichia coli* isolates coded EC3 and EC4 had the highest MAR Index of 0.7 (resistant to seven out of ten antibiotics tested), followed by EC1, EC5 and EC6 with MAR Index of 0.4 (resistant to four out of 10 antibiotics tested). All the *Klebsiella pneumoniae* isolates coded KP1, KP2, KP3, KP4, KP5, KP6 and KP7 had MAR Index of 0.3 each (resistant to three out of ten antibiotics tested, while *Escherichia coli* isolate with code EC2 had the least MAR Index of 0.2 (resistant to two out of ten antibiotics tested) (Table 6).

Table 1. Prevalence of subclinical mastitis and associated coliform bacteria among the bovines studied

S/N	Local Government Area/ Settlements/Herds	No. of lactating bovines	No. of bovines examined	No.(%) of samples positive for subclinical mastitis	No.(%) of samples positive for coliform bacteria
1.	Giwa	50	24	6(4.1)	3(2.0)
2.	Igabi	30	15	5(3.4)	2(1.4)
3.	Chikun	39	19	3(2.0)	1(0.7)
4.	Soba	27	12	1(0.6)	0(0.0)
5.	Zaria	55	26	4(2.7)	1(0.7)
6.	Sabongari	40	19	3(2.0)	2(1.4)
7.	Birnin Gwari	68	32	7(4.8)	4(2.7)
	Total	309	147	29(19.7%)	13(8.8%)

Table 2. Biochemical characterization (IMVIC) of isolates

Suspected coliform isolates	Indole test	Methyl Red test	Vogues Proskauer test	Citrate Utilization test	Probable Organism
C1	+	+	-	-	<i>Escherichia</i> sp
C2	+	+	-	-	<i>Escherichia</i> sp
C3	+	+	-	-	<i>Escherichia</i> sp
C4	+	+	-	-	<i>Escherichia</i> sp
C5	+	+	-	-	<i>Escherichia</i> sp
C6	+	+	-	-	<i>Escherichia</i> sp
C7	-	-	+	+	<i>Klebsiella</i> sp
C8	-	-	+	+	<i>Klebsiella</i> sp
C9	-	-	+	+	<i>Klebsiella</i> sp
C10	-	-	+	+	<i>Klebsiella</i> sp
C11	-	-	+	+	<i>Klebsiella</i> sp
C12	-	-	+	+	<i>Klebsiella</i> sp
C13	-	-	+	+	<i>Klebsiella</i> sp

Key: C1-C5 = Probable *Escherichia* species, C6-C13 = Probable *Klebsiella* species

Table 3. Microgen tests for the identification of the isolates up to species level

Presumptive isolates	Octal number	Final identification	Percentage probability
EC1	04600570	<i>Escherichia coli</i> inactive	96.6%
EC2	05604520	<i>Escherichia coli</i> inactive	90.2%
EC3	04604420	<i>Escherichia coli</i> inactive	86.5%
EC4	04405421	<i>Escherichia coli</i> inactive	88.3%
EC5	07600570	<i>Escherichia coli</i>	49.8%
EC6	07601370	<i>Escherichia coli</i>	92.6%
KP1	47523766	<i>Klebsiella pneumonia</i>	99.7%
KP2	47523666	<i>Klebsiella pneumonia</i>	95.1%
KP3	47523777	<i>Klebsiella pneumonia</i>	95.2%
KP4	47523757	<i>Klebsiella pneumonia</i>	99.3%
KP5	47555777	<i>Klebsiella pneumonia</i>	87.3%
KP6	47544776	<i>Klebsiella pneumonia</i>	65.1%
KP7	47544777	<i>Klebsiella pneumonia</i>	57.7%

Key: EC1-EC6 = *Escherichia coli* (6); KP1-KP7 = *Klebsiella pneumoniae* (7)

Table 4. Antimicrobial susceptibility profile of coliform bacterial isolates obtained from mastitis milk samples (n=13)

S/N	Antibiotic generic name	Dics concentration (µg /ml)	No.(%) of resistance	No.(%) of intermediate resistance	No. (%) of susceptibility	Total (%)
01.	Erythromycin (E)	15.0	12(92.3)	0(0)	1(7.7)	13(100)
02.	Ciprofloxacin (CIP)	5.0	1(7.7)	0(0)	12(92.3)	13(100)
03.	Penicillin (PEN)	10.0*(I.U)	13(100)	0(0)	0(0.0)	13(100)
04.	Imipenem (IPM)	10.0	0(0.0)	0(0)	13(100)	13(100)
05.	Tetracycline (TET)	30.0	13(100)	0(0)	0(0.0)	13(100)
06.	Sulfamethoxazole/Trimethoprim (SXT)	25.0	4(30.8)	0(0)	9(69.2)	13(100)
07.	Chloramphenicol (C)	30.0	1(7.7)	1(7.7)	11(84.6)	13(100)
08.	Streptomycin (S)	10.0	3(23.0)	4(30.8)	6(46.2)	13(100)
09.	Amoxicillin/Clavulanic acid (AMC)	30.0	1(7.7)	1(7.7)	11(84.6)	13(100)
10.	Gentamicin(GN)	30.0	1(7.7)	0(0)	12(92.3)	13(100)

*Penicillin is given in International Units (IU).

The detailed zone of inhibition for respective isolates and corresponding antimicrobial agent is presented in Fig. 2.

Table 5. Antibiotic Resistance Patterns of Coliform Bacterial Isolates obtained from Mastitis Milk Samples (n=13)

Code	Resistance pattern	No. of isolates with the pattern	Percentage of occurrence
A	TET, PEN	1	7.7
B	E, PEN, TET	5	38.5
C	E, PEN, TET, S	3	23.0
D	E, PEN, TET, SXT	2	15.4
F	E, PEN, TET, SXT, C, S, AMC	1	7.7
E	E, CIP, PEN, TET, SXT, S, GN, C, AMC	1	7.7
	Total	13	100%

Key: E – Erythromycin; CIP – Ciprofloxacin; PEN – Penicillin; TET – Tetracycline; SXT - Sulfamethoxazole/Trimethoprim; C – Chloramphenicol; S – Streptomycin; AMC - Amoxicillin/Clavulanic acid; GN - Gentamycin

The species of coliforms isolated identified were *Klebsiella pneumoniae* and *Escherichia coli*. *Klebsiella pneumonia* was the dominant species associated with bovine mastitis. This finding is in agreement with the work of Mbuk et al. [11] who isolated similar species of these organisms in Kaduna State where *Klebsiella pneumonia* was

the highest, but *Escherichia coli* was not isolated in their study. These findings also agree with Hogan and Smith [30] who reported that *Klebsiella pneumoniae* and *Escherichia coli* are the species of coliforms most frequently isolated from cases of bovine mastitis. The dominance of *Klebsiella pneumoniae* agreed with the report of

Podder et al. [31] who reported that *Klebsiella pneumoniae* is well adapted to survive in the udder and usually establishes subclinical mastitis infection of long duration which can be shed in milk, facilitating transmission to healthy animals mainly during milking process. Generally, the presence of these Coliform bacterial species in the milk is an indication of faecal and environmental contamination resulting from poor hygienic practices of rearing and milking the animals as there are no established mastitis control practices employed among the herdsman.

The results of antimicrobial susceptibility test showed that all the species of coliforms tested were sensitive to Imipenem followed by decreasing sensitivity to Ciprofloxacin, Amoxicillin/Clavulanic acid, Streptomycin, Chloramphenicol, Gentamycin and Sulfamethoxazole/Trimethoprim. This agrees with the report of Mbuk et al. [11] and Lira et al. [32] that showed similar susceptibility pattern. Susceptibility of Imipenem to all coliform species tested has proven that Carbapenems still retain considerable potency against Enterobacteriaceae. This agrees with the recommendation of CLSI [24], where this class of antibiotics was among the recommended antibiotics for treatment of infections caused by Enterobacteriaceae. The high level of susceptibility to Imipenem might be due to its rare use and abuse in cattle. However, it is worthy of note here that the Coliform species tested showed 1.4% intermediate resistance and 5.1% resistance to some CLSI [24]

recommended antibiotics (Ciprofloxacin, Amoxicillin/Clavulanic acid, Streptomycin, Chloramphenicol, Gentamycin and Sulfamethoxazole/ Trimethoprim). Therefore, irrational prescriptions and indiscriminate use of these drugs may lead to complete resistance in the future [33].

All the species of Coliform bacteria tested were completely resistant to Penicillin and Tetracycline. Coliforms are however naturally resistant to penicillin as reported previously. This susceptibility pattern however agrees with previous studies of high resistance to these same antibiotics [11,29,32]. The high degree of resistance observed might be due to prolonged and indiscriminate usage of those antibiotics which could lead to possible resistance development in humans and animals [34,35].

Moreover, all the *Klebsiella pneumoniae* and *Escherichia coli* species exhibited multidrug resistance, as they were consistently resistant to two or more classes of antibiotics among others used especially Erythromycin, Penicillin and Tetracycline. This finding agreed with the previous reports where Coliform species tested displayed multidrug resistance to Erythromycin, Penicillin and Tetracycline [32]. These findings however, contradict the report of Memom *et al.* [29] where coliform species were completely resistant to Ciprofloxacin, Gentamycin, Amoxicillin and Sulfamethoxazole/ Trimethoprim. However, based on this study, antibiotics still effective against the coliform species tested were Imipenem (100%),

Table 6. Multiple Antibiotic Resistance (MAR) index of coliform species investigated

Organisms	Isolate code	Number of antibiotics tested	MAR index	Number of classes of antibiotics tested	Number to which isolates were resistant	Resistance pattern
<i>E. coli</i>	EC1	10	0.4	6	4	E, PEN, TET, STX
	EC2	10	0.2	6	2	PEN, TET
	EC3	10	0.7	6	7	E, CIP, PEN, TET, STX, S, GN
	EC4	10	0.7	6	7	E, PEN, TET, STX, C, S, AMC
	EC5	10	0.4	6	4	E, PEN, TET, S
	EC6	10	0.4	6	4	E, PEN, TET, STX
<i>K. pneumoniae</i>	KP1	10	0.3	6	3	E, PEN, TET
	KP2	10	0.3	6	3	E, PEN, TET
	KP3	10	0.3	6	3	E, PEN, TET
	KP4	10	0.3	6	3	E, PEN, TET
	KP5	10	0.3	6	3	E, PEN, TET
	KP6	10	0.3	6	3	E, PEN, TET
	KP7	10	0.3	6	3	E, PEN, TET




Fig. 2. Measured zones of inhibition to antibiotics tested in millimeters

Key: E – Erythromycin; CIP – Ciprofloxacin; PEN – Penicillin; IPM – Imipenem; TET – Tetracycline; STX - Sulfamethoxazole/Trimethoprim; C – Chloramphenicol; S – Streptomycin; AMC - Amoxicillin/Clavulanic acid; GN – Gentamycin

Ciprofloxacin (92.3%), Gentamycin (92.3%), Chloramphenicol (84.6%), Amoxicillin/Clavulanic acid (84.6%) and Sulfamethoxazole/Trimethoprim (69.2%). Although Chloramphenicol is prohibited for use in animal treatments, it is still being applied in Nigeria since there is no legal framework for its prohibition yet. It remains one of the highly used antibiotics among the Nomads. However, the higher susceptibility of the coliforms to Imipenem, Ciprofloxacin, Amoxicillin/Clavulanic acid and Streptomycin than Chloramphenicol makes them better alternatives to it. This study therefore recommends the discouragement of the use of Chloramphenicol in animal treatment in Nigeria in line with global best practice.

Furthermore, the result of susceptibility pattern of Coliform bacterial species obtained affirms that some of CLSI [24] recommended antibiotics of choice against the treatment of infections caused by Enterobacteriaceae are increasingly becoming ineffective within the studied population. Therefore, it is always very important to conduct antimicrobial sensitivity tests before empirical therapy is initiated to avoid resistance development to other sensitive antibiotics in future. However, based on the degree of susceptibility pattern obtained in this study, Imipenem is recommended as first line drug of choice where infection by *K. pneumonia* and *E. coli* respectively is suspected within the studied area.

4. CONCLUSION

This study concludes that the prevalence of subclinical mastitis in Kaduna State is 19.7% while the prevalence of Coliform Mastitis is 8.8%. A low prevalence of Coliform mastitis was observed in this region, but the presence of *Klebsiella pneumonia* and *Escherichia coli* in raw milk samples of the studied bovine constitute serious environmental health risk to the consumers as the milk obtained from these herds are widely circulated and consumed without any form of treatment. They are also among the list of organisms classified as dangerous biological agents that have the potential to pose a severe threat to public health and safety by United States Public Health Services. The species of coliforms isolated showed decreased sensitivity to the majority of recommended antibiotics of choice by Clinical and Laboratory Standard Institute (CLSI). This phenomenon could result to complete resistance development in future if not properly handled. The high level of resistance to

some of the commonly used antibiotics by the herdsmen imply that the selection pressure imposed by the use of these antibiotics whether therapeutically in veterinary medicine or as prophylaxis in the animal production is a key driving force in the selection of antimicrobial resistance.

CONSENT

Ethical consent was obtained from the postgraduate board of the department of microbiology, faculty of life sciences, Ahmadu Bello University for Zaria to undertake the study and to publish this report and accompanying images.

ETHICAL APPROVAL

All authors hereby declare that "Principles of laboratory animal care" (NIH publication No. 85-23, revised 1985) were followed, as well as specific national laws where applicable. All experiments have been examined and approved by the appropriate ethics committee.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Beheshti R, Shaieghi J, Eshratkhan B, Ghalehkandi JG, Maheri-Sis N. Prevalence and etiology of subclinical mastitis in ewes of the Tabriz region, Iran. *Global Veterinaria*. 2010;4(3):299-302.
2. Hussain R, Javed MT, Khan A, Mahmood F, Kausar R. Mastitis and associated histopathological consequences in the context of udder morphology. *International Journal of Agriculture and Biology*. 2012;14(6): 947-952.
3. Radostis OM, Gay CC, Blood DC, Hinchliff KW. *Veterinary medicine: A textbook of the disease of cattle, sheep, pigs, goats and horses*. 9th Ed. ELBS and Baillier Tindall. 2000;563-660.
4. Sharma N, Singh NK, Bhadwal MS. Relationship of somatic cell count and mastitis: An overview. *Asian-Australian Journal of Animal Science*. 2011;24(3): 429-438.
5. Gera S, Guha A. Assessment of acute phase proteins and nitric oxide as indicator

- of subclinical mastitis in Holstein Å Haryana cattle. *Indian Journal of Animal Sciences (India)*. 2011;81(10):1029-1031.
6. Halasa T, Huijps K, Østerås O, Hogeveen H. Economic effects of bovine mastitis and mastitis management: A review. *Veterinary Quarterly*. 2007;29(1):18-31.
 7. Ashish T, Sisodia RS, Sharma RK, Misraulia KS, Garg UK. Incidence of sub-clinical mastitis in cows of Malwa region of Madhya Pradesh. *Indian Journal of Dairy Science*. 2000;53(4):328-31.
 8. Shittu A, Abdullahi J, Jibril A, Mohammed AA, Fasina FO. Sub-clinical mastitis and associated risk factors on lactating cows in the Savannah Region of Nigeria. *BMC Veterinary Research*. 2012;8(1):134.
 9. Lamey AE, Ammar AM, Zaki ER, Khairy N, Moshref BF, Refai MK. Virulence factors of *Escherichia coli* isolated from recurrent cases of clinical and subclinical mastitis in buffaloes. *Intl. J.* 2013;4(1):86-94.
 10. Suleiman AB, Umoh VJ, Kwaga JK, Shaibu SJ. Enterotoxigenicity and antibiotic resistance of *Staphylococcus aureus* isolated from sub-clinical bovine mastitis milk in Plateau State, Nigeria. *Research Journal of Microbiology*. 2013;8(2):101.
 11. Mbuk EU, Kwaga JK, Bale JO, Boro LA, Umoh JU. Coliform organisms associated with milk of cows with mastitis and their sensitivity to commonly available antibiotics in Kaduna State, Nigeria. *Journal of Veterinary Medicine and Animal Health*. 2016;8(12):228-36.
 12. Sharma N. Alternative approach to control intramammary infection in dairy cows-A review. *Asian J. Anim. Vet. Adv.* 2007;2(2): 50-62.
 13. Kivaria FM. Epidemiological studies on bovine mastitis in smallholders dairy herds in the Dar es Salaam Region, Tanzania. Doctoral Thesis, Utrecht University, The Netherlands; 2006.
 14. Pyörälä S. Indicators of inflammation in the diagnosis of mastitis. *Veterinary Research*. 2003;34(5):565-578.
 15. KDSG. Kaduna State achievement in data on estimated annual animal population, fish production and investment opportunities in Kaduna State. Kaduna State Government. 2008;16-18.
 16. Ugwu DS. Dairy production among small and medium scale farmers in Nigeria: A case study of Kaduna and Kano States. *World Journal of Agricultural Sciences*. 2010;6(1):01-6.
 17. Library of Congress. Administrative Map of Nigeria. Available:https://www.loc.gov/item/2010592721/ (Accessed 08/06/2017)
 18. Available:http://www.nationsonline.org/one-world/map/nigeria-administrative-map.htm (Accessed 08/06/2017)
 19. Sarmukaddam SS, Gerald SG. Validity of assumption while determining sample size. *Indian Journal of Community Medicine*. 2006;20-29.
 20. Middleton JR, Saeman A, Fox LK, Lombard J, Hogan JS, Smith KL. The National Mastitis Council: A global organization for mastitis control and milk quality, 50 years and beyond. *Journal of Mammary Gland Biology and Neoplasia*. 2014;19(3-4):241-51.
 21. Quinn PJ, Markey BK, Carter ME, Donnelly WJC, Leonard FC, Maghir D. *Veterinary microbiology and microbial disease*. Blackwell Science Ltd, London. 2002;191-208.
 22. Das D, Panda SK, Jena B, Sahoo AK. Somatic cell count: A biomarker for early diagnosis and therapeutic evaluation in bovine mastitis. *Int. J. Curr. Microbiol. App. Sci.* 2018;7(3):1459-63.
 23. Mekonnin E, Eshetu E, Awekew A, Thomas N. A study on the prevalence of bovine mastitis and associated risk factors in and the surrounding areas of Sodo Town, Wolaita Zone, Ethiopia. *Global Journal of Science Frontier Research: D Agriculture and Veterinary*. 2016;16(2):13-5.
 24. Clinical and Laboratory Standard Institute (CLSI). Performance standards for antimicrobial susceptibility testing. 26th Edition. CLSI Supplement M100S. Wayne PA, USA; 2016.
 25. Intrakamhaeng M, Komutarin T. Antibiotics resistance and RAPD-PCR typing of multidrug resistant MRSA isolated from bovine mastitis cases in Thailand. *Science Asia*. 2012;38:30-35.
 26. Schabauer A, Pinior B, Gruber CM, Firth CL, Käsbohrer A, Wagner M, Rychli K, Obritzhauser W. The relationship between clinical signs and microbiological species, spa type, and antimicrobial resistance in bovine mastitis cases in Austria. *Veterinary Microbiology*. 2018;227:52-60.

27. Tomazi T, Coura FM, Gonçalves JL, Heinemann MB, Santos MV. Antimicrobial susceptibility patterns of *Escherichia coli* phylogenetic groups isolated from bovine clinical mastitis. *Journal of Dairy Science*. 2018;101(10):9406-18.
28. Zhang D, Zhang Z, Huang C, Gao X, Wang Z, Liu Y, Tian C, Hong W, Niu S, Liu M. The phylogenetic group, antimicrobial susceptibility, and virulence genes of *Escherichia coli* from clinical bovine mastitis. *Journal of Dairy Science*. 2018;101(1):572-80.
29. Hogan J, Smith KL. Coliform mastitis. *Veterinary Research*. 2003;34(5): 507.
30. Podder MP, Rogers L, Daley PK, Keefe GP, Whitney HG, Tahlan K. Klebsiella species associated with bovine mastitis in Newfoundland. *PloS one*. 2014;9(9): e106518.
31. Agunbiade TB, Umoh VJ, Whong CMZ, Ella EE. Assessment of bovine milk obtained from selected farms in Zaria environs for toxigenic strains of *Escherichia coli*. *Academic Journal of Life Sciences*. 2015;1(1):8-13.
32. Memon J, Kashif J, Yaqoob M, Kiping W, Yang Y, Hongjie FC. Molecular characterization and antimicrobial sensitivity of pathogens from subclinical mastitis in Eastern China. *Pakistan Veterinary Journal*. 2013;33(2):170-174.
33. Kwaga JKP. Veterinary intervention on the global challenges of antimicrobial resistance. Paper Presented at the World Veterinary Day Celebration, 28th April, NVMA, Kaduna; 2012.
34. Sharma N, Dogara BB, Misra R, Gandham N, Sardar M, Jadhav S. Multidrug resistant *Pantoea agglomerans* in patients with septic arthritis – a rare report from India. *International Journal of Microbiological Research*. 2014;4:263-265.
35. Devriese LA, Daube G, Homme J, Haesebrouck FS. *In vitro* susceptibility of *Clostridium perfringens* isolated from farm animals to growth-enhancing antibiotics. *Journal of Applied Microbiology*. 1993;75: 55-57.

© 2019 Makolo et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

*The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/46287>*