



Top Edible Wild Plants of Eastern Mediterranean Region. Part III: Antidiabetic Activity

Abdullatif Azab^{a*}

^a Eastern Plants Company, Box 868, Arara, Israel.

Author's contribution

The sole author designed, analyzed, interpreted and prepared the manuscript.

Article Information

DOI: 10.9734/EJMP/2023/v34i91160

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/107662>

Review Article

Received: 04/08/2023
Accepted: 09/10/2023
Published: 18/10/2023

ABSTRACT

In the first two parts of this series of review articles, we presented the anticancer and anti-inflammatory activities of the most important wild edible plants of eastern Mediterranean region, which we named as the "Deca-plants" (D-P). In this review article, we will present the antidiabetic activities of these very important plants. Comparing with anticancer and anti-inflammatory activities, the overall antidiabetic activity of the D-P is moderate, but there are clear differences between the ten species. Based on this fact and on the knowledge that the D-P contain several natural products with proven antidiabetic activity, in the discussion section (4), possible future research directions will be indicated. In addition to presenting the reported antidiabetic activity of the plants, some important statistical facts about diabetes will be presented, as well as ethnomedicinal use of these plants for diabetes treatment, in the reviewed region. For the purpose of comparison and comprehensiveness, in the last part of this article, four Non-Deca-Plants with reported antidiabetic activity will be shortly reviewed, when the criteria of selection are wild and edible.

Keywords: *Diabetes mellitus; antidiabetic; hypoglycemic; insulin resistance; type II diabetes; medicinal plants; α -amylase; plant extracts; essential oils; α -glucosidase.*

*Corresponding author: E-mail: eastern.plants@gmail.com;

1. INTRODUCTION

Diabetes mellitus (DM), often referred to as diabetes, is one of the worst chronic diseases of the modern era. According to the World Health Organization (WHO), DM is the ninth cause of human deaths globally [1]. According to the organization, the highest ranking of DM-caused death is in the middle-income countries, sixth.

Although DM is clearly a modern times disease, archeological research found that the remains of a 61-year-old (2050-1911 BC) male in ancient Egypt, clearly indicate that this person had severe conditions of DM [2]. The artifacts found with the mummy of this person, even though some were looted in antiquity, and more important, the fact that he lived to such a long age (at that time) with a few chronic diseases, lead to two conclusions. First, this person was wealthy and could afford medical care. Second and more important, there was treatment for DM at that time. The researchers indicate that the first documentation of DM was also in ancient Egypt, around 1530 BC, in the Ebers papyrus.

In a very detailed and comprehensive report, K.L. Ong and many contributing authors, presented current time (2021) statistics about DM, with prediction until 2050 [3]. According to this report, in 2021, there are 529 million diabetic people around the world (6.1%), and in 2050 there will be 1.31 billion diabetics (more than 10%). The lowest current prevalence of diabetes in 2021 was found in Sub-Saharan Africa (2%), and the highest in North Africa and the Middle East (9.3%). Nationally, Qatar is with a shocking 76.1% of DM prevalence.

P. Zhang and his colleagues predicted that the global expenditure on DM is expected to rise from USD 490 billion in 2020 to USD 561 billion in 2030, keeping in mind that it was USD 418 billion in 2010 [4]. And according to the 2022 report of the American Diabetes Association (for 2019), the relation between DM prevalence and ethnicity in the USA, is consistent with the global trends [5]. In 2019, 37.3 million Americans, or 11.3% of the population, had DM. 1.4 million Americans are diagnosed with diabetes every year. The ethnicities of the diabetics are: 14.5% American Indians/Alaskan natives, 12.1% non-Hispanic blacks, 11.8% Hispanics, 9.5% Asian Americans, and, 7.4% non-Hispanic whites. In the same year, 87,647 Americans died because of DM. The total spending on DM was \$327

billion, where \$237 billion was for direct medical costs and \$90 billion was in reduced productivity. And to finalize the presentation for the USA, J.E. Rodríguez and K.M. Campbell found that prevalence of DM among the different ethnic groups is consistent with other reports, but there are clear differences in health care for DM patients and the drugs used for treatment, in the different ethnic groups [6].

In the Arab world (21 countries), there is a direct proportion between DM and two variables: daily calories intake and gross domestic product (GDP) [7]. Based on this, it is easy to understand the ranking of the Gulf countries on the top of diabetics list, while poor Arab countries are on the bottom. Interestingly, the situation in Israel is partially contradicting: Jews (78% of the population), who have higher income but less carbohydrate consumption, have 12% diabetics, compared with Arabs (22% of the population), who have 24% diabetics [8].

Damages of DM are direct, including death, but the disease complications are life quality damaging, no less than DM itself, reports A.D. Deshpande and her colleagues [9]. These complications are shown in Fig. 1.

D. Tomic and her colleagues refer to the DM-complications presented in Fig. 1. as "traditional complications" [10]. They add non-traditional DM-complications such as cancer (several types), infections, liver disease, functional disability, cognitive disability, and affective disorders. And S. Hill Golden added another severe DM-complication: depressive symptoms [11]. They found that 22.0 out of 1000 DM patients were diagnosed with depressive symptoms, while 16.6 out of 1000 DM patients were diagnosed with elevated depressive symptoms.

To conclude this introduction, we will present a very important phenomenon in DM: insulin resistance. To illustrate the importance of this severe health disorder, we will cite M. Li and her colleagues that defined the discovery of insulin as: the transformation of diabetes from a fatal diagnosis into a medically manageable chronic condition [12]. They present major mechanisms of insulin resistance (excellent figures) and possible future therapies. S-H. Lee and her colleagues define insulin resistance as a key pathogenic component of many metabolic diseases and is defined as a state of reduced responsiveness of insulin-targeting tissues to physiological levels of insulin [13]. In addition to

excellent schemes and explanations, they emphasize the role of insulin resistance mainly in DM, but in other diseases as well. This relation is also highlighted by X. Zhao and her colleagues, but they focus on the possible induction of metabolic diseases by insulin resistance [14]. And D.J. Rader summarized the major factors that produce insulin resistance and the major outcomes of it [15], and these are presented in Fig. 2.

The D-P are *Arum palaestinum* (Araceae), *Cichorium pumilum* (Syn. *Cichorium endivia*, Asteraceae), *Cyclamen persicum* (Primulaceae), *Foeniculum vulgare* (Apiaceae), *Gundelia tournefortii* (Asteraceae), *Majorana syriaca* (Syn. *Origanum syriacum*, Lamiaceae), *Malva sylvestris* (Malvaceae), *Micromeria fruticosa* (Lamiaceae), *Salvia fruticosa* (Syn. *S. triloba*, *S. libanotica*, *S. cypria*, *S. lobryana*, Lamiaceae), *Sinapis alba* (Brassicaceae).

2. ETHNOMEDICINAL ANTIDIABETIC ACTIVITIES OF THE DECA-PLANTS

When comparing the antidiabetic activity of domesticated and wild plants, it is correct to say that the first group is more active than the second. Most of the commonly used plants for DM treatment are domesticated, as listed by M. Przeor: *Morus alba*, *Cinnamomum zeylanicum*, *Trigonella foenum-graecum*, *Phaseolus vulgaris*, *Zingiber officinale*, and *Panax ginseng* [16].

Based on this, and especially in countries where the D-P are rare or not found, it is clear why they are not mentioned in ethnomedicinal literature, such as the extensive review article of A.M. Abu-Odeh and W.H. Talib [17]. This article presents a partial state of knowledge and practice, and it focuses on Arabian Peninsula. So, six of D-P are not mentioned, but four of them are: *Arum palaestinum* (Jordan), *Cichorium pumilum* (Jordan), *Gundelia tournefortii* (Iran), *Salvia fruticosa* (Lebanon, Turkey).

Several publications presented the antidiabetic activity in Jordanian ethnomedicine. S.A. Ootom and his colleagues mentioned only *Majorana syriaca* in their review article [18]. Contrary to that, A. Al-Aboudi1 and F.U. Afifi reported that *C. pumilum*, *G. tournefortii*, *M. syriaca* and *S. fruticosa*; are used in Jordanian traditional medicine [19]. And in another review article published in the same year, F.U. Afifi and her colleagues reported only the use of *C. pumilum*

and *M. syriaca* for DM treatment in Jordanian ethnomedicine [20]. And to make these reports even more ambiguous, F.U. Afifi and V. Kasabri, coauthors of references 19 and 20, reported two years later that only *S. fruticosa* is “claimed” to have antidiabetic activity [21]. R. Issa and her colleagues investigated mixtures of medicinal plants used to treat DM in Jordanian folk medicine and reported *S. fruticosa* as a component of a mixture containing seven other plants [22].

Among the very few reports of Lebanese plants used in traditional medicine to treat DM, Z. Jeambey and her colleagues mentioned the use of *C. pumilum* and *G. tournefortii* [23]. They mentioned two other plants of the D-P, *F. vulgare* and *M. sylvestris*, but not in the context of antidiabetic activity. Interestingly, Z. Yaniv and her colleagues from Israel, reported the use of *M. syriaca* and *S. fruticosa* in Israeli folk medicine [24].

Palestinian traditional medicine uses four plants (like the Jordanian) according to M.S. Ali-Shtayeh and his colleagues [25]. Their antidiabetic plants list includes *A. palaestinum*, *C. pumilum*, *F. vulgare*, *M. syriaca* and *S. fruticosa*.

3. ANTIDIABETIC ACTIVITIES OF THE DECA-PLANTS AND THEIR NATURAL PRODUCTS

As we mentioned above, traditional medicines of the reviewed region used either edible domesticated plants for antidiabetic treatments, or edible and inedible wild plants that did not include the D-P. Those played a minor role. The modern scientific research followed the ethnomedicines, but the findings were surprising for some of these plants. These published findings are presented in **Table 1**.

4. DISCUSSION

The search for antidiabetic plants products started as early as humans recognized diabetes as a disease, and this search is still going on, and it is increasing continuously since humans realized the fatal nature of this disease and its complications. And as we mentioned earlier and in the previous article of this series, the search for antidiabetic plant agent with the Mediterranean diet, resulted in discovering notable advantages of this nutritional style [80].

Table 1. Published Antidiabetic Activities of the D-P in Eastern Mediterranean region

Testing Method, Results and Reference/s
<p><i>Arum palaestinum</i></p> <p>Leaves and flowers were separately extracted with ethanol and water. Both extracts were tested for antidiabetic activity: inhibition of pancreatic lipase, inhibition of α-amylase, inhibition of α-glucosidase, glucose lowering in starch/glucose fed rats. In all tests, aqueous extract was more potent than ethanolic, and the overall activities were weak. [26]</p> <p>Leaves were separately extracted with <i>n</i>-hexane and 50% aqueous ethanol. The <i>n</i>-hexane extract had notable α-amylase inhibition activity ($IC_{50} = 25.34 \mu\text{g/mL}$). The polar extract had weak activity ($IC_{50} = 573.72 \mu\text{g/mL}$). [27]</p>
<p><i>Cichorium pumilum</i></p> <p>Dry leaves water suspension was orally administered (500 mg/kg) to STZ-induced diabetic rats (STZ, streptozotocin). Testing several biomarkers showed that the suspension had similar activity to antidiabetic drug of glibenclamide (600 mg/kg). [28]</p> <p>80% Aqueous ethanol leaves extract was prepared and had clear activity in alloxan-induced diabetic rats. [29]</p>
<p><i>Cyclamen persicum</i></p> <p>No published studies (see Discussion).</p>
<p><i>Foeniculum vulgare</i></p> <p>Seeds powder was soaked in diethyl ether, and the solvent was removed to obtain an extract, the authors name "fixed oil". This material had negligible effect in alloxan-induced diabetic rats. [30]</p> <p>Fruits methanolic extract had clear effect against dexamethasone-induced insulin resistance in rats. [31]</p> <p>Commercial essential oil (EO, plant part/s not indicated) was orally fed (30 mg/kg) to STZ-induced diabetic rats, resulting clear positive effect in tested biomarkers. [32]</p> <p>Fruits were successively extracted with petroleum ether, chloroform, and dichloromethane. The petroleum ether extract had notable positive effect in STZ-induced diabetic rats (10 mg/kg, twice a day, 45 days). Effect was measured by lowering blood glucose, inhibition of aldose reductase, and the kinetics of <i>trans</i>-anethole (Fig. 3). concentration, which was indicated as the active compound. [33]</p> <p>Seeds were separately extracted with chloroform, ethyl acetate, acetone, methanol, ethanol, <i>n</i>-butanol, and water, and all extracts were analyzed for general chemical composition. The aqueous extract was administered to STZ-induced diabetic rats (orally, 150, 300 mg/kg, 35 days). Both doses were active, but 300 mg/kg was more potent, by measuring several biomarkers. [34]</p> <p>Leaves methanolic extract was used to treat alloxan-induced diabetic mice: 1.2, 2.2 mg/kg, 30 days. The higher dose was more active, as shown by measuring several biomarkers. [35]</p> <p>Fruits aqueous and 96% ethanolic extracts were prepared and tested for glycogen phosphorylase inhibition activity. Both extracts showed negligible potency. [36]</p> <p>Aerial parts <i>n</i>-hexane extract was prepared and used to treat STZ-induced diabetic rats: 100 mg/kg, 14 days. Compared with antidiabetic drug of glibenclamide, the extract had significant activity found in 8 different biochemical tests. [37]</p> <p>Commercial <i>trans</i>-anethole (Fig. 3.) was orally administered to STZ-induced diabetic rats (180 mg/kg, 45 days), resulting positive effect measured by concentrations of three biomarkers: blood glucose, insulin, blood and tissue glycoprotein. [38]</p> <p>Commercial seeds EO and <i>trans</i>-anethole were used to prepare nanoemulsions (2%), and these were used to treat STZ-induced diabetic rats. Positive effects were measured compared with control, and non-emulsified EO and <i>trans</i>-anethole. [39]</p> <p>Seeds 70% aqueous ethanolic extract was prepared and used to treat STZ- induced diabetic rats (50, 75 and 100 mg/kg/day, 30 days). Five biochemical parameters were tested showing positive effect of treatment with the extract, where all doses were active, and the highest was most potent. [40]</p> <p>Seeds aqueous extract was used to treat alloxan-induced diabetic rats, separately and in combination with propolis. In both cases, positive effects were measured (4 biochemical tests), and the combination proved to be more active. [41,42,43]</p>

Seeds aqueous extract was used to treat STZ-induced diabetic rats, separately and in combination with *Cassia angustifolia* leaves aqueous extract (150 mg/kg/day, 30 days). In both cases, positive effects were measured (10 biochemical tests), and the combination proved to be more active. [44] Fruits were extracted with 90% aqueous methanol, and so was the whole plant of *Tagetes erecta*. The extracts used for treatment of STZ-induced diabetic rats, separately and in combination (orally 500 and 100 mg/kg/day, 28 days, respectively). The positive control in this study was glibinclamide, and seven biomarkers were tested, resulting significant activity of the *F. vulgare* and higher for the combination. [45]

α -Pinene (Fig. 1). was used to treat alloxan-induced diabetic mice: intraperitoneal injection, 0.25 mL/kg, and fasting blood glucose was measured compared with glibinclamide as positive control. Results showed notable effect. [46]

Seeds were separately extracted with ethyl acetate, benzene, and *n*-butanol. Extracts in 5 different concentrations were tested for α -amylase and α -glucosidase, with acarbose as a reference. Ethyl acetate and benzene extracts were most active, especially for α -glucosidase inhibition. [47]

Seeds powder was administered to human subjects fed with glucose, smokers and nonsmokers, 50 mg/kg of body weight, in a single dose. Blood glucose was measured after 2 h showing significant effect of lowering blood glucose, compared with control group of untreated subjects. Effect was higher in smokers group. [48]

Seeds were extracted with 80% aqueous ethanol and extract was tested for dipeptidyl peptidase-IV inhibition, with sitagliptin a reference. Activity was low. [49]

Leaves aqueous extract (single dose, orally, 10 mg/kg) reduced blood glucose in STZ-induced diabetic rats. [50]

Seeds 70% ethanolic extract and *trans*-anethole were used to treat STZ-induced diabetic rats: orally, 200, 400 mg/kg/day, 35 days. Testing several biomarkers, especially blood glucose concentration, showed significant positive effect in both treatments. [51]

Roots and leaves were separately extracted with water, and extracts had notable α -glucosidase and α -amylase inhibition. Extracts were analyzed for chemical composition. Five compounds with highest concentrations in each extract are shown in Fig. 4. [52]

Seeds were extracted with 20% aqueous ethanol, and cookies that contained 3% extract were supplemented to hyperglycemic human subjects (4 cookies in a single dose, mass not indicated). Tests indicated significant reduction of blood glucose concentrations [53].

Computational prediction of α -glucosidase inhibition activity of several major ingredients of the plant. [54]

Seeds EO had very weak α -glucosidase inhibition activity. [55]

Seeds aqueous extract had significant positive effect in STZ-induced diabetic rats: 150 mg/kg/day, 30 days. [56]

Leaves ethanolic extract had notable α -amylase and α -glucosidase inhibition activity, with acarbose as a reference. [57]

Gundelia tournefortii

Leaves were extracted with petroleum ether and extract had efficient α -amylase inhibition activity. [58]

Roots aqueous extract was used to treat dexamethasone-induced diabetic mice: 75, 150 and 300 mg/kg/day, 22 days. All doses lowered blood glucose and the highest dose was most active. [59]

Shoots aqueous extract was used to treat STZ-induced diabetic rats: 200, 400 mg/kg/day, 21 days. Both treatments reduced blood glucose and the higher dose was more effective. [60]

Aerial parts were extracted with *n*-hexane and methanol. Both extracts were *in vitro* tested for glucose transporter 4 (GLUT4) translocation in rat L6 muscle cell lines. Both extracts were active and methanolic extract was more potent. Both extracts were analyzed for chemical compositions, where stigmasterol (Fig. 5). was the only compound present in both extracts. [61]

Aqueous extract of aerial parts was used to treat alloxan-induced diabetic mice: 5, 10, 20, 40 mg/kg/day, 20 days. All doses reduced blood glucose like glibenclamide that was used as a reference. [62,63]

Aerial parts were extracted with ethanol, and extract was fractionized with water, ethyl acetate and *n*-hexane. Extract and fractions were tested for aldose reductase inhibition, where ethyl acetate fraction was most active. [64]

Roots and stems were separately extracted with ethanol, and both extracts were tested for α -

amylase and α -glucosidase inhibition. Stems extract was more active with IC_{50} (mg/mL) 4.22 for α -amylase, and 10.37 for α -glucosidase. [65]

A follow up study of the research presented in reference [61], with more extracts, and identification of the compounds responsible for GLUT4 translocation activity. Four of these compounds are shown in Fig. 6. [66]

Majorana syriaca

Leaves aqueous extract and EO were prepared, but only the extract was tested for α -amylase and α -glucosidase inhibition, resulting significant activity for both enzymes, with acarbose as a reference. The EO composition was determined showing that it contains two major compounds: γ -terpinene and *p*-cymene, 31.6%, 39.1%, respectively (Fig. 7). [67]

Malva sylvestris

Leaves were separately extracted with *n*-hexane and 50% aqueous ethanol. The hydrophilic extract had notable α -amylase inhibition activity (IC_{50} = 38.55 μ g/mL). The lipophilic extract had very weak activity. [27]

Flowers ethanolic extract had high α -amylase and α -glucosidase inhibition activities (IC_{50} = 7.8, 11.3 μ g/mL, respectively), with acarbose as a reference. [68]

Leaves were extracted with 40% aqueous methanol and the extract was fed to common carp (*Cyprinus carpio*), resulting increase of α -amylase activity. [69]

Pasta was enriched with 3, 6% dried leaves, then fed to healthy volunteers. Pasta glycemic index was reduced by 21.6%, 24.3%, respectively. The 50% aqueous ethanol extract of these dried leaves had significant α -amylase and α -glucosidase inhibition activities (IC_{50} = 12.95, 27.55 mg/g, respectively), with acarbose as a reference. [70]

Micromeria fruticosa

Volatile oils were obtained from aerial parts harvested in three locations, using ultrasonic-assisted extraction, and their chemical compositions were determined. Major three compounds are presented in Fig. 8. Oils had very high α -amylase inhibition activities compared with acarbose as a reference. [71]

Salvia fruticosa

Dry leaves powder was used to prepare 10% infusion, that was orally supplemented to alloxan-induced diabetic rabbits, resulting notable reduction of blood glucose. [72]

80% Aqueous methanolic and chloroform extracts of aerial parts were prepared. Chemical analysis resulted two new phenolics (Fig. 9), known NPs that were isolated for the first time from the genus *Salvia*, and other NPs that were isolated for the first time from *S. fruticosa*. The aqueous methanolic extract had blood glucose lowering activity in alloxan-induced diabetic rats, where glibenclamide was the reference. [73]

80% Roots aqueous ethanolic extract was prepared and had blood glucose lowering activity in alloxan-induced diabetic mice, where glibenclamide was the reference. [74]

Aqueous leaves extract and was tested *in vivo*: alloxan-induced diabetic rats, 50, 150, 450 mg/kg/day, 17 days, orally; and *in vitro* for inhibition of α -amylase and α -glucosidase inhibition. Significant activity was measured in both tests, and in the *in vivo* test, blood glucose concentration was measured, with acarbose as a reference. The highest dose was most effective. [75]

Aerial parts methanolic extract had blood glucose lowering effects in normal and STZ-nicotinamide-induced diabetic rats: oral administration, 200 mg/kg/day, 21 days. [76]

A comparative study of α -glucosidase inhibition by leaves EO of *S. fruticosa* and *S. officinalis*. For high tested concentrations (50, 75%) of EO, both species had similar activities, despite the fact their chemical compositions are different, qualitatively and quantitatively. For lower concentrations (5, 10, 20%), EO of *S. officinalis* was more active. Acarbose was reference in this study. [77]

A comparative study of α -glucosidase and α -amylase inhibition activities by aerial parts extract of organically cultivated *S. fruticosa* and *S. officinalis*. For low tested concentrations, *S. fruticosa* was much more active, especially for α -amylase inhibition. For higher concentrations, *S. fruticosa* was also more active. Acarbose was reference in this study. [78]

Sinapis alba

Diet containing 10% seeds powder had blood glucose lowering effect in STZ-induced diabetic rats. Normal rats were the control group. [79]




Fig.1. Prevalence of DM-related complications [9]




Fig. 2. Major causes of insulin resistance and major diseases resulting from it [15]




Fig. 3. Antidiabetic agents from *F. vulgare* [33,46]




Fig. 4. Acids with highest concentrations in leaves and roots of *F. vulgare* [52]




Fig. 5. Stigmasterol from *n*-hexane and methanolic extracts of *G. tournefortii* [61]




Fig. 6. Natural products with GLUT4 translocation activity from *G. tournefortii* [66]




Fig. 7. Major components of leaves EO of *M. syriaca* [67]




Fig. 8. Major components of aerial parts volatile oil of *M. fruticososa* [71]




Fig. 9. New phenolics from aqueous methanolic extract of *S. fruticososa* [73]

The reviewed region of Eastern Mediterranean includes several climate zones, and consequently, diverse biome plant groups. But it is very well known that the Deca-Plants grow in moderately rainy areas and mild temperatures. This can explain their absence from herbalist use in dry, desert regions such as Oman and Saudi Arabia, even though Oman has some tropical climate in the mountainous regions [81,82].

In recent years, the world was under the COVID-19 pandemic, which worsened the situation of diabetes mellitus (DM), and it was found that the Mediterranean diet was one of the successful nutritional methods to ease this complication [83]. In this sense, the search for antidiabetic agents in foods and traditional medicines is very interesting, keeping in mind that there are more than 60 FDA-approved synthetic drugs [84]. Among these synthetic drugs, metformin is the most used [85]. The structure of this compound is shown in Fig. 10, and based on that, it is easy to understand the prevalence of its adverse effects, especially iron deficiency and anemia [86,87].

In section 3, we cited only two publications about the antidiabetic activity of *A. palaestinum* [26,27]. Both studies reported that polar extracts,

aqueous, ethanolic and 50% aqueous ethanolic; had low activity, while *n*-hexane extract had significantly α -amylase inhibition activity ($IC_{50} = 25.34 \mu\text{g/mL}$). Based on the chemical composition of the plant [88], we find these reports notably interesting. This composition includes several polar compounds, that will be easily extracted with polar solvents, and have published, well known antidiabetic activity. One of the most important of these compounds is vicianin-2, shown in Fig. 11. [89,90].

The number of published reports about the antidiabetic activity of *C. pumilum* is like that of *A. palaestinum*: two [28,29]. We find this scarcity of publications no less unexpected. *C. pumilum* is rich with potentially antidiabetic natural products. Among these, the structurally close related phenolic acids: chicoric, chlorogenic and caffeic (Fig. 12). [91,92,93].

Each one of these acids was reported for antidiabetic activity in several publication, and we will cite two of them here [94,95].

In the second article of this series, anti-inflammatory activity of the D-P, we highlighted the fact that *Cyclamen pesicum* was not published for this activity, despite the fact that it contains anti-inflammatory agents [96]. This

“puzzle” continues here as well: no published article about antidiabetic activity of *C. pesicum*, even though it contains NPs with published such activity. For this discussion we chose only three compounds contained in this plant with significant concentrations: β -carotene [97], farnesol [98] and peonidin 3-O-glucoside [99] (Fig. 13).

β -Carotene was published for antidiabetic activity in many research and review articles. For example, it had clear positive effect in STZ-induced diabetic rats [100]. Farnesol had similar effect in STZ-induced diabetic mice [101], and peonidin 3-O-glucoside inhibited α -amylase [102].

Among the D-P, *Foeniculum vulgare* is the published for its antidiabetic activity [30-57]. This can be easily understood based on its chemical composition that includes many NPs with notable antidiabetic activity, such as caffeic acid [94,103], and chlorogenic acid [104]. But among these twenty-eight reports [30-57], we found a few that are attention drawing like the work of D.T. Al-Aridhi [35]. In this article, authors reported interestingly very low active doses of leaves extract that treated alloxan-induced diabetic mice: 1.2, 2.2 mg/kg, 30 days. In addition, this report clearly contradicts the report of Y. Hilmi and her colleagues, that found the ethanolic toxic [36].




Fig. 10. Structure of metformin




Fig. 11. Structure of vicenin-2, a NP with notable antidiabetic activity [89,90]




Fig. 12. Three phenolic acids from *C. pumilum* with antidiabetic activity [91-93]




Fig. 13. NPs with published antidiabetic activity from *C. persicum* [100-102]

Inhibition of α -amylase is one of the most important and indicative antidiabetic activities [105]. A. Bashkin and his colleagues found that two of the D-P have very weak α -amylase inhibition: *F. vulgare* and *G. tournefortii* [106]. This contradicts previous reports for *F. vulgare* [47,52,57], and even more for *G. tournefortii* [65], where M. Keskin and her colleagues reported very high activity of this enzyme inhibition: $IC_{50} = 4.22$ mg/mL. And the last contradiction for *F. vulgare*, we find in the report of F. Amir Khanloo and her colleagues that the essential oil of this plant could not improve insulin resistance in women with polycystic ovary syndrome, PCOs

[107]. This contradicts previously cited reports [31,38], but can be understood based on model difference and the fact that women were under hypocaloric diet.

S. Kadan and his colleagues prepared the methanolic extract of the aerial parts *G. tournefortii* and fractionized it in two studies and found that the extract and some of its components have glucose transporter 4 (GLUT4) translocation activity [61,66]. Two of these compounds, *myo*-inositol and D-pinitol (Fig. 6), were published for their notable insulin regulation [108,109].

In the first study [61], the researchers prepared the *n*-hexane extract as well, and they reported that stigmasterol (Fig. 5). was the only compound present in both extracts. So, the GLUT4 translocation activity of *G. tournefortii* can also be referred to this compound since it is known to have this property from other studies [110]. In fact, analyzing *G. tournefortii* for phytosterols leads to the discovery that six of them are contained in this seed EO of this plant with significant concentrations [111]. Stigmasterol is only second (11.7%) to β -sitosterol (35.25%, of total phytosterols content, Fig. 14). This phytosterol was published for strong antidiabetic activity [112].

The fatty acids fraction of this oil contains two compounds with very high concentrations: oleic acid (28%) and linoleic acid (54.6%, of total fatty acids content, Fig. 14). These acids were also published for significant antidiabetic activities [113,114].

Searching for literature of antidiabetic activity of *Majorana syriaca* (Syn. *Orgianum syriacum*) resulted only one reliable article of F.U. Afifi and her colleagues [67]. It is no wonder then, that J. Mesmar and his colleagues did not mention this property in their comprehensive review article of the phytochemistry and the pharmacology of this plant [115]. But in the article of F.U. Afifi and her colleagues, the major two NPs are γ -terpinene and *p*-cymene, 31.6%, 39.1%, respectively (Fig. 7), while in the work of M. Farhat and her colleagues, carvacrol (Fig.15). is way exceeding the other two compounds [116]. This difference has special importance since carvacrol is known for its antidiabetic activity, among many other [117].

We will conclude the discussion of *M. syriaca* with the report of A.R. Al-Assi and his colleagues [118]. They fed rats with 100 and 400 mg/kg of aerial parts "whole extract", which is a

combination of extracts prepared by using three solvents. The report that they obtained "significant decrease" in blood glucose. But examining the results (graph on the second page of the publication) shows a minor decrease that can be included in error range.

The number of published studies about the antidiabetic activity of *Malva sylvestris* is surprisingly low, four [27,68-70]. But attempts to utilize this plant to treat DM are performed in indirect ways. U. Koca-Caliskan and her colleagues prepared copper nanoflowers using methanolic extract of *M. sylvestris* as a reductant [119]. These nanoflowers successfully inhibited α -amylase and α -glucosidase.

In our thorough search for published articles about the antidiabetic activity of *M. sylvestris* we came upon the work of A. Ben Saad and his colleagues [120]. In this work, the research group did not study the antidiabetic activity of *M. sylvestris*, but in the introduction of their article they wrote that the plant has this property, and they cited the work of F.Z. Sabri and her colleagues [121]. So, we found the cited article to discover for our disappointment that it does not mention antidiabetic or any related activity.

Micromeria fruticosa is one of the most important plants in the reviewed region in terms of herbal teas, decoctions, and infusions, that are used for numerous medicinal and nutritional purposes. So, we found it not clear that there was just one report of the antidiabetic activity of this plant [71]. Moreover, I. Telci and M. Ceylan reported the chemical composition of the EO this plant [122], and I.M. Abu-Reidah and his colleagues reported the phenolics composition [123]. In the first report, linalool was the compound with the highest concentration, and in the second report, quercetin was one of the major components (Fig. 16).




Fig. 14. NPs with published antidiabetic activity from *G. tournefortii* [112-114]




Fig. 15. Carvacrol: antidiabetic agent from the leaves EO of *M. syriaca* [116,117]




Fig. 16. Linalool and quercetin, NPs with antidiabetic activity in *M. fruticosa* [122-125]

Linalool was published for antidiabetic activity in a few articles [124], while quercetin was published for the same activity in many research and review articles [125].

The importance of *Salvia fruticosa* for peoples of Eastern Mediterranean region goes beyond being a medicinal nutritional plant, it is a symbol. The presentation and discussion of this topic is out of this article. So, in our search for publications of *S. fruticosa* antidiabetic activity, we found the review article of N.G. Etsassala and his colleagues [126]. They cited two publications. First, the research article of A. Al-Aboudi and F.U. Affi, which we also cited [19]. Second, the review article of R. Mahdizadeh and his colleagues [127]. This article cites a chapter in a book (31 in their reference list) that we will not cite here and the article of M.R. Loizzo and her colleagues (34 in their reference list) [128]. This article does not report any research of *S. fruticosa* antidiabetic activity but mentions the work of M. Perfumi *et al.* that we cited as well [72]. Interestingly, in the article of R. Mahdizadeh and his colleagues, we

found the citation of the work of M.F. Azevedo and her colleagues, that used *S. fruticosa* tea and one of its major components, rosmarinic acid (Fig. 17), to treat STZ-induced diabetic rats [129]. They also that both materials modulate levels of intestinal Na⁺/glucose cotransporter-1 (SGLT1).

A single report was found about antidiabetic activity of *Sinapis alba*: diabetic rats (STZ-induced) were fed with food containing 10% seed powder, resulted lowering blood glucose [79]. And when analyzing the chemical composition of these seed oil [130] or the green, dried aerial parts (that clearly depends on harvest time) [131], we discover plenty of NPs that have published antidiabetic activity. Four of them are presented in Fig. 18 [132-136].

To conclude the discussion section, we present the structures (Fig.19). of the compounds that were used to induce diabetes and were mentioned in this article, as well as the structures of the compounds that were used as references for treatment.

5. SELECTED WILD EDIBLE NON-DECA-PLANTS WITH ANTIDIABETIC ACTIVITY

Screening the top medicinal plants used for treatment of DM in the reviewed region, reveals the fact that most of them are either domesticated or inedible [16,25]. So, in the

search of wild, edible, Non-Deca-Plant for treatment of DM, we will list the most important four, shown in **Table 2**.

Finally, it is important to indicate that none of the D-P is a tree, while of the mentioned above non-D-P plants are trees: *Crataegus aronia* and *Zizyphus spina-christi*.




Fig. 17. Rosmarinic acid, NP with antidiabetic activity in *S. fruticosa* [129]





Fig. 18. NPs with antidiabetic activity in *S. alba* [132-136]

Table 2. Published antidiabetic activities of the Non-D-P in eastern mediterranean region

Testing Method, Results and Reference/s
<i>Crataegus aronia</i> Fruits pulp was extracted with water and orally administered to STZ-induced diabetic rats. Blood glucose concentration was reduced, and memory was improved. [137] Leaves were extracted with methanol and orally administered to STZ-induced diabetic rats. Positive effects were measured by several biomarkers including reduction of blood glucose concentration. [138]
<i>Portulaca oleracea</i> Aerial parts aqueous extract was supplemented to alloxan-induced diabetic rats resulting in increase of β -cells production and decrease of blood glucose concentration. [139] Aerial parts were extracted with 70% aqueous ethanol and the extract was orally fed to STZ-induced diabetic male rats. Blood glucose concentration was decreased, while sperm count and sex hormones concentrations were increased. [140]
<i>Silybum marianum</i> Seeds aqueous extract was supplemented to diabetic humans for four months. Results were measured by several biomarkers including blood glucose, glycosylated hemoglobin, insulin, total cholesterol. Positive effects were observed. [141] Seeds powder was defatted with petroleum ether and extracted with 95% aqueous ethanol. Extract had strong α -glucosidase inhibition activity. [142]
<i>Zizyphus spina-christi</i> Leaves 70% aqueous ethanolic extract was prepared and orally administered to STZ-induced diabetic rats. Positive effects were recorded in blood glucose and insulin concentrations. [143] Fruits were extracted with 80% aqueous ethanol. <i>In vitro</i> , the extract had notable α -amylase and α -glucosidase inhibition activities. <i>In vivo</i> , the extract was orally supplemented to alloxan-induced diabetic rats, resulting significant decrease of blood glucose concentration. [144]

**Fig. 19. Chemicals that induce diabetes (red) and those for treatment/reference (green)**

6. CONCLUSIONS

- 1) Most of the Deca-Plants have weak to moderate antidiabetic activity, according to published research so far.
- 2) *Foeniculum vulgare* is reported for excellent antidiabetic activity and should be used more frequently in homeopathy and alternative medicine.
- 3) Some of the D-P like *Cyclamen persicum* and *Sinapis alba* should be studied more, since they contain natural products with proven antidiabetic activity.
- 4) The chemical compositions of some of the D-P are not completely known, and there is a need to reveal and study them.
- 5) Accuracy of scientific reporting and citation are vital for scientific research.

CONSENT AND ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Author has declared that they have no known competing financial interests or non-financial interests or personal relationships that could have appeared to influence the work reported in this paper.

REFERENCES

1. World Health Organization (WHO): The top 10 cause of death. [WHO]; 2020.
2. Dupras TL, Williams LJ, Willems H, Peeters C, Pathological skeletal remains from ancient Egypt: the earliest case of diabetes mellitus?. *Pract. Diabetes Int.* 2010;27:358-363. DOI: 10.1002/pdi.1523
3. GBD 2021 Diabetes collaborators. Global, regional, and national burden of diabetes from 1990 to 2021, with projections of prevalence to 2050: A systematic analysis for the Global Burden of Disease Study 2021. *Lancet.* 2023; 402:203-234. DOI: 10.1016/S0140-6736(23)01301-6
4. Zhang P, Zhang X, Brown J, Vistisen D, Sicree R, Shaw J, Nichols G, Global healthcare expenditure on diabetes for 2010 and 2030. *Diabetes Res. Clin. Pract.* 2010;87:293-301. DOI: 10.1016/j.diabres.2010.01.026
5. American diabetes association; 2022.
6. Rodríguez JE, Campbell KM. Racial and ethnic disparities in prevalence and care of patients with type 2 diabetes. *Clin. Diabetes.* 2017;35:66-70. DOI: 10.2337/cd15-0048
7. Meo SA, Usmani AM, Qalbani E, Prevalence of type 2 diabetes in the Arab world: impact of GDP and energy consumption. *Eur. Rev. Med. Pharmacol. Sci.* [ResearchGate] 2017;21:1303-1312.
8. Jaffe A, Giveon S, Wulffhart L, Oberman B, Baidousi M, Ziv A, Kalter-Leibovici O, Adult Arabs have higher risk for diabetes mellitus than Jews in Israel. *PLoS ONE.* 2017;12:e0176661. DOI: 10.1371/journal.pone.0176661
9. Deshpande AD, Harris-Hayes M, Schootman M, Epidemiology of diabetes and diabetes-related complications. *Phys. Ther.* 2008;88:1254-1264. DOI: 10.2522/ptj.20080020
10. Tomic D, Shaw JE, Magliano DJ. The burden and risks of emerging complications of diabetes mellitus. *Nat. Rev. Endocrinol.* 2022;18:525-539. DOI: 10.1038/s41574-022-00690-7
11. Hill Golden S, Lazo M, Carnethon M, Bertoni AG, Schreiner PJ, Diez Roux AV, Lee HB, Lyketsos C. Examining a bidirectional association between depressive symptoms and diabetes. *JAMA.* 2008;299:2751-2759. DOI: 10.1001/jama.299.23.2751
12. Li M, Chi X, Wang Y, Setrerrahmane S, Xie W, Xu H, Trends in insulin resistance: insights into mechanisms and therapeutic strategy. *Signal Transduct. Target. Ther.* 2022;7:216. DOI: 10.1038/s41392-022-01073-0
13. Lee SH, Park SY, Choi CS. Insulin resistance: From mechanisms to therapeutic strategies. *Diabetes Metab. J.* 2022;46:15-37. DOI: 10.4093/dmj.2021.0280
14. Zhao X, An X, Yang C, Sun W, Ji H, Lian F. The crucial role and mechanism of insulin resistance in metabolic disease. *Front. Endocrinol.* 2023;14:1149239. DOI: 10.3389/fendo.2023.1149239
15. Rader DJ. Effect of insulin resistance, dyslipidemia, and intra-abdominal adiposity on the development of cardiovascular disease and diabetes mellitus. *Am. J. Med.* 2007;120:S12-8. DOI: 10.1016/j.amjmed.2007.01.003
16. Przeor M, Some common medicinal plants with antidiabetic activity, Known and

- Available in Europe (A Mini-Review).
Pharmaceuticals. 2022;15:65.
DOI: 10.3390/ph15010065
17. Abu-Odeh AM, Talib WH, Middle east medicinal plants in the treatment of diabetes: A review. *Molecules*. 2021; 26:742.
DOI: 10.3390/molecules26030742
 18. Otoom SA, Al-Safi SA, Kerem ZK, Alkofahi A. The use of medicinal herbs by diabetic Jordanian patients. *J. Herb. Pharmacother.* [ResearchGate]. 2006;6:31-41.
 19. Al-Aboudi A, Afifi FU. Plants used for the treatment of diabetes in Jordan: A review of scientific evidence. *Pharm. Biol.* 2011;49:221-239.
DOI: 10.3109/13880209.2010.501802
 20. Afifi-Yazar FU, Kasabri V, Abu-Dahab R. Medicinal plants from Jordan in the treatment of diabetes: traditional uses vs. *in vitro* and *in vivo* evaluations - part 2. *Planta Med.* 2011;77:1210-1220.
DOI: 10.1055/s-0031-1279983
 21. Afifi FU, Kasabri V. Pharmacological and phytochemical appraisal of selected medicinal plants from Jordan with claimed antidiabetic activities. *Sci. Pharm.* 2013;81:889-932.
DOI: 10.3797/scipharm.1212-20
 22. Issa R, Khatlaba A, Alkarem TA, Altameemi O, The use of antidiabetic herbal remedies by Jordanian herbalist: A comparison of folkloric practice vs. evidence-based pharmacology. *Jordan J. Pharm. Sci.* [ResearchGate]. 2019;12:23-37.
 23. Jeambey Z, Johns T, Talhouk S, Batal M. Perceived health and medicinal properties of six species of wild edible plants in north-east Lebanon. *Public Health Nutr.* 2009;12:1902-1911.
DOI: 10.1017/S1368980009004832
 24. Yaniv Z, Dafni A, Friedman J, Palevitch D. Plants used for the treatment of diabetes in Israel. *J. Ethnopharmacol.* 1987;19:145-151.
DOI: 10.1016/0378-8741(87)90038-9
 25. Ali-Shtayeh MS, Jamous RM, Jamous RM. Complementary and alternative medicine use amongst Palestinian diabetic patients. *Complement. Ther. Clin. Pract.* 2012;18:16-21.
DOI: 10.1016/j.ctcp.2011.09.001
 26. Afifi FU, Kasabri V, Litescu SC, Abaza IM. *In vitro* and *in vivo* comparison of the biological activities of two traditionally and widely used arum species from Jordan: *Arum dioscoridis* Sibth & Sm. and *Arum palaestinum* Boiss. *Nat. Prod. Res.* 2015;30:1777-1786.
DOI: 10.1080/14786419.2015.1072713
 27. Hawash M, Jaradat N, Elaraj J, Hamdan A, Lebdeh SA, Halawa T. Evaluation of the hypoglycemic effect of seven wild folkloric edible plants from Palestine. *J. Complement. Integr. Med.* 2019; 17:20190032.
DOI: 10.1515/jcim-2019-0032
 28. Kamel ZH, Daw I, Marzouk M. Effect of *Cichorium endivia* leaves on some biochemical parameters in streptozotocin-induced diabetic rats. *Aus. J. Basic App. Sci.* 2011;5:387-396.
[ResearchGate]
 29. Alkofahi AS, Abdul-Razzak KK, Alzoubi KH, Khabour OF. Screening of the anti-hyperglycemic activity of some medicinal plants of Jordan. *Pak. J. Pharm. Sci.* 2017;30:907-913.
[ResearchGate]
 30. Özbek H, Öztürk M, Bayram I, Uğraş S, Çitoğlu GS. Hypoglycemic and hepatoprotective effects of *Foeniculum vulgare* miller seed fixed oil extract in mice and rats. *East. J. Med.* 2003;8:35-40.
[ResearchGate]
 31. Dongare VR, Arvindekar AU, Magadum CS. Hypoglycemic effect of *Foeniculum vulgare* Mill. fruit on dexamethasone induced insulin resistance rats. *Res. J. Pharmacog. Phytochem.* [Google Scholar]. 2010;2:163-165.
 32. Abou El-Soud N, El-Laithy N, El-Saeed G, Wahby MS, Khalil M, Morsy F, Shaffie N. Antidiabetic activities of *Foeniculum Vulgare* Mill. Essential oil in streptozotocin-induced diabetic rats. *Maced. J. Med. Sci.* 2011;4:139-146.
DOI: 10.3889/MJMS.1957-5773.2011.0173
 33. Dongare V, Kulkarni C, Kondawar M, Magdum C, Haldavnekar V, Arvindekar A. Inhibition of aldose reductase and anti-cataract action of trans-anethole isolated from *Foeniculum vulgare* Mill. fruits. *Food Chem.* 2012;132:385-390.
DOI: 10.1016/j.foodchem.2011.11.005
 34. Anitha T, Balakumar C, Ilango KB, Jose CB, Vetrivel D. Antidiabetic activity of the aqueous extracts of *Foeniculum vulgare* on streptozotocin-induced diabetic rats. *Int. J. Adv. Pharm. Biol. Chem.* 2014;3:487-494.
[Google Scholar]
 35. Al-Aridhi DT. The effect of *Foeniculum Vulgare* Alcoholic extract on some metabolic changes in liver and kidneys of

- alloxan diabetic mice. Iraqi J. Med. Sci. 2014;12:119-125. [Google Scholar]
36. Hilmi Y, Abushama MF, Abdalgadir H, Khalid A, Khalid H. A study of antioxidant activity, enzymatic inhibition and *in vitro* toxicity of selected traditional Sudanese plants with anti-diabetic potential. BMC Complement. Altern. Med. 2014;14:149. DOI: 10.1186/1472-6882-14-149
 37. Mhaidat NM, Abu-zaiton AS, Alzoubi KH, Alzoubi W, Alazab RS. Antihyperglycemic properties of *Foeniculum vulgare* extract in streptozocin-induced diabetes in rats. Int. J. Pharmacol. 2015;11:72-75. DOI: 10.3923/ijp.2015.72.75
 38. Pari L, Sheikh BA. Antihyperglycemic effect of *trans*-anethole in streptozotocin induced diabetic rats with special reference to glycoprotein components. Int. J. Adv. Res. Biol. Sci. 2015;2:28-34. [Google Scholar]
 39. Mostafa DM, Abd El-Alim SH, Asfour MH, Al-Okbi SY, Mohamed DA, Awad G. Transdermal nanoemulsions of *Foeniculum vulgare* Mill. essential oil: Preparation, characterization and evaluation of antidiabetic potential. J. Drug Deliv. Sci. Technol. 2015;29:99-106. DOI: 10.1016/j.jddst.2015.06.021
 40. Parsaeyan N, The effect of *Foeniculum vulgare* (fennel) extract on lipid profile, lipid peroxidation and liver enzymes of diabetic rat. Iran. J. Diabetes Obes. 2016;8:24-29. [Google Scholar]
 41. Zaahkook SA, Ibrahim DF, Elarabi BE. Hypolipidemic studies on Egyptian propolis and *Foeniculum Vulgare* on alloxan induced diabetic rats. Int. J. Adv. Res. 2016;4:1180-1186. [Journal Website]
 42. Zaahkook SA, Ibrahim DF, Elarabi BE. Antioxidants and hypolipidemic studies on Egyptian propolis and *Foeniculum Vulgare* on alloxan induced diabetic rats. Int. J. Anim. Biol. 2016;2:1-10. [Journal Website]
 43. El Araby B, Ibrahim DF, Zahkook SA. Effect of *Foeniculum vulgare* and propolis on liver in alloxan diabetic rats. Adv. Biol. Res. 2017;11:311-318. DOI: 10.5829/idosi.abr.2017.311.318
 44. Osman NN, Jambi EJ, Aseri NH. Assessment of antidiabetic and antioxidant activities of *Cassia angustifolia* and *Foeniculum vulgare* in diabetic rats. Int. J. Pharm. Res. Allied Sci. 2017;6:149-162. [ResearchGate]
 45. Basha SG, Ahamed MF, Farnaaz S. Evaluation of *Foeniculum vulgare* and *Tagetes erecta* for Synergic anti diabetic activity in streptozotocin induced diabetic rats. World J. Pharm. Res. 2017;6:1401-1428. DOI: 10.20959/wjpr20178-9000
 46. Özbek H, Sever Yilmaz B. Anti-inflammatory and hypoglycemic activities of alpha-pinene. Acta Pharm. Sci. 2017;55:7-14. DOI: 10.23893/1307-2080.APS.05522
 47. Godavari A, Amutha K, Moorthi NM. *In-vitro* hypoglycemic effect of *Foeniculum vulgare* Mill. Seeds on the carbohydrate hydrolysing enzymes, α -amylase and α -glucosidase. Int. J. Pharm. Sci. Res. 2018;9:4441-4445. DOI: 10.13040/IJPSR.0975-8232.9(10).4441-45
 48. Zulifqar S. Influence of *Foeniculum vulgare* Mill in the Management of Hyperglycemia. Int. J. Innov. Sci. Res. Technol. 2019;4:1117-1122. [Google Scholar]
 49. Setyaningsih EP, Saputri FC, Mun'im A. The antidiabetic effectivity of Indonesian plants extracts via DPP-IV inhibitory mechanism. J. Young Pharm. 2019;11:161-164. DOI: 10.5530/jyp.2019.11.34
 50. El-Ouady F, Lahrach N, Ajebli M, Haidani AE, Eddouks M. Antihyperglycemic effect of the aqueous extract of *Foeniculum vulgare* in normal and streptozotocin-induced diabetic rats. Cardiovasc. Hematol Disord. Drug Targets. 2020;20:54-63. DOI: 10.2174/1871525717666190612121516
 51. Samadi-Noshahr Z, Hadjzadeh MA, Moradi-Marjaneh R, Khajavi-Rad A. The hepatoprotective effects of fennel seeds extract and *trans*-Anethole in streptozotocin-induced liver injury in rats. Food Sci. Nutr. 2021;9:1121-1131. DOI: 10.1002/fsn3.2090
 52. Sayah K, El Omari N, Kharbach M, Bouyahya A, Kamal R, Marmouzi I, Cherrah Y, Faouzi ME. Comparative study of leaf and rootstock aqueous extracts of *Foeniculum vulgare* on chemical profile and *In vitro* antioxidant and antihyperglycemic activities. Adv. Pharmacol. Pharm. Sci. 2020; 8852570. DOI: 10.1155/2020/8852570
 53. Shabbir R, Imran M, Arshad S, UI-Hassan F. Assessment and effectiveness of fennel

- powder extract based cookies in hyperglycemic subjects. Open J. Nutr. Food Sci. 2020;2:1008. [Journal Website]
54. Rohman F, Putra WE. The bioinformatics perspective of *Foeniculum vulgare* fruit's bioactive compounds as natural anti-hyperglycemic against alpha-glucosidase. Biodiversitas. 2021;22. DOI: 10.13057/biodiv/d220111
 55. Servi H, Şen A, Yildirim Servi E, Doğan A. Chemical composition and biological activities of essential oils of *Foeniculum vulgare* Mill. and *Daucus carota* L. growing wild in Turkey. J. Res. Pharm. 2021;25:142-152. DOI: 10.29228/jrp.5
 56. Osman NN, Al-Ahmadi AM, Ghazwani AH, Alhoraibi HM, Alanbari KH, Backer WS. Role of senna, *Cassia angustifolia* and fennel, *Foeniculum vulgare* in ameliorating nephropathy in diabetic rats. Biosci. Biotechnol. Res. Commun. 2022;15:208-216. DOI: 10.21786/bbrc/15.1.32
 57. Kumar PY, Subramanayaan M. Inhibitory Effect of *Foeniculum vulgare* leaf extract on alpha-glucosidase and alphaamylase activity. Acta Sci. Med. Sci. 2022;6:26-31. DOI: 10.31080/ASMS.2022.06.1115
 58. Hamad NS, Hasan HG. Inhibitory effect of *Gundelia* extract on urinary α -amylase activity of type-I diabetes mellitus. Zanco J. Med. Sci. 2010;14:1-6. [Google Scholar]
 59. Azeez OH, Kheder AE. Effect of *Gundelia tournefortii* on some biochemical parameters in dexamethasone-induced hyperglycemic and hyperlipidemic mice. Iraqi J. Vet. Sci. 2012;26:73-79. [ResearchGate]
 60. Alimoradi M, Jalili C, Kakeh-Baraei S, Tajehmiri A, Khodarahmi R. Effects of aqueous extract of gunnera (*Gundelia tournefortii* L.) on the blood serum sugar levels and changes in the streptozotocin-induced diabetic pancreatic tissue of rat. Int. J. Sci. Stud. 2017;5:186-191. DOI: 10.17354/ijssl/2017/123
 61. Kadan S, Sasson Y, Saad B, Zaid H. *Gundelia tournefortii* antidiabetic efficacy: Chemical composition and GLUT4 translocation. Evid. Based Complement. Alternat. Med. 2018; 8294320. DOI: 10.1155/2018/8294320
 62. Mohammadi G, Zangeneh MM, Rashidi K, Zangeneh A. Evaluation of nephroprotective and antidiabetic effects of *Gundelia tournefortii* aqueous extract on diabetic nephropathy in male mice. Res. J. Pharmacog. 2018;5:65-73. DOI: 10.22127/rjp.2018.69223
 63. Goorani S, Koochi MK, Seydi N, Zangeneh A, Zangeneh MM. Protection of alloxan monohydrate-induced testicular toxicity by *Gundelia tournefortii* aerial parts aqueous extract in male mice. Iranian J. Pharmacol. Ther. 2018;16:1-9. [ResearchGate]
 64. Asareh Z, Bahramikia S. Inhibitory effect of the effective fraction of *Gundelia tournefortii* L and several anti-diabetic drugs on the aldose reductase activity isolated from cow lens eye: A comparative study. J. Anim. Res. (Iran. J. Biol.). 2019;32:198-209. [Google Scholar]
 65. Keskin M, Kaya G, Keskin Ş. *Gundelia Tournefortii* L. (Kenger): Determination of *in vitro* Antidiabetic Activities. Prog. Nutr. 2021;23:e2021163. DOI: 10.23751/pn.v23i4.11079
 66. Kadan S, Melamed S, Bernalid S, Tietel Z, Sasson Y, Zaid H. *Gundelia tournefortii*: Fractionation, Chemical composition and GLUT4 translocation enhancement in muscle cell line. Molecules. 2021;26:3785. DOI: 10.3390/molecules26133785
 67. Afifi FU, Kasabri V, Beltran S, Abuhammad A, Abaza IF, Ganado O, Al-Gabbiesh AH. Comparison of different methods in determination of essential oil composition of *Origaum syriacum* L. from Jordan and its modulation of pancreatic enzymes. Rev. Roum. Chim. 2017;62:15-21. [ResearchGate]
 68. Loizzo MR, Pugliese A, Bonesi M, Tenuta MC, Menichini F, Xiao J, Tundis R. Edible flowers: A rich source of phytochemicals with antioxidant and hypoglycemic properties. J. Agric. Food Chem. 2016;64:2467-2474. DOI: 10.1021/acs.jafc.5b03092
 69. Bilen S, Filogh AM, Ali AB, Kenanoğlu ON, Zoral MA. Effect of common mallow (*Malva sylvestris*) dietary supplementation on growth performance, digestive enzyme activities, haematological and immune responses of common carp (*Cyprinus carpio*). Aquac. Int. 2020;28:73-84. DOI: 10.1007/s10499-019-00444-9
 70. Ombra MN, Nazzaro F, Fratianni F. Pasta fortification with leaves of edible wild plants to lower the p glycaemic index of handmade fresh noodles. Recent Progress Nutr. 2023;3:1-21.

- DOI: 10.21926/rpn.2302008
71. Salameh N, Shraim N, Jaradat N. Chemical composition and enzymatic screening of *Micromeria fruticosa serpyllifolia* volatile oils collected from three different regions of West Bank, Palestine. *Biomed. Res. Int.* 2018; 6536919.
DOI: 10.1155/2018/6536919
 72. Perfumi M, Arnold N, Tacconi R. Hypoglycemic activity of *Salvia fruticosa* Mill. from Cyprus. *J. Ethnopharmacol.* 1991;34:135-140.
DOI: 10.1016/0378-8741(91)90030-h
 73. Moharram FA, Mahmoud II, Mahmoud MR, Sabry SA. Polyphenolic profile and biological study of *Salvia fruticosa*. *Nat. Prod. Commun.* 2006;1:745-750
DOI: 10.1177/1934578X0600100909
 74. Raafat K, Boukhary R, Aboul-Ela M, El-Lakany A. Endogenous lebanese plants treating diabetes and related complications. *Nat. Prod. Chem. Res.* 2013;1:1000112.
DOI: 10.4172/2329-6836.1000112
 75. Bassil M, Daher CF, Mroueh M, Zeeni N, *Salvia libanotica* improves glycemia and serum lipid profile in rats fed a high fat diet. *BMC Complement. Altern. Med.* 2015;15:384.
DOI: 10.1186/s12906-015-0917-8
 76. Cam ME, Yildiz S, Ertas B, Acar AE, Taskin T, Kabasakal L. Antidiabetic effects of *Salvia triloba* and *Thymus praecox* subsp. *skorpilii* var. *skorpilii* in a rat model of streptozotocin/nicotinamide-induced diabetes. *Marmara Pharm. J.* 2017;21:818-827.
DOI: 10.12991/mpj.2017.8
 77. Chehade S, Kobeissy M, Kanaan H, Haddad M. Comparison between the chemical compositions and the *in-vitro* antidiabetic and anti-inflammatory activities of *Salvia Libanotica* and *Salvia Officinalis* leaves essential oils. *Eur. J. Pharm. Med. Res.* 2022;93:34-43.
[ResearchGate]
 78. Özüpek B, Pekacar S, Orhan DD. Evaluation of phytochemical contents and biological activities of *Salvia officinalis* and *Salvia triloba* grown with organic farming. *Fabad J. Pharm. Sci.* 2023;48:125-138.
DOI: 10.55262/fabadezczacilik.1175781
 79. Amer HM, Aly TA, Tobgy KM, Abdallah MM, El-Shahat N. Lipid metabolism in diabetic rats as affected by canola and mustard seed sprouts. *Arab Univ. J. Agric. Sci.* 2019;27:649-657.
DOI: 10.21608/AJS.2019.43678
 80. Martín-Peláez S, Fito M, Castaner O. Mediterranean diet effects on type 2 diabetes prevention, Disease progression, and related mechanisms. A Review. *Nutrients.* 2020;12:2236.
DOI: 10.3390/nu12082236
 81. Divakar MC, Al-Siyabi A, Varghese SS, Rubaie MA. The practice of ethnomedicine in the northern and southern provinces of Oman. *Oman Med. J.* 2016;31: 245-252.
DOI: 10.5001/omj.2016.49
 82. Alsanad S, Aboushanab T, Khalil M, Alkhamees OA. A descriptive review of the prevalence and usage of traditional and complementary medicine among saudi diabetic patients. *Scientifica (Cairo).* 2018;6303190.
DOI: 10.1155/2018/6303190
 83. Ochoa Esteban D, Martin-Ridaura C, Berlinches-Zapero C, Ruiz-Fernández D, Sanz-Martín V, Gavira-Izquierdo R, Muñoz-Haba A, March S, Ceinos-Arcones M. Impact of COVID-19 confinement on the health-related habits of people at high risk of type 2 diabetes. *Nutrients.* 2023;15:841.
DOI: 10.3390/nu15040841
 84. Dahlén AD, Dashi G, Maslov I, Attwood MM, Jonsson J, Trukhan V, Schiöth HB. Trends in antidiabetic drug discovery: FDA approved drugs, new drugs in clinical trials and global sales. *Front. Pharmacol.* 2022;12:807548.
DOI: 10.3389/fphar.2021.807548
 85. Hotta N. A new perspective on the biguanide, metformin therapy in type 2 diabetes and lactic acidosis. *J. Diabetes Investig.* 2019;10:906-908.
DOI: 10.1111/jdi.13090
 86. Nasri H, Rafieian-Kopaei M. Metformin: Current knowledge. *J. Res. Med. Sci.* 2014;19:658-664. [ResearchGate]
 87. Wu J, Yang R, Yu H, Qin X, Wu T, Wu Y, Hu Y. Association of metformin use with iron deficiency anemia in Urban Chinese patients with type 2 diabetes. *Nutrients.* 2023;15:3081.
DOI: 10.3390/nu15143081
 88. Abu-Reidah IM, Ali-Shtayeh MS, Jamous RM, Arráez-Román D, Segura-Carretero A. Comprehensive metabolite profiling of *Arum palaestinum* (Araceae) leaves by using liquid chromatography–tandem mass

- spectrometry. Food Res. Int. 2015;70:74-86.
DOI: 10.1016/j.foodres.2015.01.023
89. Islam N, Ishita IJ, Jung HA, Choi JS. Vicenin 2 isolated from *Artemisia capillaris* exhibited potent anti-glycation properties. Food Chem. Toxicol. 2014;69:55-62.
DOI: 10.1016/j.fct.2014.03.042
 90. Ku SK, Bae JS. Vicenin-2 and scolymoside inhibit high-glucose-induced vascular inflammation in vitro and *in vivo*. Can. J. Physiol. Pharmacol. 2015;94:287-295.
DOI: 10.1139/cjpp-2015-0215
 91. Lee J, Scagel CF. Chicoric acid: Chemistry, distribution, and production. Front. Chem. 2013;1:40.
DOI: 10.3389/fchem.2013.00040
 92. Hegazy AK, Ezzat SM, Qasem IB, Ali-Shtayeh MS, Basalah MO, Ali HM, Hatamleh AA. Diversity of active constituents in *Cichorium endivia* and *Cynara cornigera* extracts. Acta Biol. Hung. 2015;66:103-118.
DOI: 10.1556/ABiol.66.2015.1.9
 93. El-Shafey NM, AbdElgawad HR. Antioxidants released from *Cichorium pumilum* Jacq. amendment mitigate salinity stress in maize. Jordan J. Biol. Sci. 2020;13:525-533. [ResearchGate]
 94. Salau VF, Erukainure OL, Ijomone OM, Islam MS. Caffeic acid regulates glucose homeostasis and inhibits purinergic and cholinergic activities while abating oxidative stress and dyslipidaemia in fructose-streptozotocin-induced diabetic rats. J. Pharm. Pharmacol. 2022;74:973-984.
DOI: 10.1093/jpp/rgac021
 95. Ferrare K, Bidet LP, Awwad A, Poucheret P, Cazals G, Lazennec F, Azay-Milhou J, Tournier M, Lajoix AD, Tusch D. Increase in insulin sensitivity by the association of chicoric acid and chlorogenic acid contained in a natural chicoric acid extract (NCRAE) of chicory (*Cichorium intybus* L.) for an antidiabetic effect. J. Ethnopharmacol. 2018;215:241-248.
DOI: 10.1016/j.jep.2017.12.035
 96. Azab A. Top edible wild plants of eastern Mediterranean region Part II: Anti-inflammatory activity. Eur. J. Med. Plants. 2023;34:1-24.
DOI: 10.9734/EJMP/2023/v34i91155
 97. Cornea-Cipcigan M, Bunea A, Bouari CM, Pamfil D, Páll E, Urcan AC, Margaoan R. Anthocyanins and carotenoids characterization in flowers and leaves of cyclamen genotypes linked with bioactivities using multivariate analysis techniques. Antioxidants. 2022;11:1126.
DOI: 10.3390/antiox11061126
 98. Ishizaka H, Yamada H, Sasaki K. Volatile compounds in the flowers of *Cyclamen persicum*, *C. purpurascens* and their hybrids. Sci. Hortic. 2002;94:125-135.
DOI: 10.1016/S0304-4238(01)00362-4
 99. Boase MR, Lewis DH, Davies KM, Marshall GB, Patel D, Schwinn KE, Deroles SC. Isolation and antisense suppression of flavonoid 3', 5'-hydroxylase modifies flower pigments and colour in cyclamen. BMC Plant Biol. 2010;10:107.
DOI: 10.1186/1471-2229-10-107
 100. Nimbalkar V, Joshi U, Shinde S, Pawar G. *In-vivo* and *in-vitro* evaluation of therapeutic potential of β -Carotene in diabetes. J. Diabetes Metab. Disord. 2021;20:1621-1630.
DOI: 10.1007/s40200-021-00912-1
 101. Valdés M, Calzada F, Mendieta-Wejebe JE, Merlín-Lucas V, Velázquez C, Barbosa E. Antihyperglycemic effects of *Annona diversifolia* Safford and its acyclic terpenoids: α -glucosidase and selective SGLT1 inhibitors. Molecules. 2020;25:3361.
DOI: 10.3390/molecules25153361
 102. Sui X, Zhang Y, Zhou W. *In vitro* and *in silico* studies of the inhibition activity of anthocyanins against porcine pancreatic α -amylase. J. Funct. Foods. 2016;21:50-57.
DOI: 10.1016/j.jff.2015.11.042
 103. Oršolic N, Sirovina D, Odeh D, Gajski G, Balta V, Šver L, Jazvinšcak Jembrek M. Efficacy of caffeic acid on diabetes and its complications in the mouse. Molecules. 2021;26:3262.
DOI: 10.3390/molecules26113262
 104. Yan Y, Zhou X, Guo K, Zhou F, Yang H. Use of chlorogenic acid against diabetes mellitus and its complications. J. Immunol. Res. 2020;9680508.
DOI: 10.1155/2020/9680508
 105. Kashtoh H, Baek KH. New insights into the latest advancement in α -amylase inhibitors of plant origin with anti-diabetic effects. Plants. 2023;12:2944.
DOI: 10.3390/plants12162944
 106. Bashkin A, Ghanim M, Abu-Farich B, Rayan M, Miari R, Srouji S, Rayan A, Falah M. Forty-one plant extracts screened for dual antidiabetic and antioxidant functions: Evaluating the types of

- correlation between α -amylase inhibition and free radical scavenging. *Molecules*. 2021;26:317.
DOI: 10.3390/molecules26020317
107. Amir Khanloo F, Esmaeilzadeh S, Mirabi P, Abedini A, Amiri M, Saghebi R, Golsorkhtabaramiri M. Comparison of *Foeniculum Vulgare* versus metformin on insulin resistance and anthropometric indices of women with polycystic ovary, an open-label controlled trial study. *Obes. Med.* 2022;31:100401.
DOI: 10.1016/j.obmed.2022.100401
 108. Antony PJ, Gandhi GR, Stalin A, Balakrishna K, Toppo E, Sivasankaran K, Ignacimuthu S, Al-Dhabi NA. Myoinositol ameliorates high-fat diet and streptozotocin-induced diabetes in rats through promoting insulin receptor signaling. *Biomed. Pharmacother.* 2017;88:1098-1113.
DOI: 10.1016/j.biopha.2017.01.170
 109. Azab A, D-Pinitol – Active natural product from carob with notable insulin regulation. *Nutrients*. 2022;14:1453.
DOI: 10.3390/nu14071453
 110. Wang J, Huang M, Yang J, Ma X, Zheng S, Deng S, Huang Y, Yang X, Zhao P. Anti-diabetic activity of stigmaterol from soybean oil by targeting the GLUT4 glucose transporter. *Food Nutr. Res.* 2017;61:1364117.
DOI: 10.1080/16546628.2017.1364117
 111. Khanzadeh F, Haddad Khodaparast MH, Elhami Rad AH, Rahmani F. Physicochemical properties of *Gundelia tournefortii* L. seed oil. *J. Agric. Sci. Tech.* 2012;14:1535-1542.
[ResearchGate]
 112. Gupta R, Sharma AK, Dobhal MP, Sharma MC, Gupta RS. Antidiabetic and antioxidant potential of β -sitosterol in streptozotocin-induced experimental hyperglycemia. *J. Diabetes*. 2011;3:29-37.
DOI: 10.1111/j.1753-0407.2010.00107.x
 113. López-Gómez C, Santiago-Fernández C, García-Serrano S, García-Escobar E, Gutiérrez-Repiso C, Rodríguez-Díaz C, Ho-Plágaro A, Martín-Reyes F, Garrido-Sánchez L, Valdés S, Rodríguez-Cañete A, Rodríguez-Pacheco F, García-Fuentes E. Oleic acid protects against insulin resistance by regulating the genes related to the pi3k signaling pathway. *J. Clin. Med.* 2020;9:2615.
DOI: 10.3390/jcm9082615
 114. Yoon SY, Ahn D, Hwang JY, Kang MJ, Chung SJ. Linoleic acid exerts antidiabetic effects by inhibiting protein tyrosine phosphatases associated with insulin resistance. *J. Funct. Foods*. 2021;83:104532.
DOI: 10.1016/j.jff.2021.104532
 115. Mesmar J, Abdallah R, Badran A, Maresca M, Baydoun E. *Origanum syriacum* Phytochemistry and Pharmacological Properties: A comprehensive review. *Molecules*. 2022;27:4272.
DOI: 10.3390/molecules27134272
 116. Farhat M, Tóth J, Héthelyi BÉ, Szarka S, Czige S. Analysis of the essential oil compounds of *Origanum syriacum* L. *Eur. Pharm. J.* 2012;59:6-14.
DOI: 10.2478/v10219-012-0020-x
 117. Aljelehaway QH, Maroufi Y, Javid H, Mohammadi MR, Mal Allah OR, Taheri SV, Mohammadzade H. Anticancer, antineurodegenerative, antimicrobial, and antidiabetic activities of carvacrol: recent advances and limitations for effective formulations. *Nano Micro Biosystems*. 2022;2:1-10.
DOI: 10.22034/nmbj.2023.380207.1009
 118. Al-Assi AR, Al-Qirim TM, Abuomair MS, Shahwan MS, Bader AM. Short-term feeding effects of *Origanum syriacum* crude extract on blood constituents in rats. *Int. J. Res. Ayurveda Pharm.* 2017;8:118–120.
DOI: 10.7897/2277-4343.08124
 119. Koca-Caliskan U, Donmez C, Eruygur N, Ayaz F, Altinkaynak C, Ozdemir N. Synthesis and characterization of copper-nanoflowers with the utilization of medicinal plant extracts for enhanced various enzyme inhibitory activities. *Chem. Biodivers.* 2022;19:e202200476.
DOI: 10.1002/cbdv.202200476
 120. Ben Saad A, Rjeibi I, Alimi H, Ncib S, Smida A, Zouari N, Zourgui L. Lithium induced, oxidative stress and related damages in testes and heart in male rats: The protective effects of *Malva sylvestris* extract. *Biomed. Pharmacother.* 2017;86:127-135.
DOI: 10.1016/j.biopha.2016.12.004
 121. Sabri FZ, Belarbi M, Sabri S, Alsayadi Muneer MS. Phytochemical screening and identification of some compounds from mallow. *J. Nat. Prod. Plant Resour.* 2012;2:512-516. [ResearchGate]
 122. Telci I, Ceylan M, Essential oil composition of *Micromeria fruticosa* Druce from

- Turkey. Chem. Nat. Compd. 2007;43:629-631.
DOI: 10.1007/s10600-007-0212-0
123. Abu-Reidah IM, Arráez-Román D, Al-Nuri M, Warad I, Segura-Carretero A. Untargeted metabolite profiling and phytochemical analysis of *Micromeria fruticosa* L. (Lamiaceae) leaves, Food Chem. 2018;279:128-143.
DOI: 10.1016/j.foodchem.2018.11.144
124. More TA, Kulkarni BR, Nalawade ML, Arvindekar AU. Antidiabetic activity of linalool and limonene in streptozotocin-induced diabetic rat: A combinatorial therapy approach. Int. J. Pharm. Pharm. Sci. 2014;6:159-163. [ResearchGate]
125. Dhanya R. Quercetin for managing type 2 diabetes and its complications, an insight into multitarget therapy. Biomed. Pharmacother. 2022;146:112560.
DOI: 10.1016/j.biopha.2021.112560
126. Etsassala NG, Hussein AA, Nchu F. Potential application of some lamiaceae species in the management of diabetes. Plants. 2021;10:279.
DOI: 10.3390/plants10020279
127. Mahdizadeh R, Moein S, Soltani N, Malekzadeh K, Moein M. Study the molecular mechanism of *Salvia species* in prevention of diabetes. Int. J. Pharm. Sci. Res. 2018;9:4512-4521.
DOI: 10.13040/IJPSR.0975-8232.9(11).4512-21
128. Loizzo MR, Saab AM, Tundis R, Menichini F, Bonesi M, Piccolo V, Statti GA, De Cindio B, Houghton PJ, Menichini F. *In vitro* inhibitory activities of plants used in Lebanon traditional medicine against angiotensin converting enzyme (ACE) and digestive enzymes related to diabetes. J. Ethnopharmacol. 2008;119:109-116.
DOI: 10.1016/j.jep.2008.06.003
129. Azevedo MF, Lima CF, Fernandes-Ferreira M, Almeida MJ, Wilson JM, Pereira-Wilson C. Rosmarinic acid, major phenolic constituent of Greek sage herbal tea, modulates rat intestinal SGLT1 levels with effects on blood glucose. Mol. Nutr. Food Res. 2011;55:S15–S25.
DOI:10.1002/mnfr.201000472
130. Peng C, Zhao SQ, Zhang J, Huang GY, Chen LY, Zhao FY. Chemical composition, antimicrobial property and microencapsulation of Mustard (*Sinapis alba*) seed essential oil by complex coacervation. Food Chem. 2014;165:560-568.
DOI: 10.1016/j.foodchem.2014.05.126
131. Sadowska U, Jewiarz K, Kopak M, Dziadek K, Francik R, Kopeć A. Proximate analysis and antioxidant properties of young plants of *Sinapis alba* L. Depend on the time of harvest and variety. Appl. Sci. 2023;13:7980.
DOI: 10.3390/app13137980
132. Liu W, Li K, Zheng M, He L, Chen T. Genipin attenuates diabetic cognitive impairment by reducing lipid accumulation and promoting mitochondrial fusion via FABP4/Mfn1 signaling in microglia. Antioxidants. 2023;12:74.
DOI: 10.3390/antiox12010074
133. Yao DD, Yang L, Wang Y, Liu C, Wei YJ, Jia XB, Yin W, Shu L. Geniposide promotes beta-cell regeneration and survival through regulating β -catenin/TCF7L2 pathway. Cell Death Dis. 2015;6:e1746.
DOI: 10.1038/cddis.2015.107
134. Dusabimana T, Park EJ, Je J, Jeong K, Yun SP, Kim HJ, Kim H, Park SW. Geniposide improves diabetic nephropathy by enhancing ULK1-Mediated autophagy and reducing oxidative stress through AMPK activation. Int. J. Mol. Sci. 2021;22:1651.
DOI: 10.3390/ijms22041651
135. Niture NT, Ansari AA, Naik SR. Anti-hyperglycemic activity of rutin in streptozotocin-induced diabetic rats: an effect mediated through cytokines, antioxidants and lipid biomarkers. Indian J. Exp. Biol. 2014;52:720-727. [ResearchGate]
136. Cherng YG, Tsai CC, Chung HH, Lai YW, Kuo SC, Cheng JT. Antihyperglycemic action of sinapic acid in diabetic rats. J. Agric. Food Chem. 2013;61:12053-12059.
DOI: 10.1021/jf403092b
137. Pirmoghani A, Salehi I, Moradkhani S, Karimi SA, Salehi S. Effect of *Crataegus* extract supplementation on diabetes induced memory deficits and serum biochemical parameters in male rats. IBRO Rep. 2019;7:90-96.
DOI: 10.1016/j.ibror.2019.10.004
138. Al-Mobideen OK, Alqudah AA, Al-Mustafa A, Alhawarat F, Mizher H. Effect of *Crataegus aronia* on the biochemical parameters in induced diabetic rats. Pharmacog. J. 2022;14:587-595.
DOI: 10.5530/pj.2022.14.140

139. Ramadan BK, Schaalán MF, Tolba AM, Hypoglycemic and pancreatic protective effects of *Portulaca oleracea* extract in alloxan induced diabetic rats. BMC Complement. Altern. Med. 2017;17:37. DOI: 10.1186/s12906-016-1530-1
140. Rakhshandeh H, Rajabi Khasevan H, Saviano A, Mahdinezhad MR, Baradaran Rahimi V, Ehtiati S, Etemad L, Ebrahimzadeh-bideskan A, Maione F, Askari VR. Protective effect of *Portulaca oleracea* on streptozotocin-induced type i diabetes-associated reproductive system dysfunction and inflammation. Molecules. 2022;27:6075. DOI: 10.3390/molecules27186075
141. Huseini HF, Larijani B, Heshmat R, Fakhrzadeh H, Radjabipour B, Toliat T, Raza M. The efficacy of *Silybum marianum* (L.) Gaertn.(silymarin) in the treatment of type II diabetes: a randomized, double-blind, placebo-controlled, clinical trial. Phytother. Res. 2006;20:1036-1039. DOI: 10.1002/ptr.1988
142. Qin NB, Jia CC, Xu J, Li DH, Xu FX, Bai J, Li ZL, Hua HM. New amides from seeds of *Silybum marianum* with potential antioxidant and antidiabetic activities. Fitoterapia. 2017;119:83-89. DOI: 10.1016/j.fitote.2017.04.008
143. Abdel-Zaher AO, Salim SY, Assaf MH, Abdel-Hady RH. Antidiabetic activity and toxicity of *Zizyphus spina-christi* leaves. J. Ethnopharmacol. 2005;10s1: 129-138. DOI: 10.1016/j.jep.2005.04.007
144. Nuru KA, Garkuwa UA, Yusuf H, Tijjani H, Kura AU. Hypoglycaemic potential of *Zizyphus Spina-Christi* fruit on alloxan induced hyperglycaemic rats. J. Bioeq. Stud. 2022;8:102s. [Journal Website]

© 2023 Azab et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/107662>