



Nutritional and Food Security through Kitchen Gardening in Sawai Madhopur District of Rajasthan, India

**Chamandeep Kaur ^{a++*}, B. L. Dhaka ^{a#}, B. L. Meena ^{a†}
and Suresh Bairwa ^{a‡}**

^a KVK, Sawai Madhopur, AU, Kota, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/CJAST/2023/v42i474311

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/110015>

Original Research Article

Received: 25/10/2023

Accepted: 02/12/2023

Published: 13/12/2023

ABSTRACT

Nutritional deficiency like Anaemia, PEM a serious nutritional problem in rural and urban areas in women, pregnant women and adolescent girls; resulting in different types of diseases, negative effects on physical growth and retarding brain development. 77 percent of families in Indian households; vegetables and fruits are often the only source of micronutrients in their diet. Homestead production of fruits and vegetables provides the people the direct access to important nutrients that may not be readily available or within their economic rich. Hence, kitchen gardening is an important strategy to improve household nutritional and food security. Present study was conducted in Sawai Madhopur district of Rajasthan to quantify the impact of 120 kitchen gardens to improve nutritional security of households in rural areas. The result suggests that kitchen gardening

⁺⁺SMS (Home Science);

[#]Senior Scientist & Head;

[†]Associate Professor (Planr Protection);

[‡]SMS (Horticulture);

^{*}Corresponding author;

has proved a Feasibility, Accessibility and Replicability to livelihood strategy for resource Farming Community in terms of nutrient as well as calorie intake and economic performances. In 2019-2020 around 100 Kitchen Garden kits have been distributed among women beneficiaries of Padli villages, in Sawai Madhopur block under NARI programme of KVK, Sawai Madhopur.

Keywords: Nutritional security; kitchen gardening; Rural areas.

1. INTRODUCTION

“Protein Energy Malnutrition (PEM) is a deficiency disease caused in the infants due to 'Food Gap' between the intake and requirement. It affects children under 5 mostly belonging to the poor underprivileged communities. PEM is particularly serious during the post-weaning stage and is often associated with infection. Clinical signs and symptoms of protein-energy malnutrition (PEM) include Poor weight gain, Slowing of linear growth, Behavioural changes - Irritability, apathy, decreased social responsiveness, anxiety, and attention deficits. Protein–energy malnutrition (PEM), sometimes called protein-energy undernutrition (PEU), is a form of malnutrition that is defined as a range of conditions arising from coincident lack of dietary protein and/or energy (calories) in varying proportions. The condition has mild, moderate, and severe degrees. EM is fairly common worldwide in both children and adults and accounts for about 250 000 deaths annually” [1,2]. “In the industrialized world, PEM is predominantly seen in hospitals, is associated with disease, or is often found in the elderly” [3]. Note that PEM may be secondary to other conditions such as chronic renal disease [4] or cancer cachexia [5] in which protein energy wasting (PEW) may occur.

“Protein–energy malnutrition affects children the most because they have less protein intake. The few rare cases found in the developed world are almost entirely found in small children as a result of fad diets, or ignorance of the nutritional needs of children, particularly in cases of milk allergy” [6]. “It retards growth, increases the risk and duration of illness, reduces work output and slows social and mental development” [7]. “For poor households, vegetables and fruits are often the only sources of micronutrients in the family diet. Home gardening is one of the world’s most ancient food production practices and is practiced throughout the world” (Landauer and Brazil, 1985). “Homestead production of fruits

and vegetables provides the households with direct access to important nutrients that may not be readily available or within their economic reach. So, home gardening would be a good means to improve household food security” (Talukder et al., 2002). Therefore, present study was conducted to see the impact of kitchen gardening in improving the nutritional security of households in rural areas. The basic nutritional need of women age group between 18-40 years. In between this age group maximum women’s grow-up, conceives child and do maximum house hold works. This table clearly shows daily nutritional requirement for women likes Energy, protein, fat, iron, B carotene, Vitamin C in form of different cereals, pulses, green leafy vegetables, fruits.

2. METHODOLOGY

The study was conducted in Sawai Madhopur district of Rajasthan. In Sawai Madhopur, there were total 5 blocks namely Sawai Madhopur, Chouth ka Barwara, Khandaar, Bonli, Malarna. For the study sawai madhopur block was selected as NARI programme Village “Padli”. There were all 100 nutritional garden seed kits were distributed to selected families Randomly. “Different capacity building activities including training, exposure visit and farmer’s scientist’s interaction on various aspects including vegetable grown in homestead, homestead vegetable utilization, average vegetable consumption, nutrient contribution from homestead vegetable gardening were planned and undertaken. Pre-survey was conducted to obtain information regarding profile and respondent’s dietary food habits and nutritional deficiency diseases were also pre-surveyed. After one year of establishment of nutritional garden, a post-survey was done to analyses the impact of kitchen gardens on nutritional status of selected families. Data were collected by interview schedule Map of Sawai Madhopur shows production and productivity of major recipes” [8,9].



Fig. 1. Map showing study location

3. RESULTS AND DISCUSSION

Socio-economic characteristics of respondents were analyzed and presented in Table 1. The table indicates that majority (75%) of respondents were belonged to joint family and followed by (25%) nuclear family. "It was found that majority of the families (65%) were from medium sized families followed by small size (20%) and big size (15%). Results on family income

showed that majority (76 %) of respondents belonged to income group of more than Rs. 1.0 lack. While looking at their educational status, results revealed that 70 per cent heads of the family were literate to primary level educated. Results on land holding depicts that majority (60%) had medium scale land followed by small (30%) scale land and only (10%) had large scale land" [8].

Table 1. Socio-economic status of families N=100

S. No	Variables	Category	Number	Percent
1	Type of Family	Joint Family	75	75
		Nuclear family	25	25
2	Size of Family	Small size (1-4)	20	20
		Medium size (5-7)	65	65
		Large size (more than 7)	15	15
3	Annual Income (Rs)	< 50000	10	10
		50000-100000	14	14
		>100000	76	76
4	Literacy	No School	70	70
		(Up to 5h)	8	8
		(Up to 8)	9	9
		(Up to 10)	11	11
		Graduation	2	2
5	Land Holding (ha)	Small	30	30
		Medium	60	60
		Large	10	10

Table 2. Vegetable production and utilization

S. No.	Particular	Production (kg)	Purchase (kg)	Distribution (kg)	Consumption (kg)
1	Before Intervention	205	72	12	265
2	After Intervention	990	0	91	563
3	Change	785	-72	79	298
4	Per cent of Change	193.27	100	450	89.55

Table 3. Change in nutritional status through homestead vegetable production

S. No.	Essential Nutrients/Food Groups	Per Head Per Day Consumption through Vegetables/ Oils/ Fruits/ Nutrients		
		Balanced diet	Before intervention	After intervention
1.	Cereals (g/d)	270	201.01	223.12
2.	Pulses (g/d)	60	32.30	39.00
3.	Green leafy vegetable (g/d)	100	36.2	50.00
4.	Root-tubers (g/d)	200	55	72
5.	Other vegetables (g/d)	250	85.01	120
6.	Fruits	100	25.3	50.3
7.	Milk & milk products (g/d)	300	134	142
8.	Fat & oil (g/d)	20	30	26.8
9.	Sugar (g/d)	20	40	22
10.	Energy (Kcal)	2230	1512	1945
11.	Protein (g)	55	25.35	39.33
12.	Fat(g)	25	22.30	21.00
13.	Calcium (mg)	600	280	382
14.	Iron (mg)	21	10.53	16.00
15.	B Carotene (ug)	4800	1130	3120
16.	Vitamin C (mg)	40	24	35.2

(ICMR 2017) Nutrition to Women between 18-45 years of Age

conceives child and do maximum house hold works. This table clearly shows daily nutritional requirement for women like Energy, protein, fat, iron, B carotene, Vitamin C in form of different cereals, pulses, green leafy vegetables, fruits. Kitchen gardens predict as best source of nutritional and Food security in contexts of Accessibility, Replicability and Utility.

Accessibility as Kitchen gardens are helpful to provide nutritional & economically advantages. Nutritionally & economically beneficial for farm families & Urban Households. Replicability as Kitchen gardening is only given at KVK campus. Utility as by growing kitchen gardens a family can save about 50 thousand rupees yearly.

4. CONCLUSION

It may be concluded that establishment of kitchen gardens had immense role in tackling the problem of malnutrition and micronutrients deficiencies in rural areas. Farm families were highly satisfied by growing kitchen gardens which provide them daily best meal in their thali,

nutrition, variety, economically well saving and easily available. After intervention they had grown 21 items of vegetables like that bottle gourd, bitter gourd, green chilli, brinjal, summer squash, tomato, cucumber, ridge gourd, okra, cluster bean, cow pea, spinach, coriander, cauliflower, onion, cabbage, carrot, pea, fenugreek, radish, papaya, guava, improved variety of millets of bajara and jowar in *Kharif*, Rabi and Zaid seasons. Women of Padli were efficiently doing kitchen gardening which contributes maximum nutritional meals in their daily Nutri -thali with fruits. Production of vegetables at beneficiaries increased 193.27 per cent which resulted in increased consumption (89.55%) and distribution (450%) and money saving. Many Household women started nutria garden and achieve nutritional food security. Above study reveals cereals increased by 223.12, pulses by 39.00, green leafy vegetables 50.00, root tubers by 72, fruits by 50.3. Table further depicts that after intervention farming families receive energy 1945, fat 21.00, calcium 382, Iron 16.00, B carotene 3120, Vitamin C 35. Kitchen gardens predict as best source of

nutritional and Food security in contexts of Accessibility, Replicability and Utility. Accessibility as Kitchen gardens are helpful to provide nutritional & economically advantages. Nutritionally & economically beneficial for farm families & Urban Households. Replicability as Kitchen gardening are only given at KVK campus. Utility as by growing kitchen gardens a family can save about 50 thousand rupees yearly.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Franco V, Hotta JK, Jorge SM, Dos Santos JE. Plasma fatty acids in children with grade III protein-energy malnutrition in its different clinical forms: Marasmus, marasmic kwashiorkor, and kwashiorkor. *Journal of Tropical Pediatrics*. 1999;45(2):71–5.
DOI: 0.1093/tropej/45.2.71
PMID 10341499.
2. Deaths from protein-energy malnutrition, by age - Our World in Data.
3. Jump up to: Dietary reference Intake: The Essential Guide to Nutrient Requirements, published by the Institute of Medicine. Available: <https://www.nap.edu/read/11537/chapter/14?term=protein-energy+malnutrition#151>
4. Muscaritoli Maurizio, Molfino Alessio, Bollea Maria Rosa, Fanelli, Filippo Rossi. Malnutrition and wasting in renal disease. *Current Opinion in Clinical Nutrition and Metabolic Care*. 2009;12(4): 378-83.
DOI:10.1097/MCO.0b013e32832c7ae1
PMID 19474712. S2CID 32472183.
5. Bosaeus, Ingvar. Nutritional support in multimodal therapy for cancer cachexia. *Supportive Care in Cancer*. 2008;16(5): 447–51.
DOI: 10.1007/s00520-007-0388-7
PMID 18196284. S2CID 7078558.
6. Liu T, Howard RM, Mancini AJ, Weston WL, Paller AS, Drolet BA, Esterly NB, Levy ML, et al. Kwashiorkor in the United States: Fad diets, perceived and true milk allergy, and nutritional ignorance. *Archives of Dermatology*. 2001;137(5):630–6.
PMID 11346341.
7. Singh R, Singh VK, Singh S. Organic backyard gardening: a promising approach to enhance household food security and wellbeing. *The Pharma Innovation Journal*. 2018;7(4):169-72.
8. Arya S, Prakash S, Joshi S, Tripathi KM, Singh V. Household food security through kitchen gardening in rural areas of western Uttar Pradesh, India. *International Journal of Current Microbiology and Applied Sciences*. 2018;7(2):468-74.
9. ICMR, Guidelines e-manual. #(ICMR, 2017)**Nutrition to Women Between 18-45 years of Age; 2017.

© 2023 Kaur et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/110015>