



Blood Plasma Biochemical Changes and Conception Rate in Anoestrus Buffaloes Treated with Various Hormone Protocols

Janak Panchal ^{a*}, Ravindra Jadav ^a, Sachin Kalaswa ^a,
Vipul Solanki ^a, Keshav ^b and B.N. Suthar ^a

^a Department of Veterinary Gynaecology and Obstetrics, College of Veterinary Science and Animal Husbandry, Sardarkrushinagar, Dantiwada Kamdhenu University, Gujarat, India.

^b Department of Veterinary Gynaecology and Obstetrics, College of Veterinary Science, Assam Agricultural University, Khanapara, Guwahati, Assam, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JABB/2024/v27i5788

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/115357>

Original Research Article

Received: 01/02/2024

Accepted: 04/04/2024

Published: 06/04/2024

ABSTRACT

To study certain blood biochemical parameters (total cholesterol, total protein, calcium and inorganic phosphorus) at various periods of treatment. Total 40 anoestrus buffalo heifers with history of anoestrus were selected from the semi-arid area of Banaskanth and randomly divided into four equal groups. (1)Group-I:Heatsynch protocol (n=10), (2)Group-II:Ovsynch protocol(n=10), (3)Group-III: Doublesynch protocol (n=10) and (4)Group-IV: Control(n=10) followed by fixed time artificial insemination in treated buffalo heifers and insemination at spontaneous estrus in buffalo heifers of group-IV. The overall conception rate following 3 consecutive services was maximum in Group II (50%) followed by Group III (40%), Group I (30%) and Group IV (30%). The blood plasma

*Corresponding author: E-mail: janakpanchal.1990@gmail.com;

was collected at the time of initiation of treatment (P1), at the time of inj. PGF_{2α} (P2), at the time of 1st FTAI (P3) and day 21st post insemination (P4) for the estimation of biochemical parameters. The blood biochemical profiles include total protein (range: 4.54± 0.19 to 6.11± 0.52 g/dl) which in all group did not show any significant difference during the different period but significant (P<0.05) difference between the groups was noticed. Similarly calcium and phosphorus found ranging between 5.73± 0.20 to 9.53± 0.95 mg/dl and 5.46± 0.22 to 6.95± 0.30 mg/dl, respectively in all four groups. The cholesterol estimated values (range: 34.12 ± 1.75 to 41.97± 2.43 mg/dl) were found not-significantly varying between the groups as well as among the different periods. Heatsynch, Ovsynch and Doublesynch protocol can be successfully used for the induction of estrus in true anoestrus buffalo heifers with Better conception rate would be achieved following Ovsynch protocol in buffalo heifers.

Keywords: Artificial insemination; reproductive cycle; reproductive efficiency; nutrition; buffalo; calcium.

1. INTRODUCTION

The buffalo farmers usually face challenges in detecting estrus as buffaloes have a tendency to show silent estrus and anestrus. Prolonged postpartum acyclicity and anestrus are also responsible for huge economic losses to buffalo breeders. The incidence of anestrus is higher (56.0%) in buffalo heifers than in cow heifers (36.0%) [1]. Anestrus is one of the most commonly occurring reproductive problems in cattle and buffalo of India. It is a functional disorder of the reproductive cycle which is characterized by absence of overt signs of estrus manifested either due to lack of expression of estrus or failure of its detection. In heifers, it poses a herd problem possibly due to low plane of nutrition, stress of seasonal transition or extremes of climatic conditions. Seren *et al.* [2] and Ohashi [3] reported that widespread use of AI in buffalo is still limited due to relatively low expression of estrus behavior in buffaloes. This consideration indicates need for estrus synchronization using fixed time insemination for implementation of breeding programs in buffaloes [4]. To improve reproductive efficiency, several protocols of estrus induction and ovulation synchronization have been developed. These procedures are based on manipulating the Corpus luteum, either to induce premature luteolysis using prostaglandins or to prolong the luteal phase using progestogens. The keys for successful estrus synchronization are closely synchronized rapid decline in circulating progesterin concentrations and synchronous growth and ovulation of a viable follicle. Estrus synchronization has many advantages and is becoming mandatory in modern animal husbandry practices in indigenous cattle and buffaloes, which are known for anoestrus and silent estrus.

Biochemical constituents of blood have great diagnostic value in evaluating the physiological status as well as to improve post-partum fertility in female. Lack of protein or insufficient intake of protein was considered to be a cause of failure or delay in estrus cycle [5]. Cholesterol is synthesized from acetate and is an essential precursor for steroid hormone synthesis. Mineral like calcium and phosphorus have influence on certain enzyme system and affects reproductive efficiency [6] which might be reflected in lower blood level of them. Lack of calcium and phosphorus upsets the proper functioning of genital organs [7]. Looking to the above fact the comparative study of various synchronization protocols and their influence on plasma biochemicals in anestrus buffalo heifers was decided.

2. MATERIALS AND METHODS

2.1 Location and Climatic of the Experimental Area

The study was carried out at Banaskantha district. The district is situated in the north western part of the Gujarat state and lies between north latitudes 23°33' & 24°25' and east longitude 71°07' & 73°02'. The district has semi-arid climate. The recorded maximum and minimum temperature was 32.1 °C and 15.5 °C. Whereas, Relative humidity were 70.5 and 46.15 per cent, respectively during the period of study.

2.2 Experimental Protocol

2.2.1 Selection of the buffalo heifer

A total 40 Anestrus buffalo heifers with the history of anestrus were selected from

villages of Banaskantha milk-shed areas of Gujarat. The buffalo heifers were screened gynaeco-clinically for their reproductive status. Detailed history and rectal palpation findings were recorded. The anestrous buffalo heifers having small, smooth, inactive ovaries with normal genitalia and no palpable CL on either ovary, relaxed uterine horn was selected for study. The reproduction status of their buffalo heifers were re-examined after 10 days for conformation the condition. All these selected buffalo heifers were dewormed.

2.2.2 Grouping of buffalo heifers

All the 40 buffalo heifers were randomly divided in four groups as follow.

2.2.2.1 Group-I (HeatSynch Protocol, n=10)

The Selected buffalo heifers of this group were administered with Intramuscular Inj. of 10 µg of GnRH analogue i.e. Buserelin Acetate 10 µg (Receptal , @ 2.5 ml, Intervet India Pvt Ltd, New delhi) on day 0, followed by Inj. of 500 µg PGF_{2α} analogue i.e. Cloprostenol sodium (Estrumate, @ 2 ml, Intervet India Pvt Ltd) and estradiol benzoate 1 mg I/M (Sigma, USA) on days 7 and 8, respectively, and were fix time artificial inseminated(FTAI) twice at 48 and 60 hrs post-Estradiol injection.

2.2.2.2 Group-II (Ovsynch Protocol, n=10):


The selected buffalo heifers of this group were administered with Intramuscular Inj. of 10 µg of Buserelin acetate on day 0, Inj. of 500 µg Cloprostenol sodium, on day 7 and second Inj. of 10 µg of Buserelin acetate on day 9 followed by fix time artificial insemination twice at 12 and 24 hrs later.

2.2.2.3 Group-III (Double Synch Protocol, n=10)

The buffalo heifers of this group were administered with Inj. of 500 µg Cloprostenol sodium on -3 day (3 days before the first injection of GnRH), I/M followed Inj. of 10 µg Buserelin acetate on day 0, Inj. of 500 µg Cloprostenol sodium on 7th day and second Inj. of 10 µg Buserelin acetate on 9th day, the buffalo heifers were fix time artificial inseminate twice at 12 and 24 hrs later following the second injection of buserelin acetate.

2.2.2.4 Group-IV (Anoestrus control, n=10)

The selected buffalo heifers of this group were injected with normal saline 5 ml, i.m on days 0, 7 and 9 and were observed for onset of natural estrus and artificial insemination was performed on detection of estrus. The group served as untreated control.



2.3 Induction of Estrus and Estrus Intensity

The buffalo heifers were closely observed for the exhibition of estrus symptoms at the time of fix time insemination. The intensity of induce estrus was recorded based on the expressions of estrus symptoms. Estrus symptoms of group IV were recorded at the spontaneous estrus. The estrus intensity of selected buffalo heifers were monitored on the basis of their behavior changes i.e. bellowing, excitement, off feed, estrus mucus discharge as well as edema, erythema and wetness of vulva.

2.4 Conception Rate / Fertility Response

The buffalo heifers which did not show signs of estrus following the treatment were examined per rectally for pregnancy diagnosis on 60th day post insemination to confirm the pregnancy. The buffalo heifers which did not conceive at the fixed time insemination were re-inseminated up to two subsequent estrus cycles. The conception rate up to 3rd insemination was recorded.

2.5 Blood Plasma Collection

2.5.1 Method of blood plasma collection

The 5 ml blood samples from buffalo heifers under treatment were collected in heparinized vials through jugular vein puncture. The samples were centrifuged at 3000 rpm for 15 min and plasma was separated out and was stored deep freeze at -20°C until analyzed.

2.5.2 Schedule of blood plasma collection

Blood plasma collection was as under;

- P1 = before initiation of treatment
- P2 = at the time of injection of PGF_{2α}
- P3= on the day of 1st timed artificial insemination
- P4 = on the 21st day post insemination.

The blood samples from control group were collected on 0, 7, 1st (at spontaneous estrus) insemination and 21st day post insemination.

2.6 Blood Plasma Biochemicals

Total cholesterol, total protein, calcium and phosphorus were quantified in blood plasma by

using diagnostic kits (Coral Clinical System (India) Pvt. Ltd., Goa) and analyzed by using Bio-chemistry Analyzer RX-50V (Micro Lab, India).

2.7 Statistical Analysis of Data

The data collected were suitably tabulated and analyzed following standard statistical method shown by Steel and Torrie [8]. While the test of significance between and within the treatment groups and periods for biochemical profile were made by factorial completely randomized block design and Duncan's new multiple range test. The Duncan's new multiple range test was performed at 5 per cent level of significance.

3 RESULTS AND DISCUSSION

3.1 Intensity and Induction of Estrus

Intensity of estrus was recorded on fix time insemination in different groups of treatment. Different estrus symptoms were bellowing, excitement, off-feed, estrus mucus discharge as well as edema, erythema and wetness of vulva. The finding (bellowing, excitement, mucus discharge and swollen) of present study are similar of Mohan *et al.* [9] in buffaloes following Heatsynch protocol. The recorded estrus intensity at the time of artificial Insemination in Group-I was 90 per cent (9/10), Group-II express 80per cent (8/10), In Group-III estrus induction was 80per cent (8/10), Whereas in group-IV (control anestrous buffalo heifer) it was 50 per cent (5/10). Neglia *et al.* [10] and Atanasov *et al.* [11] observed mucus discharge in buffaloes at the time of AI following Ovsynch protocol. Similar Malik *et al.* [12] and Yotov *et al.* [13] also observed 40 and 66.7 per cent anestrous buffaloes showing estrus mucus discharge following Ovsynch protocol. Karen and Darwish, [14] observed estrus mucus discharge in cyclic heifers and non-cyclic cows following Ovsynch protocol. Hoque *et al.* [15] observed diverse intensity of estrus signs water buffaloes with different protocols like Ovsynch, modified Ovsynch and Double-ovsynch.

When the records of all 40 buffalo heifers were pooled it was observed that 75 per cent (30/40 buffalo heifer) heifers showed induction of estrus. In Heatsynch protocol total 90 per cent (9/10) buffalo heifer showed estrus symptoms at the time of insemination. Similar result was recorded by Buhecha *et al.* [16] in anestrues buffaloes.

However, 100 per cent estrus was recorded following heatsynch protocol in buffalo by Ali et al. [17] and Mohan et al. [18] In Ovsynch protocol total 80 per cent (8/10) buffalo heifers showed estrus induction at the time of insemination. Similar result was recorded by Jabeen et al. [19] Thorat et al. [20] , Savalia et al. [21] and Buhecha et al. [16] in anoestrus buffaloes. In Doublesynch protocol total 80 per cent (8/10) buffalo heifer showed estrus symptoms at the time of insemination. Similar results were recorded by Ozturk et al. [22] in dairy cow. However, 100 per cent estrus was recorded following doublesynch protocol in buffalo by Mirmahmoudi and Prakash, [23] In control group total 50 per cent (5/10) buffalo heifer showed estrus symptoms at different time interval within 90 days. Whereas, Lower estrus response was observed by Naikoo et al. [24] Parmar et al. [25], Nakrani et al. [26] and Buhecha et al. [16] in aneestrus buffaloes. In present study maximum estrus induction was obtained in Heatsynch group followed by Ovsynch and Doublesynch group.

3.2 Conception Rate

The buffalo heifers of treatment group were inseminated at fix time and the control group was inseminated at spontaneous estrus by semen of buffalo bull. The buffalo heifers which failed to conceive at 1st insemination were re-insemination for two subsequent estrus. The recorded overall conception rate of group-I, II, III and IV was 30.00, 50.00, 40.00 and 30.00 per cent, respectively. The 1st insemination conception rate of group I, II, III&IV was 10.00, 20.00, 20.00 and 10.00 per cent, respectively. Whereas the similar fingers for 2nd insemination conception rate was 22.22, 25.00, 12.50 and 22.22 per cent, respectively. Following the third insemination recorded conception rate was 0, 16.66, 14.28 and 0 per cent, respectively in group I, II, III and IV.

In present study 30 per cent conception rate was achieved by using Heatsynch protocol in aneestrus buffalo heifers. Similar conception rate was obtained by Mohan et al. [9] using Heatsynch protocol during summer and winter season in aneestrus buffaloes. Almost similar conception rate was obtained by Buhecha et al. [16] in buffaloes using Heatsynch protocol. However, higher (80%) conception rate was reported by Cevik et al. [27] in cows. Conception rate was 50 per cent following Ovsynch protocol in aneestrus buffalo heifers. Similar results were obtained by Berber et al. [28] Naikoo et al. [24] in buffalo. Derar et al. [29] obtained 62.5 per cent conception rate in buffalo heifers which was higher than the reported in present study. Ramakrishnan et al. [30] obtained 66.66 per cent conception in post-partum aneestrus Gir cow. In Doublesynch protocol obtained conception rate was 40 per cent in aneestrus buffalo heifers of present study. Higher conception rate was obtained by Mirmahmoudi and Prakash, [23] in aneestrus buffaloes. Ozturk et al. [22] reported 72 per cent conception rate in aneestrus dairy cow following Doublesynch protocol. In control group overall conception rate was 30 per cent in aneestrus buffalo heifers. Contrary to the present study higher conception rate [31,26,16] in aneestrus buffaloes and Lower conception rate (10.00%) in buffaloes [13] was observed.

3.3 Biochemical Profile

3.3.1 Total protein (g/dl)

The concentrations of total protein found at various periods in four different groups are presented in Table.1 and Fig. 1.

In the present study the mean plasma total protein level on different period in buffalo heifers under Heatsynch, Ovsynch and Doublesynch revealed that the profile did not vary significantly

Table 1. Total protein concentrations (g/dl) in blood plasma at various periods in groups of anoestrus buffalo heifers (Mean ± SEM).

Period Groups	Total protein (g/dl)			
	P1	P2	P3	P4
Group-I	4.630±0.15 ^d	4.901±0.27 ^d	4.549±0.19 ^d	5.077±0.22 ^c
Group-II	5.243±0.21 ^b	5.489±0.26 ^b	5.123±0.13 ^c	5.594±0.18 ^b
Group-III	4.849±0.25 ^{cd}	4.884±0.22 ^{cd}	5.719±0.38 ^b	4.706±0.31 ^d
Group-IV	5.902±0.20 ^a	6.073±0.24 ^a	6.112±0.52 ^a	5.990±0.49 ^a

Note: - Means bearing different superscripts differ significantly (P<0.05) within Columns

but significant different was observed between groups. It shows close agreement with Savalia et al. [21], Nakrani et al. [26], Buhecha et al. [16] in anestrus buffaloes. Contrary to the present finding Kumari et al. [32] reported significant difference between before and after Ovsynch protocol in responded Murrah buffaloes, however this difference has not been observed in non-responded groups. Higher total protein level was recorded in Anestrus buffaloes [17,26] and anestrus cow [33] was in agreement with the present study.

3.3.2 Total cholesterol (mg/dl)

The concentrations of total cholesterol was found at various periods in four different groups are presented in Table 2 and Fig. 2.

In the present study mean level of plasma total cholesterol level recorded in buffalo heifers under treatment of Heatsynch, Ovsynch and Doublesynch protocol did not showed significant

different in profile between period of blood collection. Similar result was reported by Parmar et al. [25], Savalia et al. [21] and Buhecha et al. [16] in anoestrus buffaloes. Kumari et al. [32] reported the total cholesterol level increase in responded buffaloes after treatment of Ovsynch protocol.

Total cholesterol concentration remained fluctuating 41.97 ± 2.43 mg/dl to 34.12 ± 1.75 mg/dl in all four groups of buffalo heifers during different period. Similar total cholesterol level was recorded by Ghuman et al. [34]. Higher total cholesterol concentration was recorded by Parmar et al. [25], Savalia et al. [21] and Buhecha et al. [16] in anestrus buffalo.

3.3.3 Calcium (mg/dl)

The concentrations of calcium found at various periods in four different groups are presented in Table 3 and Fig. 3.




Fig. 1. Total protein levels (g/dl) at various periods in different treatment groups

Table 2. Total Cholesterol concentration (mg/dl) in blood plasma at various periods in different groups of anoestrus buffalo heifers (Mean + SEM)

Period Group	Total cholesterol (mg/dl)			
	P1	P2	P3	P4
Group-I	37.75±1.46	36.96±1.62	35.20±1.02	39.61±1.54
Group-II	36.07±2.64	35.89±1.29	35.17±2.21	41.97±2.43
Group-III	37.26±2.26	39.77±3.80	39.81±4.34	34.12±1.75
Group-IV	38.98±0.81	40.30±1.66	41.56±3.36	37.88±0.93




Fig. 2. Total cholesterol levels (mg/dl) at various periods in different treatment groups

Table 3. Calcium concentrations (mg/dl) in blood plasma at various periods in different groups of buffalo heifers (Mean± SEM)

Period Group	Calcium (mg/dl)			
	P1	P2	P3	P4
Group-I	5.97±0.27 ^c	5.73±0.20 ^c	5.95±0.22 ^d	6.02±0.22 ^c
Group-II	7.77±0.84 ^a	7.16±0.74 ^{ab}	7.31±0.89 ^c	7.94±0.84 ^b
Group-III	8.33±0.79 ^a	7.85±0.66 ^a	9.53±0.95 ^a	8.80±0.83 ^a
Group-IV	6.81±0.16 ^{bc}	7.01±0.26 ^{ab}	8.24±1.01 ^b	8.48±0.65 ^{ab}

Note: - Means bearing deferent superscripts differ significantly ($P<0.05$) within column




Fig. 3. Calcium levels (mg/dl) at various periods in different treatment groups

Table 4. Inorganic phosphorus concentration in blood plasma at various periods in different groups of buffalo heifers (Mean ± SEM)

Period Group	Phosphorus (mg/dl)			
	P1	P2	P3	P4
Group-I	5.54±0.27 ^c	5.88±0.30 ^{bc}	5.29±0.28 ^c	5.19±0.24 ^c
Group-II	5.76±0.28 ^{bc}	5.81±0.35 ^c	5.66±0.33 ^{bc}	6.11±0.44 ^{ab}
Group-III	6.95±0.30 ^a	6.81±0.41 ^a	6.54±0.24 ^a	6.31±0.45 ^a
Group-IV	5.91±0.26 ^{bc}	6.01±0.39 ^{bc}	5.91±0.35 ^b	5.46±0.22 ^{ab}

Note: - Means bearing different superscripts differ significantly ($P<0.05$) within column




Fig. 4. Phosphorus levels (mg/dl) at various periods in different treatment groups

The plasma calcium level obtained in anestrus buffalo heifer during Heatsynch, Ovsynch and doublesynch protocols did not differ significantly within group. Similar results were obtained by Buhecha et al. [16] in anestrus buffaloes following PRID, Ovsynch and Heatsynch protocol in anestrus buffaloes. Ali et al. (2013) reported significant difference in calcium concentration on day 0 and 10th day in anestrus buffaloes following Ovsynch, Heatsynch and CIDR.

The concentration of plasma calcium in anestrus buffalo heifers during study was obtained in a range of 5.73±0.20 to 9.53±0.95 mg/dl in all groups. Similar results were obtained by Jayachandran et al., [35,36] in anestrus buffaloes. Kumari et al. [32] reported higher calcium concentration during Ovsynch protocol in anestrus buffaloes. Higher calcium concentration was recorded by Paul et al., [37] in anestrus Nili Ravi buffaloes. Ahmed et al. [38]

reported 9.07±0.10 mg/dl in delayed buffalo heifers.

3.3.4 Inorganic Phosphorus (mg/dl)

The concentrations of phosphorus found at various periods in four different groups are presented in Table 4 and Fig. 4.

The plasma inorganic phosphorus level was obtained in anestrus buffalo heifers during Heatsynch, Ovsynch and doublesynch protocols did not reveal significant difference within groups. Similar result obtained by Buhecha et al. [16] in anestrus buffaloes following PRID, Ovsynch and Heatsynch protocol in anestrus buffaloes. Ali et al. [17] reported significant difference in inorganic phosphorus concentration on day 0 and 10th day in anestrus buffaloes following ovsynch, heatsynch and CIDR. Parmar et al. [25] reported the inorganic phosphorus level significantly higher at induce estrus than

pre-treatment in post-partum anoestrus buffaloes. Kumar *et al.* [39] reported not-significant different in inorganic phosphorus level in Murrah buffaloes before and after ovsynch protocol.

The concentration of plasma inorganic phosphorus in anoestrus buffalo heifers during study was obtained in rang (5.73 ± 0.20 to 9.53 ± 0.95 mg/dl) in all groups. Similar result obtained by [40,41,36] in aneustrous buffaloes. Higher phosphorus concentration was recorded by Paul *et al.* [37] in aneustrous Nili Ravi buffaloes. Ahmed *et al.*[38] reported 9.07 ± 0.10 mg/dl in delayed buffalo heifers. Also lower concentration of inorganic phosphorus reported in acyclic Murrah buffaloes [42], aneustrous buffaloes [43,35].

4. SUMMARY

The study included the response of different hormonal protocols like Heatsynch (Group I), Ovsynch (Group II) and Doublesynch (Group-III) for the induction of estrus, intensity of estrus, conception rate, plasma biochemical (total protein, total cholesterol, calcium and phosphorus). The recorded estrus induction was 90, 80 and 80 per cent in Group I, II and III, respectively. The buffalo heifers were inseminated at fixed time with good quality of buffalo bull semen and the overall conception rate following 3 consecution services was maximum in Group II (50%) followed by Group III (40%) and I (30%). The total protein was found significantly ($P<0.05$) different between the groups. The cholesterol estimated values were found not-significantly varying between the groups as well as among the different periods. The levels of calcium in all the four groups were estimated and statistical analysis revealed the significant ($P<0.05$) difference in calcium concentration among the groups. The estimation value of Inorganic phosphorus was differed significantly ($P<0.05$) among the groups but not within the periods.

5. CONCLUSIONS

The present study concluded that the Heatsynch, Ovsynch and Doublesynch protocol can be successfully used for the induction of estrus in true anoestrus buffalo heifers. Comparatively better conception rate could be achieved following Ovsynch protocol followed by Doublesynch and Heatsynch protocol in buffalo heifers. The alteration of blood biochemical like

total protein, total cholesterol, calcium and phosphorus did not show any specific effect due to any of the synchronization protocol.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Ullah N, Anwar M, Rizwan S, Murtaza S. Blood plasma progesterone concentrations in two different veins and comparison of progesterone concentrations and rectal palpation findings to determine ovarian cyclicity in the Nili- Ravi buffalo (*Bubalus bubalis*). Pakistan Veterinary Journal. 2006;26:118–120.
2. Seren E, Parmeggiani A, Zicarelli L, Montemurro N, Pacelli C, Terzano GM. Periestrus endocrine changes in Italian buffaloes. In proceedings of international symposium on prospects of buffalo production in the mediterranean/ middle East. Cairo, Egypt. 1993;393-396.
3. Ohashi OM. (1994). Estrus detection in buffalo-cow. Buffalo Journal. 1994;10:61–64.
4. Ali A, Fahmy S. Ovarian dynamics and milk progesterone concentrations in cycling and non-cycling buffalo-cows (*Bubalus bubalis*) during Ovsynch program. Theriogenology. 2007;68:23-28.
5. Roberts SJ. (1971). Veterinary Obstetrics and genital diseases. 2nd edition; Scientific Book Agency, Calcutta, India; 1971.
6. Dhoble RL, Gupta SK. Serum calcium and inorganic phosphorus during postpartum anoestrus in buffaloes. Indian Journal of Animal Health. 1986;25(2):123-126.
7. Acharya RM. A Note on infertility in cows. Indian Veterinary Journal. 1960;37(8):417-422.
8. Steel RGD, Torrie JH. Principles and procedures of statistics, a biometric approach. 2nd Edn. Mc Graw Hill, Int. Book Agency, Singapore; 1981.
9. Mohan K, Sarkar M, Prakash BS. Efficiency of heatsynch protocol in estrous synchronization, ovulation and conception of dairy buffaloes (*Bubalus bubalis*). Asian-Australian Journal of Animal Science. 2009;22(6):774-780.
10. Neglia G, Gasparrini B, Di Palo R, De Rosa C Zicarelli L, Campanile G. Comparison of pregnancy rates with two

- estrus synchronization protocols in Italian Mediterranean Buffalo cows. *Theriogenology*. 2003;60:125-133.
11. Atanasov A Yotov S, Antonov A, Kolev P, Induction of oestrus and conception rates in bulgarian murrh buffaloes after fixed-time artificial insemination (A Preliminary Study). *Bulgarian Journal of Veterinary Medicine*. 2011;14(3):165–170.
 12. Malik RK, Singh P, Singh LJ, Sharma RK, Phulia SK, Tuli, Chandolia RK. Ovarian response and fertility of ovsynch-treated post-partum anestrous murrh buffaloes. *Buffalo Bulletin*. 2011;30(4):272-276.
 13. Yotov S, Atanasov A, Ilieva Y. Therapy of ovarian inactivity in postpartum Bulgarian Murrh buffaloes by PRID and Ovsynch estrus synchronization protocols. *Asian Pacific Journal of Reproduction*. 2012; (4):293-299.
 14. Karen M, Darwish A. Efficacy of Ovsynch protocol in cyclic and acyclic Egyptian buffaloes in summer. *Animal Reproduction Science*. 2010;119:17–23.
 15. Hoque MN, Talukder AK, Akter M, Shamsuddin M. Evaluation of ovsynch protocols for time artificial insemination in water buffaloes in Bangladesh. *Turkish Journal of Veterinary and Animal Science*. 2014;38:418-424.
 16. Buhecha KV, Dhama AJ, Theodore VK, Thakor R, Parmar SC. Effect of various ovulation synchronization protocols on estrus response, conception rate and blood biochemical profile in anoestrus buffaloes. *International Journal of Advanced Veterinary Science and Technology*. 2016;5 (1):232-238.
 17. Ali R, Shukla SP, Nema SP. Hormonal induction of ovarian cyclicity and conception rate in postpartum anoestrus buffaloes. *Indian Journal of Field veterinarian*. 2012;7(4):44-46.
 18. Mohan K, Nitu, Prakash BS. Evaluation of the heatsynch protocol in murrh buffaloes (*Bubalus bubalis*) during hot summer season. *Journal of Animal Research*. 2014;4(2):241-246.
 19. Jabeen S, Anwar M, Andrabi SMH, Mehmood A, Murtaza S, Shahab M. Determination of ovsynch efficiency for oestrus synchronization by plasma LH and P4 levels in nili ravi buffalo during peak and low breeding seasons. *Pakistan Veterinary Journal*. 2012;33(2):221-224.
 20. Thorat K, Patil AD, Kumbhar UB, Ghoke SS. Improving fertility in postpartum marathwadi anoestrus buffaloes using ovsynch and selectsynch protocols. *Indian Journal of Animal Reproduction*. 2012;33(2):58-60.
 21. Savalia KK, Dhama AJ, Hadiya KK, Patel KR, Sarvaiya NP. Influence of controlled breeding techniques on fertility and plasma progesterone, protein and cholesterol profile in true anoestrus and repeat breeding buffaloes. *Veterinary World*. 2014;7(9):727-732
 22. Ozturk OA, Cirit U, Baran A, Ak K Is doublesynch protocol a new alternative for timed artificial insemination in anestrous dairy cows, *Theriogenology*. 2010;73:568-576.
 23. Mirmahmoudi R, Prakash BS. The endocrine changes, the timing of ovulation and the efficacy of the doublesynch protocol in the murrh buffalo (*Bubalus bubalis*). *General and Comparative Endocrinology*. 2012;177:153–159
 24. Naikoo M, Patel DM, Sarvaiya NP, Killader A. Estrous synchronization in postpartum anestrous mehsana buffaloes using different hormone protocols. *Indian Journal of Field Veterinarian*. 2010;6(2):1-4.
 25. Parmar KH, Shah RG, Tank PH, Dhama AJ. Effect of hormonal and non-hormonal treatment on reproductive efficiency and plasma progesterone, bio-chemical and macro-minerals profile in postpartum anoestrus buffaloes. *Indian Journal of Field Veterinarian*. 2012;8(2):48-54.
 26. Nakarani BB, Panchal MT, Dhama AJ., Hadiya KK, Patel JA, Gosai RK. Effect of control breeding techniques on fertility and plasma profile of biochemical and mineral constituent in anestrous buffaloes. 2014;1 (4B);299-304.
 27. Cevik M, Selcuk M, Dogan S. Comparison of pregnancy rates after timed Artificial insemination in ovsynch , heatsynch and CIDR-based synchronization protocol in dairy cows. *Kafkas Universitesi Veteriner Fakultesi Dergisi*. 2010;16(1):85-89.
 28. Berber RCA, Madureira EH, Baruselli PS. Comparison of two ovsynch protocols (GnRH vs. LH) for fixed timed insemination in buffalo (*Bubalus bubalis*). *Theriogenology*. 2002;57:1421-1430.
 29. Derar R, Hussein HA, Fahmy S, El-Sherry TM, Megahed G. Ovarian response and progesterone profile during the ovsynch protocol in buffalo heifers and post-partum buffalo-cows. *Buffalo Bulletin*. 2012;31(3): 136-148.

30. Ramakrishnan A, Dhama AJ, Naikoo M, Parmar BC, Divekar BS. Estrus induction and fertility response in post-partum anestrus Gir cows. *Indian Journal of Animal reproduction*. 2012;33(1);37-42.
31. Kumar R, Butani MG, Kavani FS, Dhama AJ, Shah RG, Sarvaiya NP, Killedar A. Influence of hormonal and non-hormonal therapies on fertility and serum progesterone, Metabolites and minerals profile in anoestrus crossbred cow. *International Journal of Cow Science*. 2008;11&12(1&2):23-30.
32. Kumari K, Mishra UK, Kumar K, Praveen PK, Dalai N, Shekhar S. Changes in serum biochemical parameters before and after ovsynch treatment in Murrah buffaloes. *Journal of Cell and Tissue Research*. 2015;15(1):4833-4837.
33. Dhama AJ, Nakrani BB, Hadiya KK, Patel JA, Shah RG. Efficacy of different estrus synchronization protocols on estrus induction response, fertility and plasma progesterone and biochemical profile in crossbred anestrus cows. *Veterinary World*. 2015;1310-1316.
34. Ghuman SPS, Singh J, Honparkhe M, Ahuja CS, Dhama DS, Nazir G, Gandotra VK. Differential fertility in dairy buffalo: Role of thyroid and blood plasma biochemical milieu. *Iranian Journal of Applied Animal Science*. 2011;1(2);105-109.
35. Jayachandran S, Nanjappan K, Muralidharan J, Selvaraj P, Manoharan A. Blood biochemical and mineral status in cyclic and postpartum anestrus buffaloes. *International Journal of Food, Agricultural and Veterinary Science*. 2013;3(1):93-97.
36. Kumar R, Ghosh M, Rawat SS, Kumar P, Kumar S. Plasma mineral profile of normal cyclic and post-partum anestrus murrha buffaloes in organized farms. *Animal Science Reported*. 2016;10:43-47.
37. Paul SS, Chawla DS, Lall D. Serum mineral profile and its relationship with reproductive disorders in Nili-Ravi buffaloes. *Indian Journal Animal Nutrition*. 2000;17(4):324-327.
38. Ahmed WM, Bashandy MM, Ibrahim AK, Shalaby SIA, Abd El-Moez SI, El-Moghazy FM, Ibrahim SRE. Investigations on delayed puberty in Egyptian buffalo-heifers with emphasis on clinic-pathological changes and treatment using GnRH (Receptal®). *Global Veterinaria*. 2010; 4(1):78-85.
39. Kumar S, Sexena A, Ramsagar, Kumar S. Study of serum mineral composition in post-partum true Anoestrus buffaloes. *Buffalo Bulletin*. 2015;34(4):397-399.
40. Ahlawat AR, Derashri HJ. Micro mineral profile in cyclin and non-cyclin animals vis a vis conceiving and non- conceiving animals. *The Asian Journal of Animal Science*. 2010;4:175-178.
41. Butani MG, Dhama AJ, Rajesh Kumar. Comparative blood profile of progesterone, metabolites and minerals in anoestrus, suboestrus, repeat breeding and normal cyclic buffaloes. *Indian Journal of Field veterinarian*. 2011;7(2):20-24.
42. Kumar S, Saxena A, Ramsagar. Comparative studies on metabolic profile of anoestrus and normal cyclic Murrah buffaloes. *Buffalo Bulletin*. 2010;29(1):7-11.
43. Chaurasia R, Kushwaha HS, Chaurasia D., Gendley MK, Santra AK. Comparative studies of certain macro minerals during various reproductive states in buffaloes. *Buffalo Bulletin*. 2010;29(4):291-298.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/115357>