



Laparoscopic Cholecystectomy in Acute Cholecystitis: An Updated Review

Kumar Hari Rajah ^{a++*}

^a Taylor's University School of Medicine and Health Science, Selangor-47500, Malaysia.

Author's contribution

The sole author designed, analysed, interpreted and prepared the manuscript.

Article Information

DOI: 10.9734/AJMAH/2024/v22i61033

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/116680>

Review Article

Received: 01/03/2024
Accepted: 02/05/2024
Published: 04/05/2024

ABSTRACT

Laparoscopic cholecystectomy is the gold standard for the management of gallstone disease, but it has recently been incorporated into the treatment of acute cholecystitis. It is divided into early and delayed laparoscopic cholecystectomy. Early laparoscopic cholecystectomy is done within 72 hours of the onset of symptoms, and delayed laparoscopic cholecystectomy is done after eight weeks. As there is no consensus on the timing of early laparoscopic cholecystectomy in adults and elderly patients, we have conducted this review article to examine the timing of early laparoscopic cholecystectomy, the risk factors for conversion to open cholecystectomy, and the role of laparoscopic subtotal cholecystectomy in the management of acute cholecystitis.

Keywords: *acute cholecystitis; early laparoscopic cholecystectomy; timing of laparoscopic cholecystectomy; conversion of laparoscopic cholecystectomy for acute cholecystitis; surgical treatment of acute cholecystitis.*

⁺⁺ Associate Professor of Surgery;

^{*}Corresponding author: E-mail: kharirajah@yahoo.com.my;

1. INTRODUCTION

Acute cholecystitis is characterized by inflammation of the gallbladder secondary to gallstones. It is diagnosed by a combination of clinical presentation and investigations, including blood investigations and ultrasound imaging. The initial treatment consists of intravenous fluids and antibiotics, but definitive treatment is performed surgically via cholecystectomy. Surgical management can be divided into open or laparoscopic cholecystectomy with laparoscopic cholecystectomy being the current most popular mode of treatment [1].

As laparoscopic cholecystectomy was considered the treatment of choice for acute cholecystitis, the timing of cholecystectomy was often debated, with the patient being managed conservatively, and laparoscopic cholecystectomy was performed after six to eight weeks. However, evidence suggests that early cholecystectomy is safe and feasible for the management of acute cholecystitis. Early laparoscopic cholecystectomy can be performed within 72 hours of admission [2].

Surgical treatment of acute cholecystitis was divided into early and delayed laparoscopic cholecystectomy. Early laparoscopic cholecystectomy involves performing cholecystectomy during the initial admission, and delayed cholecystectomy involves performing the surgery six-to eight weeks after surgery. Early laparoscopic cholecystectomy is being encouraged now as it is associated with reduced cost and prevents readmission for recurrent attacks of acute cholecystitis [3].

The optimal timing for performing laparoscopic cholecystectomy in acute cholecystitis is controversial, with more cases managed with early laparoscopic cholecystectomy rather than delayed laparoscopic cholecystectomy. The optimal timing of early laparoscopic cholecystectomy is also controversial, with no clear definition of early cholecystectomy. The most common time is to perform laparoscopic cholecystectomy within 72 hours from the onset of symptoms [4].

The Tokyo guidelines (TG) 13 and 18 have recommended that patients who present with grades 1 and 2 disease should be managed with early laparoscopic cholecystectomy, and patients with grade 3 can be managed with delayed laparoscopic cholecystectomy after initial management of the patient with percutaneous

cholecystostomy and conservative treatment. Early laparoscopic cholecystectomy was defined as performing the procedure in less than 72 hours from the onset of symptoms [5,6].

The World Society of Emergency Surgeons (WSES) has recommended that laparoscopic cholecystectomy is the treatment of choice for acute cholecystitis, and early laparoscopic cholecystectomy should be offered with the definition of early laparoscopic cholecystectomy being surgery performed in patients who present with symptoms within 72 hours or seven days. Laparoscopic cholecystectomy is also safe, feasible and associated with reduced morbidity and hospital stay [7,8].

There is no current consensus on the definition of early laparoscopic cholecystectomy, the optimal time for performing early laparoscopic cholecystectomy, and whether laparoscopic cholecystectomy should be performed as an early or delayed procedure. Conversion from laparoscopic cholecystectomy is also discussed along with its indications. We conducted this review article to answer all these questions regarding the role of laparoscopic cholecystectomy in the management of acute cholecystitis. We conducted a literature review using PUBMED, the Cochrane database of systemic reviews, Google Scholar, and semantic scholars to search for randomized control trials, non-randomized trials, observational and cohort studies, clinical reviews, systemic reviews, and meta-analyses from 1995 to 2023. The following keywords were used: "acute cholecystitis", "early laparoscopic cholecystectomy", "timing of laparoscopic cholecystectomy", "conversion of laparoscopic cholecystectomy for acute cholecystitis," and "surgical treatment for acute cholecystitis". All articles were in English and assessed by manual cross-referencing of the literature. Commentaries, case reports, and editorials were excluded from this review. Only adult patients were included in this study and pregnant patients with acute cholecystitis were excluded. Pediatric patients were excluded from this study.

2. EARLY LAPAROSCOPIC CHOLECYSTECTOMY

Early laparoscopic cholecystectomy is often performed within 72 hours of admission for acute cholecystitis. It was found to be safe and was associated with reduced morbidity and length of hospital stay. The conversion rates are not affected if early laparoscopic cholecystectomy is

performed, and there is also an added economic benefit. The operative time may be longer than that of delayed laparoscopic cholecystectomy [9–19].

The inflammatory changes that occur during acute cholecystitis make dissection of the gallbladder during early laparoscopic cholecystectomy easier as the edema and adhesion are not dense and fibrotic [20].

A retrospective study was conducted by Ambe et al., looking at the critical time for performing early cholecystectomy. A total of 152 patients were included in this study and were divided into those who underwent laparoscopic cholecystectomy within 24 hours of admission and those who underwent laparoscopic cholecystectomy within 72 hours of admission. This study concluded that laparoscopic cholecystectomy can be conducted within 72 hours of admission [21].

A randomized controlled trial comparing early laparoscopic cholecystectomy versus delayed laparoscopic cholecystectomy for acute cholecystitis was conducted by Johansson et al, 145 patients were included of which 74 underwent early laparoscopic cholecystectomy and 71 underwent delayed laparoscopic cholecystectomy after six weeks. There were no differences in conversion rates, bile duct injury, and mortality. The length of hospital stay was shorter in the early laparoscopic cholecystectomy group. This was also confirmed by another prospective randomized controlled study on early versus delayed laparoscopic cholecystectomy for acute cholecystitis by Ozkardes et al. [22,23].

A meta-analysis by Wu et al. compared early and delayed laparoscopic cholecystectomy for acute cholecystitis and included 16 studies and 1625 patients. This study concluded that early laparoscopic cholecystectomy is associated with lower wound infection rates, reduced hospital stays, and reduced costs. There was no

difference in mortality, bile duct injury, and conversion to open cholecystectomy. [24].

Gurusamy et al. conducted a systematic review and meta-analysis of randomized control trials on the safety and effectiveness of early versus delayed laparoscopic cholecystectomy for acute cholecystitis. Five studies with a total of 451 patients were included in this study and this study concluded that there was no difference with regards to bile duct injury and conversion to open cholecystectomy between the groups and the early laparoscopic cholecystectomy was associated with reduced hospital stay. [25].

An up-to-date meta-analysis of randomized controlled trials on early versus delayed laparoscopic cholecystectomy by Lyu et al. also showed no significant difference in wound infection rates, bile duct injury, and total complication rates. The operation time was increased in early laparoscopic cholecystectomy, but the hospital stay was reduced. This was also confirmed by further meta-analyses of randomized control trials on early versus delayed cholecystectomy for acute cholecystitis by Shikata et al, Siddiqui et al, and Borzellino et al. [26–29].

A meta-analysis of case-control studies by Cao et al. compared early and delayed laparoscopic cholecystectomy for acute cholecystitis, and 77 studies were included. The results from the study showed that early laparoscopic cholecystectomy was associated with reduced complication rates, bile duct injury, wound infection rate, and length of hospital stay [30].

Meta-analyses by Zhou et al and Papi et al compared the safety and efficacy of early laparoscopic cholecystectomy versus delayed laparoscopic cholecystectomy and they concluded that early laparoscopic cholecystectomy was safe, and it reduces the stay in the hospital [31,32].

Table 1. Definition of early laparoscopic cholecystectomy

Study	Study type	year	Definition of early laparoscopic cholecystectomy
Johansson et al	Randomized control trial	2004	Within 48 hrs. of admission
Okardes et al	Prospective randomized study	2014	Within 24 hrs. of admission
Agrawal et al	Prospective randomized study	2014	Within 24 hrs. of admission
Cao et al	Meta-analysis	2015	From 72hrs to 7days from admission
Wiggins et al	Prospective randomized study	2019	Within 7 days from admission

3. EARLY LAPAROSCOPIC CHOLECYSTECTOMY IN ELDERLY PATIENTS

A systematic review and meta-analysis were conducted by Loozen et al. on early cholecystectomy for acute cholecystitis in the elderly population. Eight studies with 592 patients were included in this study. The conversion, morbidity, and mortality rates were 23 %, 24%, and 3.5 %, respectively. This meta-analysis concluded that cholecystectomy is feasible and safe for the management of acute cholecystitis in elderly patients, although patient selection is important [33].

The factors affecting early cholecystectomy for acute cholecystitis in older patients include patient and hospital factors. Patient factors included comorbidities such as history of diabetes mellitus, hypertension, heart disease, and personal and social reasons. The hospital factors include the experience of the surgeon, the availability of emergency theatre time, and intensive care services. [34].

Early laparoscopic cholecystectomy in the elderly was associated with better postoperative recovery, reduced morbidity, mortality, and better outcomes than delayed laparoscopic cholecystectomy. The factors that affect the recovery of elderly patients are independent factors that are related to the age and co-morbidities of the patient [35,36].

A systemic review and meta-analysis of early versus delayed laparoscopic cholecystectomy following percutaneous cholecystostomy in elderly patients was conducted by Kourounis et al. 29 studies were included and there was no difference with regards to conversion rates, morbidity, and mortality between the groups. This study concluded that the timing of cholecystectomy after percutaneous

cholecystostomy has no impact on the management of acute cholecystitis [37].

4. CONVERSION TO OPEN CHOLECYSTECTOMY

This is often performed in patients who undergo laparoscopic cholecystectomy where the anatomy of the structures in the Callot's triangle are obscured by acute inflammation and risk of injury to the bile ducts are high. The rate of conversion is anywhere from 2%-23% The risk factors for conversion include male sex, leukocytosis, hyperglycemia, elevated serum amylase, and presence of co-morbidities. Delay in the presentation to the hospital with symptoms of acute cholecystitis also increases the risk of conversion [38–42].

A retrospective study by Sippey et al on the risk factors for conversion to an open procedure in acute cholecystitis include age, body-mass effect, elevated serum alkaline phosphatase, and low serum albumin [43].

A systemic review and meta-analysis on the risk factors for conversion from laparoscopic to open cholecystectomy was conducted by San Lio et al. A total of 35 studies were included and the majority were retrospective studies. The conversion rate was 6% and the following risk factors, age more than 65, male sex, history of acute cholecystitis, presence of diabetes mellitus and previous abdominal surgery increased the risk of conversion [44].

A systemic review on the preoperative and intraoperative risk factors for conversion of laparoscopic cholecystectomy to open cholecystectomy was conducted by Chin et al. A total of 30 studies and 5,477 cases were converted to open cholecystectomy. Male sex, older patients, symptoms of more than 72 hours, obesity, elevated C reactive protein, and diabetes mellitus were risk factors for conversion [45].

Table 2. The conversion rate and complication rates for early laparoscopic cholecystectomy for acute cholecystitis

Study	Year	Study type	N=numbers	Conversion rate-%	Complication rate-%
Papi et al	2004	Meta-analysis	1255	7.99%	3.11%
Lau et al	2005	Meta-analysis	504	16%	3.9%
Gurusamy et al	2009	Meta-analysis	451	20.3%	0.5%

5. LAPAROSCOPIC SUBTOTAL CHOLECYSTECTOMY

This is considered a bail-out technique or a rescue procedure to complete the cholecystectomy laparoscopically, when there is severe inflammation over the calot's triangle and performing a cholecystectomy is difficult. This procedure is an alternative to converting to an open cholecystectomy. A systemic review by Toro et al looked at the various methods of performing this procedure. 19 articles were included, and 678 patients had undergone subtotal cholecystectomy. This study concluded that closure of both the anterior and posterior walls of the gallbladder stump represented the best method to reduce complications [46].

Another systemic review by Azzawi et al on laparoscopic subtotal cholecystectomy for difficult gallbladders was conducted with 45 studies and 2,166 patients who had undergone a subtotal cholecystectomy. The most common procedure performed was the closure of the cystic duct stump with intracorporeal suturing. The most common complication was bile leak and intra-abdominal collection [47].

6. CONCLUSION

Laparoscopic cholecystectomy is considered the gold standard for the management of acute cholecystitis and early laparoscopic cholecystectomy is recommended due to its reduced morbidity, mortality, and reduced cost. The ideal time for performing early laparoscopic cholecystectomy is less than 72 hours from the onset of symptoms.

The problems of implementing early laparoscopic cholecystectomy in most general hospitals are the availability of senior, trained consultant general surgeons, the availability of emergency theatre time, the use of imaging services, and being able to convert and perform an open cholecystectomy. Patients who are undergoing an early laparoscopic cholecystectomy should be counseled about the risks of conversion to an open procedure.

Some general surgical units perform a delayed laparoscopic cholecystectomy after eight weeks for acute cholecystitis due to the problems that come when attempting to perform an early laparoscopic cholecystectomy. As we progress, early laparoscopic cholecystectomy will become

the norm in the management of acute cholecystitis in the future.

CONSENT AND ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Author has declared that no competing interests exist.

REFERENCES

1. Indar AA, Beckingham IJ. Acute cholecystitis. *BMJ*. 2002;325(7365):639-643. DOI: 10.1136/bmj.325.7365.639
2. Koti RS, Davidson CJ, Davidson BR. Surgical management of acute cholecystitis. *Langenbecks Arch Surg*. 2015;400(4):403-419. DOI: 10.1007/s00423-015-1306-y
3. Knab LM, Boller AM, Mahvi DM. Cholecystitis. *Surg Clin North Am*. 2014;94(2):455-470. DOI: 10.1016/j.suc.2014.01.005
4. Thangavelu A, Rosenbaum S, Thangavelu D. Timing of cholecystectomy in acute cholecystitis. *J Emerg Med*. 2018;54(6):892-897. DOI: 10.1016/j.jemermed.2018.02.045
5. Yamashita Y, Takada T, Strasberg SM, et al. TG13 surgical management of acute cholecystitis. *J Hepatobiliary Pancreat Sci*. 2013;20(1):89-96. DOI: 10.1007/s00534-012-0567-x
6. Okamoto K, Suzuki K, Takada T, et al. Tokyo guidelines 2018: flowchart for the management of acute cholecystitis [published correction appears in *J Hepatobiliary Pancreat Sci*. 2019 Nov;26(11):534]. *J Hepatobiliary Pancreat Sci*. 2018;25(1):55-72. DOI: 10.1002/jhbp.516
7. Ansaloni L, Pisano M, Coccolini F, et al. WSES guidelines on acute calculous cholecystitis published correction appears in *World J Emerg Surg*. 2016 Nov 4;11:52. *World J Emerg Surg*. 2016;11:25. Published 2016 Jun 14. DOI: 10.1186/s13017-016-0082-5
8. Pisano M, Allievi N, Gurusamy K, et al. World society of emergency surgery updated guidelines for the diagnosis and treatment of acute calculus cholecystitis.

- World J Emerg Surg. 2020;15(1):61. Published 2020 Nov 5.
DOI: 10.1186/s13017-020-00336-x
9. Al-Mulhim AA. Timing of early laparoscopic cholecystectomy for acute cholecystitis published correction appears in JSLS. 2008 Oct-Dec;12(4):432. JSLS. 2008;12(3):282-287.
 10. Oymaci E, Ucar AD, Yakan S, et al. Determination of optimal operation time for the management of acute cholecystitis: a clinical trial. Prz Gastroenterol. 2014;9(3): 147-152.
DOI: 10.5114/pg.2014.43576
 11. Skouras C, Jarral O, Deshpande R, Zografos G, Habib N, Zacharakis E. Is early laparoscopic cholecystectomy for acute cholecystitis preferable to delayed surgery?: Best evidence topic (BET). Int J Surg. 2012;10(5):250-258.
DOI: 10.1016/j.ijso.2012.04.012
 12. Madan AK, Aliabadi-Wahle S, Tesi D, Flint LM, Steinberg SM. How early is early laparoscopic treatment of acute cholecystitis?. Am J Surg. 2002;183(3): 232-236.
DOI: 10.1016/s0002-9610(02)00789-4
 13. Budişcă OA, Gabor FM, Ahmed S, et al. Early versus delayed laparoscopic cholecystectomy for acute cholecystitis: A single center experience. Chirurgia (Bucur). 2024;119(1):44-55.
DOI: 10.21614/chirurgia.2024.v.119.i.1.p.44
 14. Blohm M, Österberg J, Sandblom G, Lundell L, Hedberg M, Enochsson L. The sooner, the better? The importance of optimal timing of cholecystectomy in acute cholecystitis: Data from the National Swedish Registry for Gallstone Surgery, GallRiks. J Gastrointest Surg. 2017;21(1): 33-40.
DOI: 10.1007/s11605-016-3223-y
 15. Tzovaras G, Zacharoulis D, Liakou P, Theodoropoulos T, Paroutoglou G, Hatzitheofilou C. Timing of laparoscopic cholecystectomy for acute cholecystitis: A prospective non randomized study. World J Gastroenterol. 2006;12(34):5528-5531.
DOI: 10.3748/wjg.v12.i34.5528
 16. Wiggins T, Markar SR, MacKenzie H, et al. Optimum timing of emergency cholecystectomy for acute cholecystitis in England: Population-based cohort study. Surg Endosc. 2019;33(8):2495-2502.
DOI: 10.1007/s00464-018-6537-x
 17. Hartwig W, Büchler MW. Acute cholecystitis: Early versus delayed surgery. Adv Surg. 2014;48:155-164.
DOI: 10.1016/j.yasu.2014.05.008
 18. Agrawal R, Sood KC, Agarwal B. Evaluation of early versus delayed laparoscopic cholecystectomy in acute cholecystitis. Surg Res Pract. 2015;2015: 349801.
DOI: 10.1155/2015/349801
 19. Kao LS, Ball CG, Chaudhury PK; for Members of the evidence based reviews in surgery group. Evidence-based reviews in surgery: Early Cholecystectomy for Cholecystitis. Ann Surg. 2018;268(6):940-942.
DOI: 10.1097/SLA.0000000000002867
 20. Acar T, Kamer E, Acar N, et al. Laparoscopic cholecystectomy in the treatment of acute cholecystitis: Comparison of results between early and late cholecystectomy. Pan Afr Med J. 2017;26:49. Published 2017 Jan 31.
DOI: 10.11604/pamj.2017.26.49.8359
 21. Ambe P, Weber SA, Christ H, Wassenberg D. Cholecystectomy for acute cholecystitis. How time-critical are the so called "golden 72 hours"? Or better "golden 24 hours" and "silver 25-72 hour"? A case control study. World J Emerg Surg. 2014;9(1):60. Published 2014 Dec 16.
DOI: 10.1186/1749-7922-9-60
 22. Johansson M, Thune A, Blomqvist A, Nelvin L, Lundell L. Management of acute cholecystitis in the laparoscopic era: Results of a prospective, randomized clinical trial. J Gastrointest Surg. 2003;7(5): 642-645.
DOI: 10.1016/s1091-255x(03)00065-9
 23. Ozkardeş AB, Tokaç M, Dumlu EG, et al. Early versus delayed laparoscopic cholecystectomy for acute cholecystitis: A prospective, randomized study. Int Surg. 2014;99(1):56-61.
DOI: 10.9738/INTSURG-D-13-00068.1
 24. Wu XD, Tian X, Liu MM, Wu L, Zhao S, Zhao L. Meta-analysis comparing early versus delayed laparoscopic cholecystectomy for acute cholecystitis. Br J Surg. 2015;102(11):1302-1313.
DOI: 10.1002/bjs.9886
 25. Gurusamy K, Samraj K, Gluud C, Wilson E, Davidson BR. Meta-analysis of randomized controlled trials on the safety and effectiveness of early versus delayed laparoscopic cholecystectomy for acute cholecystitis [published correction appears

- in Br J Surg. 2010 Apr;97(4):624]. Br J Surg. 2010;97(2):141-150.
DOI: 10.1002/bjs.6870
26. Lyu Y, Cheng Y, Wang B, Zhao S, Chen L. Early versus delayed laparoscopic cholecystectomy for acute cholecystitis: An up-to-date meta-analysis of randomized controlled trials. *Surg Endosc.* 2018;32(12):4728-4741.
DOI: 10.1007/s00464-018-6400-0
 27. Shikata S, Noguchi Y, Fukui T. Early versus delayed cholecystectomy for acute cholecystitis: A meta-analysis of randomized controlled trials. *Surg Today.* 2005;35(7):553-560.
DOI: 10.1007/s00595-005-2998-3
 28. Siddiqui T, Mac Donald A, Chong PS, Jenkins JT. Early versus delayed laparoscopic cholecystectomy for acute cholecystitis: A meta-analysis of randomized clinical trials. *Am J Surg.* 2008;195(1):40-47.
DOI: 10.1016/j.amjsurg.2007.03.004
 29. Borzellino G, Sauerland S, Minicozzi AM, et al. Laparoscopic cholecystectomy for severe acute cholecystitis. A meta-analysis of results. *Surg Endosc.* 2008;22(1):8-15.
DOI: 10.1007/s00464-007-9511-6
 30. Cao AM, Eslick GD, Cox MR. Early laparoscopic cholecystectomy is superior to delayed acute cholecystitis: A meta-analysis of case-control studies [published correction appears in *Surg Endosc.* 2016 Mar;30(3):1183]. *Surg Endosc.* 2016;30(3):1172-1182.
DOI: 10.1007/s00464-015-4325-4
 31. Zhou MW, Gu XD, Xiang JB, Chen ZY. Comparison of clinical safety and outcomes of early versus delayed laparoscopic cholecystectomy for acute cholecystitis: A meta-analysis. *Scientific World Journal.* 2014;2014:274516.
DOI: 10.1155/2014/274516
 32. Papi C, Catarci M, D'Ambrosio L, et al. Timing of cholecystectomy for acute calculous cholecystitis: A meta-analysis. *Am J Gastroenterol.* 2004;99(1):147-155.
DOI: 10.1046/j.1572-0241.2003.04002.x
 33. Loozen CS, van Ramshorst B, van Santvoort HC, Boerma D. Early cholecystectomy for acute cholecystitis in the elderly population: A systematic review and meta-analysis. *Dig Surg.* 2017;34(5):371-379.
DOI: 10.1159/000455241
 34. Köstenbauer JK, Gandy RC, Close J, Harvey L. Factors affecting early cholecystectomy for acute cholecystitis in older people—a population-based study. *World J Surg.* 2023;47(7):1704-1710.
DOI: 10.1007/s00268-023-06968-9
 35. Xu R, Xu Y, Xu R. Effect of timing of laparoscopic cholecystectomy on postoperative efficacy and rehabilitation of elderly patients with acute cholecystitis. *Am J Transl Res.* 2022;14(2):1107-1113. Published 2022 Feb 15.
 36. Kuwabara K, Matsuda S, Fushimi K, Ishikawa KB, Horiguchi H, Fujimori K. Relationships of age, cholecystectomy approach and timing with the surgical and functional outcomes of elderly patients with cholecystitis. *International Journal of Surgery.* 2011;9(5):392–9. Available: <https://doi.org/10.1016/j.ijisu.2011.03.026>
 37. Kourounis G, Rooke ZC, McGuigan M, Georgiades F. Systematic review and meta-analysis of early vs late interval laparoscopic cholecystectomy following percutaneous cholecystostomy. *HPB (Oxford).* 2022;24(9):1405-1415.
DOI: 10.1016/j.hpb.2022.03.016
 38. Oymaci E, Ucar AD, Aydogan S, Sari E, Erkan N, Yildirim M. Evaluation of affecting factors for conversion to open cholecystectomy in acute cholecystitis. *Prz Gastroenterol.* 2014;9(6):336-341.
DOI: 10.5114/pg.2014.45491
 39. Warchałowski Ł, Łuszczki E, Bartosiewicz A, et al. The Analysis of risk factors in the conversion from laparoscopic to open cholecystectomy. *Int J Environ Res Public Health.* 2020;17(20):7571. Published 2020 Oct 18.
DOI: 10.3390/ijerph17207571
 40. Hadad SM, Vaidya JS, Baker L, et al. Delay from symptom onset increases the conversion rate in laparoscopic cholecystectomy for acute cholecystitis [published correction appears in *World J Surg.* 2008 Dec;32(12):2747. Hussain, Kashif [added]]. *World J Surg.* 2007;31(6):1298-1303.
DOI: 10.1007/s00268-007-9050-2
 41. Spohnholz J, Herzog T, Munding J, et al. Conversion cholecystectomy in patients with acute cholecystitis-it's not as black as it's painted!. *Langenbecks Arch Surg.* 2016;401(4):479-488.
DOI: 10.1007/s00423-016-1394-3
 42. Lipman JM, Claridge JA, Haridas M, et al. Preoperative findings predict conversion

- from laparoscopic to open cholecystectomy. *Surgery*. 2007;142(4):556-565.
DOI: 10.1016/j.surg.2007.07.010
43. Sippey M, Grzybowski M, Manwaring ML, et al. Acute cholecystitis: Risk factors for conversion to an open procedure. *J Surg Res*. 2015;199(2):357-361.
DOI: 10.1016/j.jss.2015.05.040
44. Magnano San Lio R, Barchitta M, Maugeri A, Quartarone S, Basile G, Agodi A. Preoperative risk factors for conversion from laparoscopic to open cholecystectomy: A systematic review and meta-analysis. *Int J Environ Res Public Health*. 2022;20(1):408. Published 2022 Dec 27.
DOI: 10.3390/ijerph20010408
45. Chin X, Mallika Arachchige S, Orbell-Smith J, Wysocki AP. Preoperative and intraoperative risk factors for conversion of laparoscopic cholecystectomy to open cholecystectomy: A systematic review of 30 studies. *Cureus*. 2023;15(10):e47774. Published 2023 Oct 27.
DOI: 10.7759/cureus.47774
46. Toro A, Teodoro M, Khan M. et al. Subtotal cholecystectomy for difficult acute cholecystitis: How to finalize safely by laparoscopy—a systematic review. *World J Emerg Surg*. 2021;16:45.
Available: <https://doi.org/10.1186/s13017-021-00392-x>
47. Al-Azzawi M, Abouelazayem M, Parmar C, et al. A systematic review on laparoscopic subtotal cholecystectomy for difficult gallbladders: A lifesaving bailout or an incomplete operation? *Ann R Coll Surg Engl*. 2024;106(3):205-212.
DOI:10.1308/rcsann.2023.0008

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/116680>