



# **Effect of Chlormequat (CCC) and Gibberellin (GA<sub>3</sub>) on Growth and Flowering of Tuberose (*Polianthes tuberosa* L.) Var. Mexican Single**

**Jujhar Singh<sup>a++\*</sup> and Puneet Kaur<sup>a#</sup>**

<sup>a</sup> Department of Agriculture, Mata Gujri College, Fatehgarh Sahib, Punjab, India.

## **Authors' contributions**

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

## **Article Information**

DOI: <https://doi.org/10.9734/jeai/2024/v46i82690>

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/119193>

**Original Research Article**

**Received: 30/04/2024**

**Accepted: 01/07/2024**

**Published: 17/07/2024**

## **ABSTRACT**

The experiment was conducted at Experimental Farm, Department of Agriculture, Mata Gujri College, Fatehgarh Sahib, Punjab during summer season 2022-2023 to study the effect of growth regulators on growth and flowering of Tuberose (*Polianthes tuberosa*) var. Mexican Single. The experiment was laid out in randomized block design with seven treatments such as no - growth regulators (Control), GA<sub>3</sub>@50ppm, GA<sub>3</sub>@ 100ppm, GA<sub>3</sub>@150ppm, CCC@1250ppm, CCC @ 1500ppm and CCC @1750ppm and the treatments were replicated thrice. Observation were recorded for vegetative and flowering parameters. Maximum plant canopy was recorded with CCC @ 1750ppm. Maximum leaf size was found in GA<sub>3</sub> @ 150ppm. Flowering parameters such as early

<sup>++</sup> Assistant Professor;

<sup>#</sup> MSc. Agriculture Student (Horticulture-Floriculture and Landscape Architecture);

<sup>\*</sup>Corresponding author: E-mail: Jujhar220@yahoo.com;

opening of first florets and Maximum spike length was found with GA<sub>3</sub>@150ppm. Diameter of spike and diameter of first floret recorded maximum with CCC@ 1500ppm. Total water uptake (ml) by flowers during post harvest found to be maximum in CCC@ 1750ppm. Therefore, GA<sub>3</sub> @ 150ppm and CCC@ 1500ppm can be recommended for growth and flowering of Tuberose.

**Keywords:** Growth regulators; flowering; tuberose; GA<sub>3</sub> @ 150ppm and CCC@ 1500ppm.

## 1. INTRODUCTION

Tuberose is an important bulbous plant or (*Polianthes tuberosa* L.), and belong to Amaryllidaceae family and is indigenous to Mexico. The 70–80 cm long, narrow, and densely packed leaves of the tuberose are a light green color. The flowering stalk that also emerges from center of the leaf cluster is between 80 and 120 cm long, bears leaves, and has tubular flower buds. The 5–6 cm long flowers have a slight bend near the base. Normally, 90 to 95 days after planting, tuberose starts to bloom. The tuberose oil is made from the flowers, which also give off a lovely fragrance. Tuberose natural flower oil, which is used in soap making and the cosmetics industry as well as is among the most expensive raw materials for perfumery. In the plains and during the rainy season (April to September), tuberose blooms profusely. On hills, it blooms from May to July. In mild climates, tuberose blooms all year long [1].

Gibberellic acid signaling is important for maintaining source-sink relationships, loading phloem with sucrose, sucrose long-distance transport by phloem cells of plant vascular systems, and metabolic activity in sink organs by sucrose unloading to sink organs or tissues that influence plant performance or growth. Gibberellic acid treatments have been known to play a significant role in promoting various plant developmental processes, including early flowering, increased plant height, and improved leaf yield and quality in various flowering crops [2]. The growth inhibitor Cycocel significantly improves the tuberose flower quality and flower stalk output [3]. Therefore, present research was carried out to study the effect of growth regulators on growth and flowering of tuberose var. Mexican Single.

## 2. MATERIALS AND METHODS

The current study, titled "Effect of growth regulators on growth and flowering of Tuberose var. Mexican single," was conducted from March 2022 to May 2023 at Mata Gujri College's

Research Farm in Fatehgarh Sahib, Punjab. The experimental site's field is located at 30.6435° North latitude and 76.3970° East longitude. The position is 246 meters above mean sea level. The uniformly sized (1-3 cm) bulbs were utilized for planting on the beds (2x2m) with a spacing of 30 cm between rows and 30 cm between bulbs. Foliar application of plant growth regulators on tuberose was done during 2022. In this study, plants of a single Mexican variety were exposed to various concentrations of plant growth regulators, including no - growth regulators (Control), GA<sub>3</sub>@ 50ppm, GA<sub>3</sub>@ 100ppm, GA<sub>3</sub>@ 150ppm, CCC@ 1250ppm, CCC@ 1500ppm, and CCC @ 1750ppm. Three replications of the experiment were used in its Randomized Block Design (RBD) design.

## 3. RESULTS AND DISCUSSION

Findings from the Table 1 shows that maximum plant canopy was found in T<sub>7</sub> (46.76 cm) i.e. CCC @ 1750ppm. It is due to that, CCC helps in reduce the plant growth and reduce apical dominance due to which the plants become bushy. Shah et al. [4] observed more plant canopy with CCC in chrysanthemum. The maximum leaf size was found in T<sub>4</sub> (27.13 cm) i.e. GA<sub>3</sub> @ 150ppm because GA helps in cell division and also helps in cell expansion. Similar results were found by Padaganur et al. [5] in Tuberose.

Flowering parameters such as early opening of first floret was registered with T<sub>4</sub> (170.67 days) i.e. GA<sub>3</sub> @ 150ppm. It is due to that high concentrations of GA<sub>3</sub> shorten the juvenile phase, and near the end of the juvenile phase, the apical meristem begins to produce flowers rather than vegetative development, leading to early flowering in tuberose. Similar result was found by Thakre et al. [6] found that GA<sub>3</sub> helps in early opening of first floret in Tuberose. Maximum length of spike was found in T<sub>4</sub> (81.87 cm) i.e. GA<sub>3</sub> @150ppm. It might be due to acceleration of the blooming spike's cells' elongation and cell division. The above result was in lined with the findings of Manisha et al. [7] in Tuberose. Maximum spike diameter was

**Table 1. Seven treatments and different parameters**

Treatments	Plant canopy (cm)	Leaf size (cm)	Days to opening 1 <sup>st</sup> floret	Spike length (cm)	Spike diameter (mm)	Diameter of 1 <sup>st</sup> floret(cm)	Total water uptake (ml)
T <sub>1</sub> no - growth regulators	31.35	17.83	181.17	63.95	6.79	2.44	71.11
T <sub>2</sub> GA <sub>3</sub> @50ppm	34.22	17.99	228.83	78.81	7.25	3.25	77.77
T <sub>3</sub> GA <sub>3</sub> @ 100ppm	35.64	18.06	184.07	79.84	7.98	3.40	75.55
T <sub>4</sub> GA <sub>3</sub> @150ppm	34.37	27.13	170.67	81.87	7.14	3.42	75.61
T <sub>5</sub> CCC @ 1250ppm	40.11	26.61	268.87	71.91	8.66	4.05	78.33
T <sub>6</sub> CCC @ 1500ppm	39.48	18.35	279.73	73.27	9.26	5.56	78.11
T <sub>7</sub> CCC@ 1750ppm	46.76	25.98	260.87	66.49	7.33	3.91	79.16
Sem±	2.25	0.86	0.28	2.07	0.37	0.09	2.46
CD (0.05)	6.95	2.64	0.86	6.38	1.14	0.29	7.57

recorded in T<sub>6</sub> (9.26 mm) i.e. CCC @ 1500ppm. It is due to that Cycocel (CCC), helps in formation of more food material, specifically carbohydrate, leading to the maximum spike diameter. According to, Baskaran and Misra [8] maximum diameter of spike was observed in CCC @ 1500ppm in Gladiolus. Maximum diameter of first floret was observed in T<sub>6</sub> (5.56 cm) i.e. CCC @1500ppm and. The foliar application of CCC may have increased biomass mobilization to flowers from the source, which in turn increased floret size by suppressing spike length. Similar results were observed by Sagar et al. [9] in Tuberose. Total water uptake (ml) by flowers during post harvest life was found maximum in T<sub>7</sub> (79.16 ml) i.e. CCC@ 1750ppm. It is due to that CCC increases the number of vascular bundles and their diameter and density as compared to control which increase the water uptake in tuberose during vase life. Similar outcomes was observed by Srinivasan et al. [10] in tuberose.

#### 4. CONCLUSION

From the above study it is concluded that GA<sub>3</sub> @ 150ppm and CCC @ 1500ppm is best for growing Tuberose var. Mexican Single in field conditions. GA<sub>3</sub>@ 150ppm and CCC @1500ppm improve the vegetative and flowering parameters of tuberose.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

- Sanap PB, Patil BA, Gondhali BV. Effect of growth regulators on quality and yield of flowers in tuberose (*Polianthes tuberosa* L.) cv. Single. Orissa Journal of Horticulture. 2000;28(1):68-72.
- Kumar P, Raghava SPS, Mishra RL, Singh KP. Effect of GA<sub>3</sub> on growth and yield of China aster. Journal of Ornamental Horticulture. 2003;6(2):110-112.
- Ponnuswami V, Sowmeya S. Flowers and fillers production technology; 2015.
- Shah SC, Gupta LK, Singh F. Effect of growth regulators on growth, yield and pyrethrin content in chrysanthemum. Recent Horticulture. 1994;1(1):84-87.
- Padaganur VG, Mokashi AN, Patil VS. Effect of growth regulators on growth and yield of tuberose Cv. Single. Karnataka Journal of Agricultural Sciences. 2005; 18(2):469-473.
- Thakre S, Panchbhai DM, Mahadik MK, Badge SA. Effect of gibberellic acid on growth and flowering tuberose. International Journal of Chemical Studies. 2019;7(3):153-156.
- Manisha N, Syamal MN, Narayan M, Misra RL, Sanyat M. Effect of gibberellic acid on tuberose. Floriculture research trends in India, Proceedings of the national symposium on Indian floriculture in the new millennium, Lal-Bagh, Bangalore. 2002;350.
- Baskaran V, Misra RL. Effect of plant growth regulators on growth and flowering of gladiolus. Indian Journal of Horticulture. 2007;64(4):479-482.
- Sagar NN, Kawarkhe VJ, Deshmukh M, Lokhande BS. Effect of preplanting growth regulator treatment of bulbs on growth and flowering

- of tuberose (*Polianthes tuberosa* L.)  
Orissa Journal of Horticulture. 2005;33  
(1):39- 42.
10. Srinivasan A, Rajadurai KR,  
Thamarai SSP, Jeyakumar P. Effect of  
pigments, phenol content and post-harvest  
life on application of plant growth  
retardants in *Celosia* spp. Chemical  
Science Review and Letters. 2017;6(23):  
1769-1772.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/119193>