



# Effect of Organic Nutrients and Zinc Biofortification on Growth and Yield of Sponge Gourd (*Luffa cylindrica* L.) CV. Mahy Harita

Agnes Jeba Sheela M <sup>a\*</sup> and Anburani A <sup>a</sup>

<sup>a</sup> Department of Horticulture, Faculty of Agriculture, Annamalai University, Annamalai Nagar, Chidambaram, Tamil Nadu, India.

## Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/ijpss/2024/v36i115140>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/126185>

Original Research Article

Received: 04/09/2024  
Accepted: 08/11/2024  
Published: 16/11/2024

## ABSTRACT

A field experiment was carried out at Orchard, Department of Horticulture, Annamalai University, Faculty of Agriculture, TamilNadu during 2023-2024. The experiment was laid out in RBD with fifteen treatments in three replications. The organic manures used in the experiment were farm yard manure (25 t ha<sup>-1</sup>) and enriched manure (1 t ha<sup>-1</sup>) as soil application along with consortium biofertilizer (2 kg ha<sup>-1</sup>) and foliar application of biostimulants viz., panchagavya (3%), sea weed extract (3%) and effective microbial inoculants (2%). The zinc in the form of *Bacillus subtilis* @ 10 and 20g was applied as soil application. The results of the experiment revealed that the growth parameters viz., vine length, number of leaves, leaf length, leaf breadth and leaf area were recorded the highest in the treatment that received the application of farmyard manure 25 t ha<sup>-1</sup>.

\*Corresponding author: E-mail: [agnesmohan2001@gmail.com](mailto:agnesmohan2001@gmail.com);

Cite as: M, Agnes Jeba Sheela, and Anburani A. 2024. "Effect of Organic Nutrients and Zinc Biofortification on Growth and Yield of Sponge Gourd (*Luffa Cylindrica* L.) CV. Mahy Harita". *International Journal of Plant & Soil Science* 36 (11):254-61. <https://doi.org/10.9734/ijpss/2024/v36i115140>.

<sup>1</sup>combined with panchagavya (3%) as foliar application and *Bacillus subtilis* @ 20g. The yield attributes viz., the highest number of fruits plant<sup>-1</sup>, fruit length, fruit girth, fruit weight, fruit yield plant<sup>-1</sup> and the zinc uptake in fruit were registered in the treatment which received the application of enriched manure @1 t ha<sup>1</sup> combined with panchagavya (3%) as foliar application and *Bacillus subtilis* @ 20g (T11) respectively.

**Keywords:** Organic manures; biostimulants; growth parameters; yield parameters; zinc; biofortification.

## 1. INTRODUCTION

“Sponge gourd (*Luffa cylindrica* L.) is one of the important tropical and subtropical cucurbitaceous crops grown extensively throughout India. It has a smooth surface and is one of the popular vegetables in India. It occupies an area of about 7.21 lakh ha with production of 12.87 lakh tones. The productivity of this crop is 10.52 tonnes per hectare” (Anon., 2022). The tender fruits are used as vegetable which is easily digestible and increase appetite when consumed. It is a highly nutritive vegetable and it contains moisture of 93.2 g, protein 1.2 g, fat 0.20 g, carbohydrate 2.9 g, minerals like calcium 36 mg, iron 1.1 mg and phosphorus 19mg and fibers (0.20 g) per 100 g of edible portion. The sponge of the mature fruit helps the skin in increasing the blood circulation and as a relief for rheumatic and arthritis sufferers. The fruits are also used to cure jaundice and diabetes. A large quantity of inorganic fertilizers are provided to vegetables in order to get higher yield and maximum income in commercial cultivation. But the application of inorganic fertilizer alone may cause human health problems and also pollute the environment. (Shah *et al.*, 2020). Organic fertilizer application may improve the growth by supplying plant nutrients including micro nutrients as well as enhances chemical, physical and biological properties of the soil, thereby providing a better environment for root development by improving the soil structure. Biostimulants are not a fertilizer because they have no direct effect on increase of plant growth and productivity rather, they improve the productivity by enhancing the efficiency of nutrient uptake of already existing nutrient in soil or externally applied nutrient (Manonmani *et al.*, 2022). “The zinc deficiency problem in food crops can be addressed through the zinc biofortification approach to provide adequate zinc content in multiple edible parts of plants. Zn scarcity affects a large portion of arable land, and about one third of the human population suffers from zinc malnutrition due to poor Zn intake” (Upadhyay *et al.*, 2022). Zinc is also critical to tissue growth, wound healing,

taste acuity, connective tissue growth and maintenance, immune system function, bone mineralization, proper thyroid function, blood clotting and cognitive functions. In this regard, the use of zinc-mobilizing bacteria with diverse abilities to promote plant growth is the current need to increase crop productivity, food security and to increase the zinc concentration in the edible parts of crops. The present investigation was undertaken to study the effect of organic nutrients and zinc biofortification on seed germination, seedling vigour, growth, yield and quality of sponge gourd cv. Mahy Harita.

## 2. MATERIALS AND METHODS

The investigation on “Effect of organic nutrients and zinc biofortification on growth and yield of sponge gourd (*Luffa cylindrica* L.) cv. Mahy Harita” was carried out at Orchard, Department of Horticulture, Annamalai University, Faculty of Agriculture, Tamil Nadu during 2022-2023. The experiment was laid out in RBD with 15 treatment combinations in three replications. The treatments are T1: Control, T2: FYM (Farmyard manure), T3: FYM + PG + *Bacillus subtilis* @ 10g, T4: FYM + PG + *Bacillus subtilis* @20g, T5: FYM + SWE + *Bacillus subtilis* @10g, T6: FYM + SWE + *Bacillus subtilis* @ 20g, T7: FYM + EMI + *Bacillus subtilis* @ 10g, T8: FYM + EMI + *Bacillus subtilis* @ 20g, T9: EM (Enriched manure), T10: EM+PG + *Bacillus subtilis* @ 10g, T11: EM+ PG + *Bacillus subtilis* @ 10g, T12: EM + SWE + *Bacillus subtilis* @10g, T13: EM + SWE + *Bacillus subtilis* @ 20g, T14: EM + EMI + *Bacillus subtilis* @10g, T15: EM + EMI + *Bacillus subtilis* @20g. The sponge gourd variety Mahy Harita (MSGH 6) produced by Mahyco private Limited, Mumbai was used for the experiment. The fruits are dark green with slender in shape and matures in 40-45 days after sowing. The zinc uptake of fruit was estimated by using triple acid digestion method described by Lindsay and Norwell (1958) with a atomic absorption spectrophotometer. The field was thoroughly ploughed and divided into plots of 3m x 3m. Six

pits per plot were formed and the seeds were sown. The organic manures viz., FYM 25 t ha<sup>-1</sup>, EM 1 t ha<sup>-1</sup> along with consortium of biofertilizers @ 2 kg ha<sup>-1</sup> were incorporated at the time of last ploughing as per the treatment schedule. The zinc was applied in the form of *Bacillus subtilis* through soil application at 10g and 20g at different levels. The required quantity of biostimulants, namely Panchagavya (3%), seaweed extract (3%), and effective microbial inoculants (2%), were prepared and sprayed according to the treatment schedule in three split doses, viz., 20, 35, and 50 days after sowing.

The crop was irrigated every fifth day, and proper drainage facilities were provided, as the crop cannot withstand waterlogging. Weeding was done 15 days after sowing. Necessary plant protection measures were carried out as per recommendations. The plants were harvested separately for each treatment at 45 DAS. The fruits of sponge gourd take 6 to 7 days from setting to reach marketable size.

### 3. RESULTS AND DISCUSSION

#### 3.1 Growth Parameters

The results revealed that the growth parameters (Table 1) viz., vine length, number of leaves, leaf length and leaf breadth were significantly influenced by the supplementation of organic manures alongwith consortium of biofertilizers and biofortification of zinc at varying levels. The highest vine length(136.40 cm, 267.37 cm, 368.46 cm), number of leaves (47.88, 85.29, 187.19), leaf length (14.16 cm) and leaf breadth

(15.42 cm) were recorded with 25 t FYM ha<sup>-1</sup> and foliar application of panchagavya (3%)along with soil application of *Bacillus subtilis* @ 20g. The least vine length (84.24 cm, 180.07 cm, 286.44cm), number of leaves (15.24, 41.51, 136.84), leaf length (9.53 cm) and leaf breadth (9.84 cm) wererecorded in control (T1).

The increase in vine length could be due to the organic manure applied in the form of FYM, which might have improved the physical and chemical properties of the soil, leading to an adequate supply of nutrients to the plants, sufficient water-holding capacity, and accelerated vine length. Alternatively, the reason for increased vine length may also be due to improved nutrient uptake by plants in this treatment, resulting in improved vegetative growth. The present findings are in close agreement with (Singh *et al.*, 2020) who reported that the application of organic manures improved vine length in cucumber.

The increase in plant growth might be attributed to the application of Panchagavya spray. (Sam Ruban *et al.*, 2019) reported similar findings in brinjal, where plant height significantly increased with the application of panchagavya that possess almost all macro, micronutrients and growth promoting hormones (IAA, GA) required for plant growth. Further, in the present study application of *Bacillus* also enhanced the vinelength. (Sreekumar and Singh 2020) reported that some of the strains of *Bacillus* were found to producemixtures of lactic acid, isovalericacid, isobutanicacid and acetic acid which might have directly or indirectly promoted the growth attributesin sponge gourd.



Fig. 1. Experimental field view of sponge gourd (*Luffa cylindrica* L.)

**Table 1. Effect of organic manures and zinc on growth parameters in sponge gourdc v. Mahy Harita**

Tr.No	Vine length (cm)			No.of leaves (cm)			Leaf length(cm)	Leaf breadth(cm)	Leaf area(cm <sup>2</sup> )
	30DAS	60DAS	90DAS	30DAS	60DAS	90DAS			
T1	84.24	180.07	286.44	15.24	41.51	136.84	9.53	9.84	47.04
T2	90.81	223.11	315.41	16.41	49.20	146.36	10.19	10.58	53.36
T3	129.07	260.22	356.34	37.16	75.62	178.06	13.19	14.55	96.27
T4	136.40	267.37	368.46	47.88	85.29	187.19	14.16	15.42	109.17
T5	107.98	241.58	336.22	21.40	59.66	160.54	11.46	12.01	68.81
T6	117.00	247.57	345.12	26.63	64.34	168.30	12.14	12.77	77.51
T7	99.48	235.48	327.59	17.91	54.57	153.59	10.72	11.39	61.05
T8	115.00	247.42	345.05	26.27	63.13	167.73	12.15	12.80	77.76
T9	87.84	209.37	302.79	15.81	45.93	142.57	9.85	10.20	50.23
T10	120.88	252.35	349.61	30.13	68.36	171.41	12.47	13.54	84.42
T11	132.52	263.28	361.78	41.62	79.15	182.59	13.84	14.94	103.38
T12	96.89	231.48	321.37	17.04	51.97	149.97	10.44	11.01	57.47
T13	103.23	237.45	332.62	19.20	57.72	156.71	11.08	11.68	64.70
T14	112.31	244.52	340.34	23.44	61.75	164.05	11.82	12.43	73.46
T15	124.76	256.38	354.62	33.50	72.10	174.68	12.79	13.89	88.82
S.ED	1.01	2.17	3.34	0.70	0.86	1.38	0.12	0.15	0.67
CD(p=0.05)	2.04	4.38	6.73	1.40	1.73	2.76	0.24	0.30	1.35

Similar findings on the increase in the number of leaves due to the application of organic nutrients have been reported by (Bhattarai and Maharajan 2013) in carrot. The increase in the number of leaves could also be due to the sudden release of a higher level of nutrients and minerals from FYM, which is readily available to plants. The nitrogen released from FYM is synthesized into amino acids which are built into complex protein and helped in better growth.

Application of consortium biofertilizer also increased the number of leaves in the present study. The increase in number of leaves due to the application of consortium biofertilizer improves better plant growth promotion ability to PGPR as the consortium, apart from the nutrient supplying potential are able to synthesise phytohormones, decompose organic matter, enlarge the soil flora and improve the soil structure for root development and better absorption of water and nutrients. Foliar spray of panchagavya increase the number of leaves in the present study. It might be due to the presence of various growth enzymes which favours rapid cell division and cell- multiplication contributing to the overall growth and development of plants resulting in better yields (Kumar and Singh 2020). According to the findings reported by (Ali *et al.*, 2024) in okra, in the present study the soil application of *Bacillus* increased the number of leaves. The reason could be due to auxins, gibberellins, and cytokinins are the growth-promoting compounds produced by *B. subtilis*. These chemicals may helpful to increase the plant growth and output by enhancing nutrient uptake and water use efficiency.

Similar findings of enhanced leaf parameters were reported by (Pathak *et al.*, 2017) in radish. The application of organic manure likely facilitated rapid cell elongation and multiplication, thanks to the adequate nitrogen supply from farmyard manure during the crop's early stages. This led to increased leaf parameters and leaf area.

The role of biofertilizers played a crucial part in crop growth by maximizing nutrient solubilization potential, efficiently transforming nutrients from unavailable to available forms. These results align with (Bhuvaneshwari and Anburani 2023) who observed similar effects in bottle gourd.

The spray of panchagavya enhanced the leaf area, could be due to the presence of growth-

regulatory substances, such as IAA, GA, cytokinin, essential plant nutrients, effective microorganisms, and biofertilizers present in panchagavya. It also been reported to contain bacteria producing plant growth-promoting substances and exhibiting biological activities. The present findings are consistent with (Esakkiammal *et al.*, 2015) who reported similar results in Dolichos lablab. Application of *Bacillus subtilis* also increased leaf area, as reported by (Singh, 2020) in cucumber.

### 3.2 Yield Parameters

The results revealed (Table 2) that the higher number of fruits plant<sup>-1</sup> (23.76), fruit length (37.30cm), fruit girth (14.64 cm), single fruit weight (234.18 g) and highest fruit yield plant<sup>-1</sup> (5.24 kg plant<sup>-1</sup>). The least number of fruits plant<sup>-1</sup> (9.22), fruit length (16.69 cm), fruit girth (11.59 cm), single fruit weight (90.40g), and fruit yield plant<sup>-1</sup>(1.02 kgplant<sup>-1</sup>) were recorded in the treatment T1(control).

The application of various organic manures, combined with foliar application of panchagavya and soil application of *Bacillus subtilis*, enhanced the yield parameters. The increase in high yield and yield parameters, *viz.*, number of fruits, fruit length, fruit girth, single fruit weight, and fruit yield per plant found in organic treatments, might be due to the synergistic interaction between enriched compost and biofertilizers consortium and the mineralization of macro and micronutrients from organic manures, which might have helped achieve higher yields. Such findings were in accordance with (Barik *et al.*, 2018) in ridge gourd.

Further, in the present study, combined application of panchagavya improved maximum fruit yield and could be due to microbes present in panchagavya that produced growth hormones, which helped increase the weight of fruit and number of fruits per plant through cell division and cell elongation by translocating more carbohydrates to developing fruits. The effect of panchagavya on yield parameters was already reported by (Sanjiv *et al.*, 2019) in tomato. Furthermore, in the present study, *Bacillus subtilis* isolates increased plant yield and induced resistance to biotrophic fungal plant pathogens in tomato. These are presumably transported into the shoot via the xylem. Intensified and prolonged synthesis of these phytohormones may be regarded as a cause of delayed senescence and improved yields.

**Table 2. Effect of organic manures and zinc on yield parameters in sponge gourd cv. Mahy Harita**

Tr.No.	No. of fruits per plant <sup>-1</sup>	Fruit length(cm)	Fruit girth(cm)	Mean single fruit weight(g)	Fruit yieldplant <sup>-1</sup> (kg)	Zinc up take in fruit (mg100g <sup>-1</sup> )
T1	9.22	16.69	11.59	90.40	1.02	0.10
T2	10.01	21.56	12.05	121.15	1.36	0.60
T3	17.61	28.77	13.92	204.21	3.41	2.03
T4	19.12	34.23	14.48	228.41	4.23	2.20
T5	11.81	26.05	12.58	161.64	1.85	1.61
T6	13.28	26.57	12.97	171.34	2.20	1.73
T7	14.01	27.18	13.43	183.91	2.53	1.82
T8	16.46	28.00	13.76	194.15	3.10	1.95
T9	10.65	22.40	12.36	138.32	1.42	0.90
T10	18.76	32.20	14.29	223.81	4.13	2.12
T11	23.76	37.30	14.64	234.18	5.24	2.31
T12	11.87	26.22	13.18	163.18	1.92	1.62
T13	15.21	27.01	13.53	188.73	2.83	1.89
T14	13.96	26.01	13.07	182.36	2.50	1.80
T15	18.22	31.20	14.12	217.34	3.90	2.08
S.ED	0.33	0.35	0.18	3.71	0.07	1.70
CD(p=0.05)	0.66	0.70	0.36	7.48	0.15	3.41

The results of this experiment were supported by the findings of (Sreekumar and Singh 2020) in sponge gourd, who recorded significantly higher fruit yield by applying *Bacillus subtilis* and increased the number and diameter of fruits.

### 3.3 Zinc Content

The result revealed (Table 2) that the zinc content in fruit were recorded highest in treatment T11 (2.31 mg100 g<sup>-1</sup>) which received the application of enriched manure 1 t ha<sup>-1</sup> + panchagavya @ 3% + *Bacillus subtilis* @ 20g. The least zinc content in fruit (0.10 mg 100 g<sup>-1</sup>) were recorded in T1 (control). The soil application of zinc in the form of *Bacillus subtilis* increased in the zinc uptake in fruits. Similar findings were reported by (Anwar *et al.*, 2021) in okra plants.

## 4. CONCLUSION

Based on the field experiments conducted, it can be concluded that the combined application of:

- Enriched manure at 1 t/ha
- Panchagavya (3% concentration)
- *Bacillus subtilis* at 20g

was the most effective treatment in enhancing growth and yield, of sponge gourd cv. Mahy Harita.

## DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

## ACKNOWLEDGEMENT

The author is thankful to Advisor and Department of Horticulture, Faculty of Agriculture, Annamalai University, Annamalai Nagar, Chidambaram, Tamil Nadu for providing necessary facilities to undertaken this research. This manuscript has not received any grant from any sources.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

Ali, H. H., S. Bibi, M.S. Zaheer, R. Iqbal, A. E. Z. M. Mustafa, and M.S. Elshikh. 2024.

Control of copper-induced physiological damage in okra (*Abelmoschus esculentus* L.) via *Bacillus subtilis* and farm yardmanure: A step towards sustainable agriculture. *Plant Stress.*,11:100309.

Anonymous. Area and production of horticulture crops. Ministry of Agriculture and Farmers Welfare, Government of India, Gurgaon, India; c2022. <https://agricoop.nic.in>

Anwar, H., X. Wang, A. Hussain, M. Rafay, M. Ahmad, M. Latif and A. Mustafa. 2021. Comparative effects of bio-wastes in combination with plant growth-promoting bacteria on growth and productivity of okra. *Agronomy.*,11(10):2065.

Barik, N., D. Phookan, V. Kumar, T. Millik and D. Nath. 2018. Organic cultivation of ridge gourd (*Luffa acutangula* Roxb.). *Current Journal of Applied Science and Technology.*,26(4):1-6.

Bhattarai, B.P., and A. Maharjan. 2013. Effect of organic nutrient management on the growth and yield of carrot (*Daucus carota* L.) and soil fertility status. *Nepalese Journal of Agricultural Sciences.*,11:16-25.

Bhuvaneshwari, S., and A. Anburani. 2023. Effect of soil and foliar application of organic nutrients on growth and yield of bottle gourd (*Lagenaria siceraria* (Molina) Standl.). *Annals of Plant and Soil Research.*,25(1):127-132.

Esakkiammal, B., C. Esaivani, K. Vasanthi, L. L. Bai and N. S. Preya. 2015. Microbial diversity of vermicompost and vermish prepared from *Eudriluseuginae*. *International Journal of Current Microbiology and Applied Sciences.*,4(9):873-883.

Kumar, C. S., G. Singh. 2020. Effect of panchagavya on growth and yield: a review. *International Journal of Current Microbiology and Applied Sciences.*,9(12):617-624.

Lindsay, W. L. and W. A. Norwell. 1958. Development of DTPA soil test for Fe, Mn, Zn and Cu. *Soil Science Society of America Journal.*,42 pp:421-428.

Manonmani, S., S. Senthikumar and S. Manivannan. 2022. Multilevel functionality of biostimulants in sustainable horticulture reform modern era. *The Pharma Innovation Journal.*, 11(10):1283-1288.

Pathak, M., P. Tripathy, S. K. Dash, G. S. Sahu and S. K. Pattanayak. 2017. Effect of source of nutrient on growth, yield and

- quality of Radish (*Raphanus sativus* L.) in radish-coriander cropping sequence. The Pharma Innovation Journal., 6(12): 496-499.
- Sam Ruban, M. R Priya, G. Barathan, S. M Suresh Kumar. 2019. Effect of foliar application of biostimulants on growth and yield of brinjal (*Solanum melongena* L.) Plant Archives., 19(2): 2126-2128.
- Sanjiv Yadav, Amit Kanawjia and Rajkumar Chaurasiya. 2019. Response of bio-enhancer on growth and yield of tomato (*Solanum lycopersicum* (L.) Mill). International Journal of Chemical Studies., 7(3):180-184.
- Shah, P. S., G. Nabi, M. Khan, S. Hussain and J. A. Jan. 2020. Performance of sponge gourd cultivars under organic and inorganic fertilizer regimes. Sarhad Journal of Agriculture., 36(2): 478-488.
- Singh, J., M. K. Singh, M. Kumar, A. Gupta and K. P. Singh. 2020. Growth, yield and quality parameters of cucumber (*Cucumis sativus* L.) as influenced by integrated nutrient management application. International Journal of Current Microbiology and Applied Sciences., 9(10):1455-1462.
- Singh, N. 2020. Effect of plant growth promoting *Rhizobacteria* on seed germination seedling vigour growth and yield of cucumber *Cucumis sativus* L and bottle gourd (*Lagenaria ciceraria*) Molina Standl. Journal of Pharmacognosy and Phytochemistry., 9(4):15559-1562.
- Sreekumar, G. and D. Singh. 2020. Study on growth and yield of sponge gourd by using plant growth promoting *Rhizobacteria* in prayagraj agro-climatic condition. International Journal of Current Microbiology and Applied Sciences., 9(08):3585-3591.
- Upadhayay, V. K., A. V. Singh, A. Khan and A. Sharma. 2022. Contemplating the role of zinc-solubilizing bacteria in crop biofortification: An approach for sustainable bioeconomy. Frontiers in Agronomy., 4903321.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/126185>