



Evaluating the Impact of Black Soldier Fly (*Hermetia illucens*) Larvae Meal on Growth Efficiency and Feed Utilization in Rainbow Trout (*Oncorhynchus mykiss*) Juvenile

Ishori Singh Mahato ^a, Krishna Paudel ^{a*}, Prem Timalsina ^b
and Sunita Chand ^a

^a Fishery Research Station, Trishuli, Nuwakot, Nepal.

^b National Fishery Research Centre, Godawari, Lalitpur, Nepal.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ajfar/2024/v26i12850>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/127454>

Original Research Article

Received: 25/09/2024

Accepted: 27/11/2024

Published: 03/12/2024

ABSTRACT

Aims: Black Soldier Fly (BSF) larvae meal is a promising alternative protein source in aquafeed, offering high nutritional value and essential amino acid profiles like fishmeal. Nepalese Rainbow trout (*Oncorhynchus mykiss*) farming currently relies on imported shrimp meal, presenting economic and ecological challenges. The focus of this research was to examine the potential of BSFL meal to replace the shrimp meal in trout feed.

*Corresponding author: Email: aceskrishna@gmail.com;

Cite as: Mahato, Ishori Singh, Krishna Paudel, Prem Timalsina, and Sunita Chand. 2024. "Evaluating the Impact of Black Soldier Fly (*Hermetia Illucens*) Larvae Meal on Growth Efficiency and Feed Utilization in Rainbow Trout (*Oncorhynchus Mykiss*) Juvenile". *Asian Journal of Fisheries and Aquatic Research* 26 (12):107-117. <https://doi.org/10.9734/ajfar/2024/v26i12850>.

Place and Duration of Study: Fisheries Research Station (FRS), Trishuli Nepal, between May 1 to August 7, 2023.

Methodology: A feeding trial was conducted using 750 healthy Rainbow trout (mean individual body weight 5.39 ± 0.55 g) randomly divided into five experimental groups, each having three replicates. The control diet contained no BSFL meal (CON), while experimental diets replaced 25% (BSF25), 50% (BSF50), 75% (BSF75), and 100% (BSF100) of shrimp meal with BSFL meal. Fish were raised in flow-through raceway tanks (300L) and fed twice daily until satiation. The growth efficiency parameters were measured.

Results: The results demonstrated that diets with 25-50% BSFL meal substitution (B25, B50) significantly enhanced growth efficiency compared to the control, reflected in higher final body weight and weight gain, and improved feed conversion ratios ($P < 0.05$). Conversely, higher replacement levels (75% and 100%) negatively impacted growth, possibly due to increased chitin content affecting digestibility. Statistical analysis confirmed the optimal BSFL inclusion range for growth efficiency, with specific growth rates peaking in the B50 group.

Conclusion: In conclusion, BSFL meal can effectively replace 25-50% of shrimp meal in rainbow trout feed, offering a viable and sustainable protein source for trout farming in Nepal. This substitution could reduce dependency on imported shrimp, promoting more sustainable practices.

Keywords: BSFL; replacement; insect meal; efficiency; bioconversion.

1. INTRODUCTION

Currently, cold-water aquaculture in Nepal is primarily focused on a single fish species, the Rainbow trout (*Oncorhynchus mykiss*). Approximately 120 farmers across 38 mid-high hill districts of Nepal produce around 420 metric tons of trout annually (Mahato et al., 2024). However, this production level is expected to rise as 56 districts have been identified as suitable for rainbow trout cultivation (FRS, 2023). With the growing popularity of rainbow trout among the populace, the demand for this species is also increasing. Nearly 100% of the trout produced in Nepal is consumed domestically. As rainbow trout farming expands, the demand for complete feed will also escalate. Presently, shrimp meal, known for its high protein content (approximately 46-60%) and excellent protein composition, as well as its natural source of astaxanthin pigment for imparting red coloration in salmonids, is a primary protein source in rainbow trout feed in Nepal (Mahato et al., 2023). This shrimp meal is predominantly imported from India. Research indicates a significant decline in marine and ocean life in recent decades, partly due to overfishing (Pauly and Christensen, 1995), resulting in shrimp becoming scarcer and more expensive as fishery products worldwide decline (Mohsen and Lovell, 1990). Furthermore, importing shrimp from India costs several hundred thousand annually, with no guaranteed consistency in quality. Consequently, the identification and production of more economic and sustainable sources of protein for trout feed is of vital importance.

In recent years, a variety of novel alternative protein sources for aquafeed have been explored. Insect protein is one of the emerging and excellent alternative protein sources that could potentially replace fishmeal or shrimp meal in aquafeed in the future. Although insect protein has been experimented with in poultry feed since 1969 (Calvert et al., 1969), it has only gained significant attention in aquaculture feed over the past two decades. Some insects like the silkworm (*Bombyx mori* L.), the black soldier fly (*Hermetia illucens* L.), the house fly (*Musca domestica* L.), the mealworm (*Tenebrio molitor* L.), the lesser mealworm (*Alphitobius diaperinus*), the banded cricket (*Grylloides sigillatus*), the house cricket (*Acheta domesticus* L.) and the Jamaican field cricket (*Gryllus assimili*) have already achieved researcher's attention (Bartucz et al., 2023). It is because, insects are deemed an economically and ecologically advantageous ingredient because they convert food into protein efficiently, require minimal arable land and water, have a low environmental impact, necessitate minimal investment in machinery, and possess rapid reproduction cycles (Nugroho and Nur, 2018).

In recent years, the insect species *Hermetia illucens*, commonly known as the black soldier fly, has garnered significant attention as an alternative protein source to replace fish meal in aquaculture feeds as it has been shown to have essential amino acid profile like that of fishmeal (Henry et al., 2015). In addition, *black soldier fly* larvae (BSFL) is packed with proteins (60–80%), essential amino acids, vitamins, and minerals,

plus significant lipids (31–43%), which can vary based on how they're raised (Meneguz et al., 2018). These larvae are great at converting organic material, eating twice their weight daily, and have high protein (36–48% DM) and fat (31–33% DM) levels before pupating (Henry et al., 2015). Recent studies have examined its effects on various fish species, including Mirror carp (*Cyprinus carpio*) (Xu et al., 2020), Atlantic salmon (*Salmo salar*) (Fisher et al., 2020), African catfish (*Clarias gariepinus*) (Fawole et al., 2020), European seabass (*Dicentrarchus labrax*) (Moutinho et al., 2021), Rainbow trout (*Oncorhynchus mykiss*) (Bruni et al., 2020), and others, focusing on growth efficiency, feed utilization, gut health, and microbiome modulation. In salmon and sea bass the *Hermetia illucens* replaced 40% of fishmeal (Lock et al., 2016; Magalhães et al., 2017), and up to 50% in trout (Magalhães et al., 2017) without negatively affecting the growth efficiency. Research also suggests *Hermetia illucens* larvae meal may positively influence the gut microbiota, promoting bacteria that produce beneficial short-chain fatty acids like butyrate through chitin fermentation critical for intestinal cells supporting intestinal development, immunity, and disease resistance in rainbow trout (Biasato et al., 2022; Terova et al., 2019). Therefore, given the evidence, it is plausible to anticipate that replacing part of the fishmeal or shrimp meal with insect meal could have a beneficial impact on the growth and intestinal microbiota of fish.

While in overseas extensive nutritional research has examined how incorporating *Hermetia illucens* meal into diets affects fish growth (Henry et al., 2015), and intestinal microbiota composition (Bruni et al., 2020), less is known about its specific influence on growth efficiency of Rainbow trout juveniles in the context of Nepal. For this reason, the present aims to explore how substituting shrimp meal with *Hermetia illucens* larvae meal in rainbow trout diets affects growth efficiency, feed utilization, and survival.

2. MATERIALS AND METHODS

The experimental protocol was developed according to the proposal submitted to the Nepal Agricultural Research Council and was approved by the Council to conduct the research. The feeding trial was carried out between May 1 to August 07, 2024, at the Outdoor Experimental Facility of the Fishery Research Station, Trishuli (FRS) located at Bidur-9 Nuwakot, Nepal.

2.1 Test Ingredients and Experimental Diets

The dried BSFL was obtained from J17 AQUA, a company specializing in fish food and feed supply, located in Kanchipuram, Tamil Nadu, India. Upon receipt, the dried insect larvae were processed into a fine powder before being incorporated into the experimental diets.

Five experimental diets were formulated to be isonitrogenous (Crude protein about 45 g/100g as fed) and isolipidic (ether extract about 15g/100g as fed) to fulfill the nutritional needs of rainbow trout. The control diet (CON) contained no BSFL meal, while four experimental diets incorporated partially defatted BSFL meal at varying levels. These experimental diets, labeled B25, B50, B75, and B100, substituted 25, 50, 75, and 100% of the Shrimp meal protein in the control diet with BSFL meal, respectively. Each diet were distributed randomly among three tanks.

The experimental diets were prepared at the Feed House of FRS. All the powdered ingredients and soyabean oil were individually weighed and mixed with blender followed by the addition of 250-500 ml/kg of water to the mixture. The feed pelleting machine was used to form pellet of 2.0 mm which was subsequently dried and stored in airtight buckets until used. The list of ingredients used to prepare experimental diets and the proximate composition of both the dry BSFL meal, and the experimental diets used in the study is presented in Table 1. The proximate composition and energy level of the BSF meal was provided by the BSFL supplier and experimental diets were measured in duplicate at the Animal Nutrition Laboratory at the Khumaltar Lalitpur.

2.2 Fish and Rearing Conditions

A 90-day feeding trial was conducted with the juvenile of rainbow trout obtained from own hatchery of the FRS, Trishuli. Since the fish were from the same hatchery, the acclimatization period was 5 days during which a locally made trout feed was fed (44 g/kg CP). Then a total of 750 healthy fish of mean individual body weight 5.39 ± 0.55 g was randomly divided into 15 flowthrough 300-L raceway tanks (three replicate tanks per diets). River water coming from siltation tank (average water temperature of $19.48 \pm 0.27^\circ\text{C}$) was supplied in flow-through raceways with each tank having a water flow of 6-8 L/min. Dissolved oxygen and pH were

measured every 2-3 days and were recorded to be 7.94±0.10 mg/L and 8.51±0.02 respectively. Feed was distributed by hand twice a day until apparent satiation. Apparent satiation was confirmed when the fish stopped eating. Mortality was checked every day. To monitor growth and through cleaning of the tanks, fish were weighed in bulk every 15 days.

2.3 Growth Efficiency

At the end of the feeding trial, after starving for 24 hours, all fish were harvested, counted and individually weighed for body weight and measured for total length. The following production efficiency indexes were calculated:

$$\text{Mortality (\%)} = (\text{number of dead fish}/\text{initial number of fish}) \times 100$$

$$\text{Individual weight gain (iWG, g)} = \text{average individual final body weight (iFBW, g)} - \text{average individual initial body weight (iIBW, g)}$$

$$\text{Specific growth rate (SGR, \%/day)} = [(\ln\text{FBW} - \ln\text{IBW})/\text{number of days}] \times 100$$

$$\text{Feed conversion ratio (FCR)} = \text{total feed supplied (g, DM)} / \text{WG (g)}$$

$$\text{Protein efficiency ratio (PER)} = \text{WG(g)}/\text{total protein fed (g, DM)}$$

Above mentioned indexes were calculated for each tank using individual initial and final body weight, and number of fish.

2.4 Statistical Analysis

To analyze the data IBM SPSS Statistics v. 25.0 for Windows was used. A one-way ANOVA test was used to compare data across experimental groups taking a statistical significance level at $P < 0.05$. Firstly, normality and homogeneity of variances were evaluated using the Kolmogorov–Smirnov test, and through Levene’s test respectively. Instead of F-test, we used Brown-Forsythe statistic test to compare group means when the assumption of equal variances was violated. Then, pairwise comparison was done by Tukey’s test for equal variances or Tamhane’s T2 for unequal variances (one-way ANOVA). Results are presented as mean and pooled standard error of the mean (SEM).

While there is no specific ethical protocols for fish research in Nepal, all experimental procedures were conducted with care to ensure minimal or no harm to the fish, adhering to scientific best practices and welfare standards.

Table 1. Ingredients and proximate composition of BSFL meal and experimental diets.

Ingredients	BSFL meal	CON	BSF25	BSF50	BSF75	BSF100
Shrimp meal		500	375	225	125	0
BSFL meal			125	225	375	500
Soybean full fat	355		355	355	355	355
Wheat Gluten	30		35	80	60	70
Corn meal	20		20	5	5	15
Wheat flour	50		50	80	65	50
Soyabean oil	40		35	25	10	5
Vitamin premix	3		3	3	3	3
Trace mineral premix	2		2	2	2	2
Total		1000	1000	1000	1000	1000
Proximate composition						
Dry matter (%)	91.67	93.81	93.39	93.00	92.66	92.34
Crude protein (%)	51.56	45.71	45.21	45.23	45.40	45.19
Lipid (%)	15.39	15.25	15.78	15.26	15.31	15.86
Digestible energy (MJ/kg)	26.46	18.75	19.90	20.89	21.92	23.10

Vitamin premix (IU or mg/kg diet): DL-α tocopherol acetate 60 IU; sodium menadione bisulphate 5 mg; retinyl acetate 15,000 IU; DL-cholecalciferol 3000 IU; thiamin 15 mg; riboflavin 30 mg; pyridoxine 15 mg; vitamin B12 0.05 mg; nicotinic acid 175 mg; folic acid 500 mg; inositol 1000 mg; biotin 2.5 mg; calcium pantothenate 50 mg
Mineral premix (g or mg/kg diet): Bicalcium phosphate 500 g, calcium carbonate 215 g, sodium salt 40 g, potassium chloride 90 g, magnesium chloride 124 g, magnesium carbonate 124 g, iron sulfate 20 g, zinc sulfate 4 g, copper sulfate 3 g, potassium iodide 4 mg, cobalt sulfate 20 mg, manganese sulfate 3 g, sodium fluoride 1 g

3. RESULTS

3.1 Experimental Diets

The chemical composition of all the experimental diets were found to be similar in terms of dry matter, crude protein, lipid content and gross energy as expected (Table 1). The results showed that the formulated experimental diets covered the nutritional requirements of the rainbow trout.

3.2 Growth Efficiency

The results of the growth efficiency parameters of Rainbow trout in each treatment group (CON, B25, B50, B75, and B100) are shown in Table 2.

The results show that initial body weight (iBW) of the rainbow trout did not significantly differ between the five treatment groups ($P > .05$) and ranged from 5.35 g to 5.43 g at the start of the experiment. The individual final body weight (iFBW) and individual weight gain (iWG) of rainbow trout were significantly higher in the B25, B50, and B75 groups compared to the control (CON) group ($P < .05$). The highest iFBW and iWG were observed in the B50 group. However, above B50, the iFBW and iWG decreased, with the B100 group showing iFBW and iWG values comparable to those of the CON group. The highest and lowest individual body weights were 23.97 g and 33.54 g respectively. Similarly, the highest and lowest individual weight gain were 28.19 g and 18.51 g respectively. Similarly, the specific growth rate (SGR) was significantly higher in B25, B50, and B75 groups than CON group ($P < .05$). Among the treatments, the highest SGR was observed in the B50 group, followed by B25 and B75 groups. The B100 group showed an SGR comparable to the CON group. The fish in the experiment had an SGR value from 1.67 %/day to 2.05 %/day. Likewise, the feed conversion ratio (FCR) was apparently lower in B25 and B50 groups compared to the CON and B75 group. However, there were no significant differences in FCR between the control group and the treatment groups except for B100. The B100 group had significantly higher FCR among the treatment groups in this experiment. The value for FCR in this experiment ranged from 1.19 to 2.74. Moreover, the result for PER revealed varying trends in values ranging from 0.92 to 2.04. The B25 group showed the statistically highest value among all groups, although its value was not statistically different from B50 group. The B50 group had statistically

similar PER to both CON and B75 groups, indicating similar protein efficiency ratio. The B100 group was observed with statistically lowest PER value among all the groups ($P < .05$) in this experiment.

Finally, the result of mortality rate revealed that groups B25 and B75 had the lowest mortality rate, although this rate was not statistically different from B75 and CON groups. However, significantly higher mortality rate was observed in B100 group compared to other treatments ($P < .05$). The mortality rate in this experiment ranged from 15 to 40%.

4. DISCUSSION

Several studies have demonstrated that insect-based protein sources, especially black soldier fly (BSF), are usable alternatives to the conventional protein sources like fish meal in aquafeed (Bondari and Sheppard, 1981; Hartviksen et al., 2014; Li et al., 2015). The application of BSF as a possible source of protein in trout feeds has been amply studied (Bolton et al., 2021; Borland et al., 2024; Caimi et al., 2021; Gasco et al., 2015; Hoc et al., 2021). Result of our study demonstrated that the diet B25 and B50, in which the shrimp meal was replaced by 25 and 50% BSFL meal respectively, resulted in the best results for most of the production parameters compared to control group, and the replacement up to 75% were possible without negative consequences. This result agrees with study of (Sealey et al., 2011), showing BSF meal could replace up 50% shrimp meal in the diet of rainbow trout with no adverse effect on growth efficiency. Gasco et al. reported that feeding rainbow trout with defatted BSF successfully substituted 25 and 50% of the fish meal protein without affecting growth efficiencies and quality parameters (Gasco et al., 2015). In line with our results, a study done by (Muin et al., 2017) (Muin et al., 2017) reported that BSF meal can replace up to 50% fish meal on Nile tilapia (*Oreochromis niloticus*) diet with no adverse effects on growth and feed utilization.

Similarly, a study in yellow catfish (*Pelteobagrus fulvidraco*) reported growth efficiency with up to 48% fishmeal protein replaced by BSF larvae meal protein (Xiao et al., 2018). In our study, there were slight significant differences observed in terms of growth efficiency and feed utilization among all the experimental diets. Final weight, weight gain, specific growth rate, feed conversion ratio, and protein efficiency ratio were

Table 2. Effect of BSF larvae meal on growth efficiency of rainbow trout (n=3)

	CON	B25	B50	B75	B100	SEM	P-value
ilBW, g	5.37 ^a	5.37 ^a	5.35 ^a	5.43 ^a	5.43 ^a	0.11	1.000
iFBW, g	23.87 ^c	31.54 ^{ab}	33.54 ^a	29.53 ^b	26.23 ^c	0.96	0.000
iWG, g	18.51 ^c	26.17 ^{ab}	28.19 ^a	24.11 ^b	20.80 ^c	0.95	0.000
SGR, %/d	1.67 ^c	1.97 ^{ab}	2.05 ^a	1.88 ^{ab}	1.75 ^{bc}	0.04	0.004
FCR	1.53 ^a	1.19 ^a	1.31 ^a	1.59 ^a	2.74 ^b	0.16	0.000
PER	1.56 ^b	2.04 ^a	1.83 ^{ab}	1.51 ^b	0.92 ^c	0.11	0.000
Mortality, %	19.00 ^a	15.00 ^a	15.00 ^a	22.00 ^a	40.00 ^b	2.73	0.001

BSF or B: black soldier Fly, SEM: standard error of the mean, ilBW: individual initial body weight, iFBW: individual final body weight, iWG: individual weight gain, SGR: specific growth rate, FCR: feed conversion ratio, PER: protein efficiency ratio

significantly improved in fish feed diet B25 and B50 compared to that in other groups. However, these parameters, especially weight gain and SGR were observed to decrease as the level of BSF larvae meal increased further. There were no significant differences recorded for FCR up to 75% replacement level. However, the 25% replacement diet (B25) generated the best FCR and PER values. The lower FCR and higher PER value indicate the good quality of feed and optimum utilization of feed. This indicates that rainbow trout could utilize the BSF larvae meal in its diet optimally up to 50% of shrimp meal replacement. Slightly different findings were reported in a previous study where author found no significant difference in FCR and PER up to 75% replacement level, improved weight gain and SGR when fish meal in rainbow trout feed was replaced by BSF prepupal meal (Hoc et al., 2021). However, some previous studies reported that above 25% replacement of the fish meal by BSF larvae or prepupal meal, the FCR and weight gain in rainbow trout adversely affected (Gasco et al., 2015; Hoc et al., 2021; Sealey et al., 2011; Muin et al., 2017; Xiao et al., 2018; St-Hilaire et al., 2007). In a study in red drum (*Sciaenops ocellatus*) author recorded impaired growth efficiency and feed utilization when BSF larvae meal was used as low as 150 g/kg (Yamamoto et al., 2022).

Such contrasting results might be due to difference in the species of fish used in the experiment, level and life stage of BSF used in the fish feed, and different in the rearing substrate used for culture of BSF. (Yamamoto et al., 2022) suspected that impaired growth and poor nutrient utilization observed in red drum compared to previous studies was due to difference in nutrient content in BSF larvae rearing in different rearing substrate because Yamamoto et al. found that BSF larvae meal derived from larvae fed a commercial diet

resulted in better growth efficiency in red drum compared to BSF larvae meal from larvae raised on brewers' spent grains, even though both diets were formulated to be isonitrogenous. Similarly, (Kroeckel et al., 2012) concluded that the feed intake, availability and nutrient digestibility of juvenile turbot (*Psetta maxima*) was impaired due to presence of higher amount of chitin in diet coming from older BSF larvae ultimately affecting the growth efficiency. Although, why results were obtained is unclear, St-Hilaire et al. mentioned that higher level of BSF meal in the fish feed could increase the chitin content in the fish feed which make feed less digestible for fish and ultimately affecting growth efficiency. Chitin, a primary component of crustacean exoskeletons and fungal cell walls, is a tough, nitrogen-containing polysaccharide with similar structure to cellulose, but more resistant to degradation and digestion by most animals, except few fish (Nunes et al., 2018).

There are several previous studies which shows that higher level of insect or chitin in fish feed, affects the digestibility and growth efficiency of fish regardless of the type of insect, rearing substrate used to culture insect, and species of fish in the experiment. Similar finding was reported in a study with rainbow trout where amount of tissue was decreased as the amount of insect (*Tenebrio molitor*) meal increased in the trout feed (Belforti et al., 2016). However, such result might have been obtained due to difference in ability of fish to utilize the fat from different sources, composition and amount in the diets (Watanabe et al., 1982) because of fatty acid composition of the BSF larvae meal used in preparation of experimental diets that ultimately affected fat metabolism of rainbow trout. According to previous research, activities of fatty acid synthase (FAS) and lipase (LPS), two lipid metabolism-related enzymes in fish, are closely related to composition of chitin in BSF (Yu et al.,

2023). Li et al reported that juvenile Jian Carp (*Cyprinus carpio* var. Jian) when fed with higher concentration of BSF larvae showed poor nutrient utilization and impacted ability of synthesizing the fatty acids (Li et al., 2016).

Therefore, a reasonable explanation for reduced growth efficiency and nutrient utilization above 50% replacement level in present study could be higher insect component in diet causing impaired activities of lipid metabolism-related enzymes, closely related to the high amount of chitin in BSF. In addition, a study reported that chitin digestibility is usually very low or completely absent as in rainbow trout (Lindsay, 1984). Similar conclusion was made in a study in which authors found above 3% chitin in a diet of rainbow trout negatively affected growth efficiency and nutrient digestibility (Pascon et al., 2024).

Despite these negative effects of high level of insect meal or chitin content above 2-3% in fish feed, insect meal contains several bioactive compounds and have positive impacts on fish metabolism (Gasco et al., 2020), resistant against pathogens (Udayangani et al., 2017) and immune response (Ringø et al., 2012). And chitin is one of the most studied bioactive compounds in aquafeed having such beneficial effects. The BSFL chitin at 0.4% was shown to promote the monounsaturated fatty acid deposition in muscle and improved health status of largemouth bass (*Micropterus salmoides*) (Hu et al., 2024). Previous studies have shown that the peptide extract of the BSF larvae possess broad-spectrum of antibacterial activity suggesting BSF as a potential source of novel antibiotic-like compounds which may control infections in fish (Elhag et al., 2017; Park et al., 2014; Zhou et al., 2013;). Studies also report that chitin, analogous to cellulose, acts as a prebiotic, promoting the growth of beneficial bacteria and chitin-degrading bacteria in the gut, resulting in the production of healthy short-chain fatty acids (SCFAs), including acetic, propionic, and butyric acids, which have been demonstrated to have beneficial effects on the gut, immune system and health of fish including rainbow trout (Gaudioso et al., 2021; Kipkoech and Beyond, 2023; Xiao et al., 2018). These positive responses could be due to the presence of chitin in the fish feed.

Previous studies have demonstrated the ability of fish to degrade chitin fiber through the action of chitinolytic enzymes (chitinase and chitobiase) (Fines and Holt, 2010; Holen et al., 2022). This knowledge can be applied to improve feed

digestibility and nutrient absorption in fish by incorporating chitin-degrading enzymes or bacteria into their diet. Specifically, chitinases can enhance the accessibility of digestive enzymes to entrapped proteins or lipids, leading to improved overall digestive health and nutrient absorption in fish (Rangel et al., 2022; Subramanian et al., 2022).

The cumulative evidence from the present study and existing research in the field collectively suggests that the incorporation of Black Soldier Fly (BSF) larvae meal into fish feed can yield several benefits, including improved growth efficiency, enhanced immune function, and better overall health of fish. Therefore, the use of BSF as a feed source in rainbow trout presents a viable alternative to traditional protein-rich ingredients like shrimp meal. Importantly, the production of BSF larvae using organic waste offers a potentially sustainable solution for the aquaculture industry ultimately reducing feed cost and leading to significant cost savings in commercial aquaculture. However, additional research is required to determine the optimal inclusion level of BSF larvae meal in fish feed and to develop effective processing methods to reduce the issues related to higher chitin content and maximize level of inclusion of BSF larvae in fish feed without affecting production efficiency.

5. CONCLUSION

In conclusion, this study demonstrates that diets B25 (25% replacement) and B50 (50% replacement) of shrimp meal with Black Soldier Fly (BSF) larvae meal significantly enhance growth efficiency and feed utilization in juvenile rainbow trout, identifying 25-50% inclusion as the optimal range. Conversely, higher inclusion levels (75-100%) resulted in reduced growth and increased mortality, emphasizing the importance of balanced inclusion levels. These findings highlight BSF larvae meal as a sustainable and effective alternative protein source for rainbow trout farming, with relevance for promoting aquaculture in Nepal's mid-hill regions. Future studies should investigate the digestibility of this innovative feed ingredient to further refine its application.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image

generators have been used during writing or editing of this manuscript.

ACKNOWLEDGEMENT

This work was supported by the Nepal Agricultural Research Council and funded by the Government of Nepal.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Mahato, I. S., Paudel, K., & Shrestha, A. (2024). Comparative growth performance of genetically improved, Chinese, and local strains of Rainbow trout (*Oncorhynchus mykiss*) in mid-hill Nepal. *Asian Journal of Fisheries and Aquatic Research*, 26, 44-51.
- FRS. (2023). *Annual Report of the fiscal year 2022/23 (2079/80)*. Nuwakot Nepal: Fishery Research Station, Trishuli.
- Mahato, I. S., Timalina, P., Paudel, K., Shrestha, A., Bhusal, C., & Kunwar, P. (2023). Effects of replacing dietary shrimp meal and soybean meal with silkworm (*Bombyx mori*) pupae meal on growth performance of rainbow trout *Oncorhynchus mykiss*. *International Journal of Fisheries and Aquatic Studies*, 11, 21-25.
- Pauly, D., & Christensen, V. (1995). Primary production required to sustain global fisheries. *Nature*, 374, 255-257.
- Mohsen, A. A., & Lovell, R. T. (1990). Partial substitution of soybean meal with animal protein sources in diets for channel catfish. *Aquaculture*, 90, 303-311.
- Calvert, C. C., Martin, R. D., & Morgan, N. O. (1969). House fly pupae as food for poultry. *Journal of Economic Entomology*, 62, 938-939.
- Bartucz, T., Csokas, E., Nagy, B., Gyurcsak, M. P., Bokor, Z., Bernath, G., et al. (2023). Black soldier fly (*Hermetia illucens*) meal as direct replacement of complex fish feed for rainbow trout (*Oncorhynchus mykiss*) and African catfish (*Clarias gariepinus*). *Life (Basel)*, 13.
- Nugroho, R. A., & Nur, F. M. (2018). Insect-based protein: Future promising protein source for fish cultured. *IOP Conference Series: Earth and Environmental Science*, 144.
- Henry, M., Gasco, L., Piccolo, G., & Fountoulaki, E. (2015). Review on the use of insects in the diet of farmed fish: Past and future. *Animal Feed Science and Technology*, 203, 1-22.
- Meneguz, M., Schiavone, A., Gai, F., Dama, A., Lussiana, C., Renna, M., et al. (2018). Effect of rearing substrate on growth performance, waste reduction efficiency and chemical composition of black soldier fly (*Hermetia illucens*) larvae. *J Sci Food Agric*, 98, 5776-5784.
- Xu, X., Ji, H., Yu, H., & Zhou, J. (2020). Influence of dietary black soldier fly (*Hermetia illucens* Linnaeus) pulp on growth performance, antioxidant capacity and intestinal health of juvenile mirror carp (*Cyprinus carpio* var. *specularis*). *Aquaculture Nutrition*, 26, 432-443.
- Fisher, H. J., Collins, S. A., Hanson, C., Mason, B., Colombo, S. M., & Anderson, D. M. (2020). Black soldier fly larvae meal as a protein source in low fish meal diets for Atlantic salmon (*Salmo salar*). *Aquaculture*, 521.
- Fawole, F. J., Adeoye, A. A., Tihamiyu, L. O., Ajala, K. I., Obadara, S. O., & Ganiyu, I. O. (2020). Substituting fishmeal with *Hermetia illucens* in the diets of African catfish (*Clarias gariepinus*): Effects on growth, nutrient utilization, haemato-physiological response, and oxidative stress biomarker. *Aquaculture*, 518.
- Lock, E. R., Arsiwalla, T., & Waagbø, R. (2016). Insect larvae meal as an alternative source of nutrients in the diet of Atlantic salmon (*Salmo salar*) postsmolt. *Aquaculture Nutrition*, 22, 1202-1213.
- Magalhães, R., Sánchez-López, A., Leal, R. S., Martínez-Llorens, S., Oliva-Teles, A., & Peres, H. (2017). Black soldier fly (*Hermetia illucens*) pre-pupae meal as a fish meal replacement in diets for European seabass (*Dicentrarchus labrax*). *Aquaculture*, 476, 79-85.
- Biasato, I., Chemello, G., Odon, S. B., Ferrocino, I., Corvaglia, M. R., Caimi, C., et al. (2022). *Hermetia illucens* meal inclusion in low-fishmeal diets for rainbow trout (*Oncorhynchus mykiss*): Effects on the growth performance, nutrient digestibility coefficients, selected gut health traits, and health status indices. *Animal Feed Science and Technology*, 290.

- Terova, G., Rimoldi, S., Ascione, C., Gini, E., Ceccotti, C., & Gasco, L. (2019). Rainbow trout (*Oncorhynchus mykiss*) gut microbiota is modulated by insect meal from *Hermetia illucens* prepupae in the diet. *Reviews in Fish Biology and Fisheries*, 29, 465-486.
- Bruni, L., Randazzo, B., Cardinaletti, G., Zarantoniello, M., Mina, F., Secci, G., et al. (2020). Dietary inclusion of full-fat *Hermetia illucens* prepupae meal in practical diets for rainbow trout (*Oncorhynchus mykiss*): Lipid metabolism and fillet quality investigations. *Aquaculture*, 529.
- Bondari, K., & Sheppard, D. C. (1981). Soldier fly larvae as feed in commercial fish production. *Aquaculture*, 24, 103-109.
- Hartviksen, M., Bakke, A. M., Vecino, J. G., Ringo, E., Krogdahl, A. (2014). Evaluation of the effect of commercially available plant and animal protein sources in diets for Atlantic salmon (*Salmo salar* L.): digestive and metabolic investigations. *Fish Physiol Biochem*, 40, 1621-1637.
- Li, L., Liang, X. F., He, S., Sun, J., Wen, Z. Y., He, Y. H., et al. (2015). Transcriptome analysis of grass carp (*Ctenopharyngodon idella*) fed with animal and plant diets. *Gene*, 574, 371-379.
- Bolton, C., Muller, N., Hyland, J., Johnson, M., Valente, C. S., Davies, S., et al. (2021). Black soldier fly larval meal with exogenous protease in diets for rainbow trout (*Oncorhynchus mykiss*) production meeting consumer quality. *Journal of Agriculture and Food Research*, 6, 100232.
- Borland, M., Riesenbach, C., Shandilya, U., Chiasson, M., Karrow, N., Huyben, D. (2024). Growth performance, hepatic gene expression, and plasma biochemistry of rainbow trout fed full-fat meal, defatted meal, oil, and chitin from black soldier flies. *Comparative Immunology Reports*, 200149.
- Caimi, C., Biasato, I., Chemello, G., Oddon, S. B., Lussiana, C., Malfatto, V. M., et al. (2021). Dietary inclusion of a partially defatted black soldier fly (*Hermetia illucens*) larva meal in low fishmeal-based diets for rainbow trout (*Oncorhynchus mykiss*). *Journal of Animal Science and Biotechnology*, 12, 1-15.
- Gasco, L., Stas, M., Schiavone, A., Rotolo, L., De Marco, M., Dabbou, S., et al. (2015). Use of black soldier fly (*Hermetia illucens*) meal in rainbow trout (*Oncorhynchus mykiss*) feeds. *Aquaculture Europe Meeting 2015 "Acquaculture, Nature and Society" Session: Nutrition focus on Insect meal*.
- Hoc, B., Tomson, T., Malumba, P., Blecker, C., Jijakli, M. H., Purcaro, G., et al. (2021). Production of rainbow trout (*Oncorhynchus mykiss*) using black soldier fly (*Hermetia illucens*) prepupae-based formulations with differentiated fatty acid profiles. *Science of the Total Environment*, 794, 148647.
- Sealey, W. M., Gaylord, T. G., Barrows, F. T., Tomberlin, J. K., McGuire, M. A., Ross, C., et al. (2011). Sensory analysis of rainbow trout (*Oncorhynchus mykiss*), fed enriched black soldier fly prepupae, *Hermetia illucens* (L.), meal. *Aquaculture Research*, 42, 1549-1555.
- Muin, H., Taufek, N., Kamarudin, M., & Razak, S. (2017). Growth performance, feed utilization and body composition of Nile tilapia, *Oreochromis niloticus* (Linnaeus, 1758) fed with different levels of black soldier fly, *Hermetia illucens* (Linnaeus, 1758) maggot meal diet. *Iranian Journal of Fisheries Sciences*, 16, 567-577.
- Xiao, X., Jin, P., Zheng, L., Cai, M., Yu, Z., Yu, J., et al. (2018). Effects of black soldier fly (*Hermetia illucens*) larvae meal protein as a fishmeal replacement on the growth and immune index of yellow catfish (*Pelteobagrus fulvidraco*). *Aquaculture Research*, 49, 1569-1577.
- St-Hilaire, S., Sheppard, C., Tomberlin, J. K., Irving, S., Newton, L., McGuire, M. A., et al. (2007). Fly prepupae as a feedstuff for rainbow trout, *Oncorhynchus mykiss*. *Journal of the World Aquaculture Society*, 38, 59-67.
- Yamamoto, F. Y., Suehs, B. A., Ellis, M., Bowles, P. R., Older, C. E., Hume, M. E., et al. (2022). Dietary fishmeal replacement by black soldier fly larvae meals affected red drum (*Sciaenops ocellatus*) production performance and intestinal microbiota depending on what feed substrate the insect larvae were offered. *Animal Feed Science and Technology*, 283.
- Kroeckel, S., Harjes, A. G. E., Roth, I., Katz, H., Wuertz, S., Susenbeth, A., et al. (2012). When a turbot catches a fly: Evaluation of a pre-pupae meal of the black soldier fly (*Hermetia illucens*) as fish meal substitute—Growth performance and chitin degradation in juvenile turbot (*Psetta maxima*). *Aquaculture*, 364-365, 345-352.

- Nunes, C. S., & Philipps-Wiemann, P. (2018). Chapter 18 - Chitinases. In C. S. Nunes & V. Kumar (Eds.), *Enzymes in Human and Animal Nutrition* (pp. 361-378). Academic Press.
- Belforti, M., Gai, F., Lussiana, C., Renna, M., Malfatto, V., Rotolo, L., et al. (2016). *Tenebrio Molitor* meal in rainbow trout (*Oncorhynchus mykiss*) diets: Effects on animal performance, nutrient digestibility and chemical composition of fillets. *Italian Journal of Animal Science*, 14.
- Watanabe, T. (1982). Lipid nutrition in fish. *Comparative Biochemistry and Physiology Part B: Comparative Biochemistry*, 73, 3-15.
- Yu, Z., Sun, Z., Ou, B., Zhou, M., Huang, Y., & Tan, X. (2023). Effects of partial replacement of fish meal with black soldier fly (*Hermetia illucens*) larvae meal on growth performance, lipid metabolism and hepatointestinal health of juvenile golden pompano (*Trachinotus ovatus*). *Aquaculture Reports*, 33.
- Li, S., Ji, H., Zhang, B., Tian, J., Zhou, J., & Yu, H. (2016). Influence of black soldier fly (*Hermetia illucens*) larvae oil on growth performance, body composition, tissue fatty acid composition and lipid deposition in juvenile Jian carp (*Cyprinus carpio* var. Jian). *Aquaculture*, 465, 43-52.
- Lindsay, G. J. H. (1984). Adsorption of rainbow trout (*Salmo gairdneri*) gastric lysozymes and chitinase by cellulose and chitin. *Aquaculture*, 42, 241-246.
- Pascon, G., Cardinaletti, G., Daniso, E., Bruni, L., Messina, M., Parisi, G., et al. (2024). Effect of dietary chitin on growth performance, nutrient utilization, and metabolic response in rainbow trout (*Oncorhynchus mykiss*). *Aquaculture Reports*, 37.
- Gasco, L., Biancarosa, I., & Liland, N. S. (2020). From waste to feed: A review of recent knowledge on insects as producers of protein and fat for animal feeds. *Current Opinion in Green and Sustainable Chemistry*, 23, 67-79.
- Udayangani, R. M. C., Dananjaya, S. H. S., Nikapitiya, C., Heo, G. J., Lee, J., & De Zoysa, M. (2017). Metagenomics analysis of gut microbiota and immune modulation in zebrafish (*Danio rerio*) fed chitosan silver nanocomposites. *Fish Shellfish Immunol.*, 66, 173-184.
- Ringø, E., Zhou, Z., Olsen, R. E., & Song, S. K. (2012). Use of chitin and krill in aquaculture - the effect on gut microbiota and the immune system: a review. *Aquaculture Nutrition*, 18, 117-131.
- Hu, Z., Xia, M., Wang, G., Jia, L., Ji, H., Sun, J., et al. (2024). A superior chitin product: Black soldier fly larvae chitin, beneficial to growth performance, muscle quality and health status of largemouth bass *Micropterus salmoides*, in comparison to shrimp chitin. *Aquaculture*, 595.
- Elhag, O., Zhou, D., Song, Q., Soomro, A. A., Cai, M., Zheng, L., et al. (2017). Screening, expression, purification and functional characterization of novel antimicrobial peptide genes from *Hermetia illucens* (L.). *PLOS One*, 12, e0169582.
- Park, S. I., Chang, B. S., & Yoe, S. M. (2014). Detection of antimicrobial substances from larvae of the black soldier fly, *Hermetia illucens* (Diptera: Stratiomyidae). *Entomological Research*, 44, 58-64.
- Zhou, Z., Karlsen, Ø., He, S., Olsen, R. E., Yao, B., & Ringø, E. (2013). The effect of dietary chitin on the autochthonous gut bacteria of Atlantic cod (*Gadus morhua* L.). *Aquaculture Research*, 44, 1889-1900.
- Gaudioso, G., Marzorati, G., Faccenda, F., Weil, T., Lunelli, F., Cardinaletti, G., et al. (2021). Processed animal proteins from insect and poultry by-products in a fish meal-free diet for rainbow trout: Impact on intestinal microbiota and inflammatory markers. *International Journal of Molecular Sciences*, 22.
- Kipkoech, C. (2023). Beyond proteins—Edible insects as a source of dietary fiber. *Polysaccharides*, 4, 116-128.
- Fines, B. C., & Holt, G. J. (2010). Chitinase and apparent digestibility of chitin in the digestive tract of juvenile cobia, *Rachycentron canadum*. *Aquaculture*, 303, 34-39.
- Holen, M. M., Sandve, S. R., Harvey, T. N., Jin, Y., Angell, I. L., Rudi, K., et al. (2022). The effect of dietary chitin on Atlantic salmon (*Salmo salar*) chitinase activity, gene expression, and microbial composition. *BioRxiv The Preprint Server for Biology*, 1.
- Rangel, F., Santos, R. A., Monteiro, M., Lavrador, A. S., Gasco, L., Gai, F., et al. (2022). Isolation of chitinolytic bacteria from European sea bass gut microbiota fed diets with distinct insect meals. *Biology*, 11, 964.

Subramanian, K., Balaraman, D., Panangal, M., Nageswara Rao, T., Perumal, E., R. A., et al. (2022). Bioconversion of chitin waste through *Stenotrophomonas maltophilia* for production of chitin derivatives as a seabass enrichment diet. *Sci Rep*, 12, 479.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/127454>