

## Tree Species Diversity of Ikogosi Warm Spring and Arinta Waterfall Watersheds, Nigeria: Implication for Sustainable Ecotourism

O. J. Olujobi<sup>1\*</sup>, O. B. Adeleke<sup>1</sup> and J. O. Orimaye<sup>1</sup>

<sup>1</sup>Department of Forest Resources and Wildlife Management, Ekiti State University, P.M.B. 5363, Ado-Ekiti, Nigeria.

### Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

### Article Information

DOI: 10.9734/AJRAF/2019/v3i230033

#### Editor(s):

(1) Dr. Cengiz Yucedag, Professor, Department of Landscape Architecture, Mehmet Akif Ersoy University, Turkey.

(2) Dr. Nebi Bilir, Professor, Forestry Faculty, Isparta University of Applied Sciences, Isparta, Turkey.

#### Reviewers:

(1) Diogo José Oliveira Pimentel, Rural Federal University of Pernambuco, Brazil.

(2) Rozainah Mohamad Zakaria, University of Malaya, Malaysia.

(3) M. Iqbal Zuberi, Gono University, Bangladesh.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/47239>

Original Research Article

Received 25 January 2019  
Accepted 10 April 2019  
Published 19 April 2019

### ABSTRACT

An in-depth knowledge of the richness, diversity and species composition of plant community is vital for providing information for planning and sustainable utilization. This study assesses the diversity of tree species of Ikogosi warm spring and Arinta waterfall watersheds in Ekiti State, Nigeria. Direct observation and vegetation assessment were used for data collection in two hectares (ha) of land divided into four plots of 50 m by 50 m in each of the two study sites (Ikogosi warm spring and Arinta waterfall). Two plots were diagonally selected within each hectare. All living trees of basal diameter  $\geq 10$  cm were identified and classified into families. Shannon-Weiner diversity index, species evenness, relative density (RD), relative dominance ( $RD_0$ ) and importance value index (IVI) were used to assess and compare tree species diversity and abundance. Sorenson's coefficient was used to compare sites for overlapping of similarity. The results revealed that seventy eight (78) species and 25 families were recorded in both watersheds with family Malvaceae having the highest species density (15). *Malacantha alnifolia* (5) and *Voacanga africana*

\*Corresponding author: E-mail: [olujobioj@yahoo.com](mailto:olujobioj@yahoo.com);

(5) were the species most frequently encountered. Species diversity indices revealed vegetation with very high tree species diversity and abundance in the two study sites. Species evenness value revealed even distribution in Arinta waterfall than in Ikogosi warm spring. Diameter and height distribution of trees at the two watersheds indicates a forest structure that is immature and still expanding. *Anthocleista vogelii*, has the highest value of RD<sub>0</sub> (15.63) and IVI (10.6) respectively. The study revealed that some species such as *Anthonotha macrophylla*, *Aningeria robusta*, *Bridelia atroviridis*, among others are threatened and endangered. Consequently, it was recommended that management strategies should be put in place to improve status of the watersheds while conservation efforts should be stepped up for species with rarity index value to prevent them from going into extinction.

**Keywords:** Species diversity; Ikogosi warm spring; Arinta waterfall; Sustainable ecotourism; Ekiti State; Nigeria.

## 1. INTRODUCTION

Tree as a major component of forest is an important natural resource capable of producing diverse tangible products and also provides services such as game viewing, recreation and tourism opportunities. It is a means of income generation for government and providing employment opportunities for people [1]. Forest trees help in checking Global warming by acting an effective sink for the CO<sub>2</sub>. Recent discoveries has vividly revealed that most of our protected areas are of sub-climax ecosystems for which management is essential if the desired characteristics of the ecosystem are to be maintained [2]. Even in those regions of the world where protected areas contain quite natural self-regenerating climax ecosystems, there is still sometimes a need for management in order to enhance goods and services derived from those forests [2].

Biodiversity refers to the variability among living organisms from all sources including, among other things, terrestrial and aquatic ecosystems. This variability includes within species and between species. Biodiversity is essential for human survival, economic well-being and ecosystem function and stability [3]. Habitat destruction, over exploitation; pollution and species introduction have been identified as major causes of biodiversity loss [4]. The disturbances created by these factors determine forest dynamics and tree diversity of an area [5]. An in-depth knowledge of the richness, diversity and species composition of plant community formation is vital for planning and conservation actions that can reduce the environmental impact of development on an ecosystem [6]. The diversity of tree species is fundamental to total forest biodiversity, because trees provide resources and habitat for almost all other forest

species. In natural resources management operations, inventories of biodiversity are used to determine the nature and distribution of biodiversity of an area being managed. This also determined the measure to be put in place in the management of such area to enable the resources to fulfill their potentials [7].

A viable ecotourism centre helps in community development by providing the alternative and sustainable source of livelihood to local community [8]. It has contributed to conservation of biodiversity; promotes small and medium tourism enterprises; stresses local participation, ownership and business opportunities, particularly for rural people; and above all includes the learning experiences [9]. Ecotourism centre has help in involving local community for conservation of the ecology and biodiversity which in turn provides the economic incentives to the local community; sustains the health and well-being of local people; involves responsible action on the part of tourist and the tourism industry [10].

Ecotourism being a nature based tourism which takes into account the natural ecological attraction, their conservation and development has attracted increasing attention in recent years, not only as an alternative to mass tourism, but also as a means of economic development and environmental conservation [11,12]. The aim of an ecosystem manager is to protect the environment, making it profitable to the community people by generating revenue, educating and serving the pleasure of tourists. In situ conservation is a veritable tool for the preservation of genetic resources currently decreasing at an alarming rate [13]. Before the development of any natural resource into an ecotourism sites, it is necessary to know much about the vegetation and other potentials

embedded in such an area as it will help to identify measures to be used for the conservation of the potentials of such area.

Forest in Ikogosi warm spring in Nigeria is valued for its high biodiversity; ecosystem and ecotourism importance [14]. Similarly the Arinta waterfall is also well-known for its ecotourism value [15]. However, the effort of the state government to upgrade the ecotourism status of the study area has led to the loss of endangered woody species and non-timber forest products thereby causing in-balance in the ecosystem of the site [16]. Although, the development of infrastructures is inevitable in as much as aesthetic values of the place is to be maximally and efficiently utilized, but measures should be put in place to ensure that the native vegetation species are conserved for posterity. Therefore, to manage these forests for ecotourism purpose on a sustainable level, there is the need to identify tree species and species diversity, land use practices to which the site is subjected to, and the conservation measure been put in place by the stake holders and government to properly manage the forest on a sustainable level to improve its ecotourism potential.

## 2. MATERIALS AND METHODS

### 2.1 Study Area

Ikogosi warm spring and Arinta waterfall watersheds are located in Ekiti West Local Government, Ekiti State, Nigeria. The two towns are about 5km distance apart and are situated between lofty; step sided and heavily wooded,

north-south trending hills underlain by metamorphic rock with undulating landscape [7]. Ikogosi warm spring (7° 35' 0" N., 4° 92' 0" E.) and Arinta waterfall (7° 40' 0" N.4° 59' 0" E.) sites have tropical climate of West Africa monsoonal type with two distinct rainy (April-October) and the dry (November-March) season. The annual rainfall ranged from 1,200 mm to 1,500 mm and temperature ranges between 21°C and 34°C with high humidity. The vegetation is of tropical rain forest [17].

### 2.2 Method of Data Collection

Direct observation and vegetation assessment were used for data collection. Inventory of infrastructural facilities at the study sites was carried out by observation while vegetation assessment was done by enumeration method as described by [14]. Two hectares (ha) of land (one close to places where visitors could easily assess and one in the undisturbed area) were measured in each of the study site. Each hectare was divided into four plots of 50 m by 50 m. Two plots were diagonally selected within each hectare to make a total of 4 plots in each study sites. Within each of the selected sampled plots, living trees of basal diameter  $\geq 10$  cm were identified with the help of a taxonomist using standard key. Where the tree's botanical name was not known it was identified by its commercial or local name and later translated to correct botanical names using standard key [18]. The diameter at breast height (dbh) was measured with diameter tape while height of each tree was measured with Spiegel relaskop.




Fig. 1. Map of Ekiti State showing the location of the study areas

## 2.3 Data Analyses

### Diversity Index

Species diversity index (H) was computed using the Shannon-Wiener diversity index given by [19]

$$H^1 = - \sum_{i=1}^S P_i \ln (P_i) \quad (1)$$

Where: H' = Shannon-Wiener diversity index; S = Total number of species in the site; P<sub>i</sub>= proportion of S made up of the i<sup>th</sup> species; ln = natural logarithm.

### Species Evenness

Species evenness in each community was determined using Shannon's equitability (E<sub>H</sub>) as stated by [20]

$$E = \frac{H^1}{\ln(S)} \quad (2)$$

S is the total number of species in each community.

$$\text{Sorenson's coefficient } (\beta) = \frac{2c}{s_1 + s_2} \quad (3)$$

Where

C = Total number of species in the two study areas

S<sub>1</sub> = Total number of species found in Ikogosi warm spring

S<sub>2</sub> = Total number of species found in Arinta waterfall

### Species Relative Density (RD)

Species relative density, which is an index for assessing species relative distribution [21], was computed with

$$RD = \left[ \frac{n_i}{N} \right] \times 100 \quad (4)$$

Where, RD (%) = species relative density, n<sub>i</sub> = number of individuals of species, N=total number of all individual trees of all species in the entire community.

### Species Relative Dominance (RDo)

Species relative dominance (RDo (%)), used in assessing relative space occupancy of a tree was estimated with

$$RDo = \frac{(\sum Ba_i \times 100)}{\sum Ba_n} \quad (5)$$

Where: Ba<sub>i</sub> = basal area of all trees belonging to a particular species, Ba<sub>n</sub> = basal area of all individual tree

### Importance Value Index

The Importance Value Index (IVI) of each species was computed with the relationship;

$$IVI = \frac{(RD + RD_0)}{2} \quad (6) [21]$$

All analysis were carried out using the computer model PAST version 3. Tree species were classified into families. Trees within strata of each forest were classified into three layers using method of [22].

### 3. RESULTS

#### 3.1 Infrastructural Facilities at the Ecotourism Centres and the Host Communities

The result in Table 1 shows that Ikogosi warm spring centre is well equipped with the presence of recreational facilities such as hotel, swimming pool, relaxation sport, fitness shop and multipurpose hall than the Arinta waterfall centre.

#### 3.2 Tree Species Composition in the Study Area

Table 2 shows that a total of 78 species were recorded in the two sites, however the vegetation in Arinta waterfall has higher number of species (66) than that of Ikogosi warm spring with 36 species. The vegetation in Arinta waterfall has

higher number of individual tree (158) than that in Ikogosi warm spring which had 72 tree species in the sampled plots. *Malacantha alnifolia* (5) and *Voacanga africana* (5) were the most frequently enchanter species while *Delonix regia* (3.8 cm) was the tree with the highest diameter at breast height (dbh) and *Ricinodendron heudelotii* (12.88 m) was tree species with the highest height. In Ikogosi warm spring, *Anthocleista vogelii*, *Antiaris africana*, *Delonix regia* were the most common species with frequency of 4 each while *Anthocleista vogelii* (1.64 cm) was tree species with highest dbh and *Brachystegia eurycoma* (12.72 m) with the highest height. In Arinta waterfall, *Malacantha alnifolia* and *Voacanga Africana* were the most common species with frequency of (5 each) while tree species with the highest dbh was *Delonix regia* (3.8 cm) and tree species with the highest height was *Ricinodendron heudelotii* (12.88 m). Table 2 also present detail results on species relative density (RD), relative dominance (RD<sub>0</sub>) and Importance Value Index (IVI) for the studied watersheds *Anthocleista vogelii*, *Delonix regia* and *Antiaris Africana* has the highest RD value with 5.6 each, while *Anthocleista vogelii* has the highest value of RD<sub>0</sub> (15.63) and IVI (10.6) respectively.

**Table 1. Facilities at the ecotourism centres and the host communities**

Infrastructural facilities	Ikogosi warm spring	Arinta waterfall
	Status	Status
Hotel and restaurants	√	√
Federal/State chalets	√	X
Relaxation sport	√	√
Swimming pool	√	X
Beauty centre	√	X
Fitness Shop	√	X
School	√	√
Health centre	√	√
Arts and crafts shops	√	X
Electricity	√	√
Shopping mall	√	X
Car parking facility	√	√
Staff quarters	√	X
Worship centre	√	√
Market	√	√
Road	√	√
Concrete walkway	√	X
Multi-purpose hall	√	X
Water bottling plant	√	X


√ = present x = absent



**Plate 1. A building showing relaxation centre with walk way at Ikogosi warm spring**



**Plate 2. Picture showing the source of warm spring at Ikogosi**



**Plate 3. Arinta waterfall**



Plate 4. Vegetation in one of the sampled plot in Arinta waterfall watershed

### 3.3 Richness of Major Plant Family in the Study Areas

Distribution of the tree in the two ecotourism centres by families revealed that a total of twenty-five (25) families of tree were identified in the two study areas with 17 and 20 families in Ikogosi warm spring and Arinta waterfall respectively (Table 3). Malvaceae has the highest species density (15). This was followed by the Fabaceae (12), *Anacardiaceae* and *Myristicaceae* with nine (9) each. At Ikogosi warm spring, the families with highest species density include *Anacardiaceae*, Fabaceae, *Malvaceae* and *Myristicaceae* with four (4) each. While in Arinta waterfall, the family *Malvaceae* has the highest species density of 11.

### 3.4 Diversity Indices of Trees in the Study Sites

Results on species diversity indices of Ikogosi warm spring and Arinta waterfall watersheds revealed that Shannon-Wiener diversity index of Arinta waterfall is 4.11 while that of Ikogosi warm spring is 3.48 (Table 4). Result on species evenness of Arinta waterfall was 0.92 while that of Ikogosi warm spring was 0.90. Sorenson's coefficient (SC) of similarity between the species showed a value 0.5. Also the Shannon-Wiener diversity index was 2.65 in Ikogosi warm spring while that of Arinta waterfall was 2.64. Family evenness in Arinta waterfall and Ikogosi warm spring are 0.70 and 0.83 respectively. Sorenson's coefficient (SC) of similarity between the families showed a value 0.71.

### 3.5 Vegetation Structure and Forest Characteristics of the Study Sites

Most of the tree species enumerated around the built-up/ recreational area of the study locations falls within the lower stratum of the forest structure. The middle layer category in Arinta waterfall has six (6) tree species while vegetation in Ikogosi has one (1). The result on lower stratum revealed more trees in Arinta waterfall (60) while that of Ikogosi warm spring has 35. The mean dbh of tree species in the lower stratum in Arinta waterfall was higher (97.78 cm) than that of Ikogosi warm spring (65.91 cm). Also, the mean height of tree species in the lower stratum in Arinta waterfall and Ikogosi warm spring are 9.55 m and 9.41m respectively.

## 4. DISCUSSION

The observed floristic richness of the vegetation in the two ecotourism sites revealed the true characteristics of a tropical forest whereby just two hectares could support 66 and 36 species. This observation corroborates the report by [23] and [24] who in separate studies recorded 67 plant species at Okomu National Park and 57 tree species at International Institute of Tropical Agriculture (IITA) forest reserve respectively. Comparatively lower number of tree species recorded at Ikogosi warm could be attributed to heavy presence of recreational/ infrastructural facilities at Ikogosi warm spring (Table 1) following the upgrading of Ikogosi warm spring to a resort centre of international standard by the State Government in the recent times. This

**Table 2. Tree species composition and classification of Ikogosi warm spring and Arinta waterfall watershed**

S/N	Species	Families	Ikogosi warm spring						Arinta waterfall					
			Freq	Dbh	Height	RD	RD <sub>0</sub>	IVI	Freq	Dbh	Height	RD	RD <sub>0</sub>	IVI
1	<i>Azelia Africana</i>	Fabaceae	3	0.63	9.15	4.2	2.3	3.24	0	0	0	0	0	0
2	<i>Albizia adianthifolia</i>	Fabaceae	3	0.54	12	4.2	1.70	2.94	0	0	0	0	0	0
3	<i>Alchornea cordifolia</i>	Fabaceae	0	0	0	0	0	0	2	0.85	9.12	1.27	0.57	0.92
4	<i>Alstonia booneii</i>	Euphorbiaceae	2	0.59	8.32	2.8	2	2.39	2	0.76	10.56	1.27	0.44	0.86
5	<i>Amphimas pterocpoides</i>	Apocynaceae	3	0.69	8.26	4.2	2.74	3.46	0	0	0	0	0	0
6	<i>Aningeria robusta</i>	Fabaceae	0	0	0	0	0	0	1	0.54	7.42	0.63	0.23	0.43
7	<i>Anthocleista vogelii</i>	Sapotaceae	4	1.64	11.52	5.6	15.63	10.6	0	0	0	0	0	0
8	<i>Anthonotha macrophylla</i>	Loganiaceae	1	0.61	9.2	1.4	2.15	1.77	0	0	0	0	0	0
9	<i>Antiaris africana</i>	Moraceae	4	0.47	9.58	5.6	1.26	3.41	3	3.5	12.52	1.9	9.51	5.71
10	<i>Artocarpus altilis</i>	Moraceae	0	0	0	0	0	0	2	3.5	12.52	1.27	9.51	5.39
11	<i>Bombax buonopozense</i>	Malvaceae	1	0.64	9.92	1.4	2.37	1.88	3	3.5	12.52	1.9	9.51	5.71
12	<i>Brachystegia evrycoma</i>	Moraceae	3	0.67	12.72	3	2.59	3.38	2	1.74	11.28	1.27	2.35	1.81
13	<i>Bridelia atroviridis</i>	Euphorbiaceae	1	0.47	9.36	1.4	1.26	1.33	0	0	0	0	0	0
14	<i>Canarium schweinfurthii</i>	Burseraceae	2	0.92	10.6	2.8	4.89	3.84	0	0	0	0	0	0
15	<i>Ceiba pentandra</i>	Malvaceae	2	0.67	10.8	2.7	2.59	2.69	0	0	0	0	0	0
16	<i>Chrysophyllum albidum</i>	Sapotaceae	1	0.59	8.77	1.4	2	1.7	1	0.64	9.92	0.63	0.32	0.48
17	<i>Cleistopholis pat</i>	Annonaceae	1	0.59	9.15	1.4	2	1.7	3	0.72	8.48	1.9	0.41	1.16
18	<i>Cola hispida</i>	Malvaceae	0	0	0	0	0	0	2	1.68	10.68	1.27	2.19	1.73
19	<i>Cola millenii</i>	Malvaceae	0	0	0	0	0	0	1	1.4	11.84	0.63	1.52	1.08
20	<i>Cola nitida</i>	Malvaceae	0	0	0	0	0	0	2	1.74	11.72	1.27	2.35	1.81
21	<i>Daniellia ogea</i>	Fabaceae	0	0	0	0	0	0	3	0.87	9.15	1.9	0.58	1.24
22	<i>Delonix regia</i>	Fabaceae	4	0.47	8.38	5.6	1.26	3.41	2	3.8	11.48	1.27	11.21	6.24
23	<i>Dracaena arborea</i>	Asparagaceae	1	0.47	8.38	1.4	1.26	1.33	0	0	0	0	0	0
24	<i>Elaeis guineensis</i>	Arecaceae	2	0.67	11.2	2.8	2.59	2.69	0	0	0	0	0	0
25	<i>Enantia chlorantha</i>	Annonaceae	0	0	0	0	0	0	1	1.4	10.88	0.63	1.52	1.08
26	<i>Entandrophragma cylindricum</i>	Meliaceae	1	0.62	8.72	1.4	2.22	1.81	3	0.54	10.5	1.9	0.23	1.07
27	<i>Ficus exasperate</i>	Moraceae	0	0	0	0	0	0	2	0.54	9.54	1.27	0.23	0.75
28	<i>Funtumia elastic</i>	Apocynaceae	0	0	0	0	0	0	2	1.37	10.44	1.27	1.45	1.36
29	<i>Gilbertiodendron dewevrei</i>	Fabaceae	2	0.78	9.52	2.8	3.56	3.17	0	0	0	0	0	0
30	<i>Gmelina arborea</i>	Labiatae	0	0	0	0	0	0	2	0.6	8.8	1.27	0.35	0.81
31	<i>Harungana Madagascariensis</i>	Hypericaceae	1	0.43	9.7	1.4	1.11	1.25	0	0	0	0	0	0
32	<i>Hollarhena floribunda</i>	Apocynaceae	0	0	0	0	0	0	1	0.85	12.68	0.63	0.57	0.60
33	<i>Hunteria umbellata</i>	Apocynaceae	0	0	0	0	0	0	3	1.64	10.36	1.9	2.09	1.99



S/N	Species	Families	Ikogosi warm spring						Arinta waterfall					
			Freq	Dbh	Height	RD	RD <sub>0</sub>	IVI	Freq	Dbh	Height	RD	RD <sub>0</sub>	IVI
34	<i>Macaranga spinosa</i>	Euphorbiaceae	2	0.67	9.88	2.8	2.59	2.69	3	0.72	9.4	1.9	0.41	1.16
35	<i>Malacantha alnifolia</i>	Sapotaceae	0	0	0	0	0	0	5	0.75	9.48	3.16	0.43	1.80
36	<i>Mallotus subulatus</i>	Euphorbiaceae	0	0	0	0	0	0	3	0.7	8.96	1.9	0.38	1.14
37	<i>Mangifera indica</i>	Anacardiaceae	1	0.78	9.58	1.4	3.56	2.48	2	0.84	9.98	1.27	0.54	0.91
38	<i>Milicia excels</i>	Moraceae	0	0	0	0	0	0	2	0.56	9.12	1.27	0.25	0.76
39	<i>Milicia regia</i>	Moraceae	1	0.67	8.64	1.4	2.59	1.99	4	0.92	10.85	2.53	0.65	1.59
40	<i>Mitragyna stipulosa</i>	Rubiaceae	2	0.51	9.68	2.8	1.48	2.13	1	0.68	9.12	0.63	0.36	0.5
41	<i>Monodora myristica</i>	Annonaceae	2	0.69	9.68	2.8	2.74	2.76	2	0.54	7.52	1.27	0.23	0.75
42	<i>Monodora tenuifolia</i>	Annonaceae	2	0.76	9.2	2.8	3.33	3.06	2	2.98	12.32	1.27	6.9	4.09
43	<i>Musanga cecropioides</i>	Urticaceae	2	0.68	10.4	2.8	2.67	2.73	2	0.74	9.38	1.27	0.43	0.85
44	<i>Myrianthus arboreus</i>	Moraceae	1	0.54	10.8	1.4	1.7	1.55	1	0.76	9.84	0.63	0.44	0.54
45	<i>Nauclea diderrichii</i>	Rubiaceae	0	0	0	0	0	0	1	0.6	9.28	0.63	0.3	0.47
46	<i>Nauclea latifolia</i>	Rubiaceae	0	0	0	0	0	0	2	0.78	8.16	1.27	0.47	0.87
47	<i>Nesogordonia papaverifera</i>	Malvaceae	2	0.78	8.16	2.8	3.56	3.17	2	1.04	9.86	1.27	0.84	1.06
48	<i>Newtonia buchanani</i>	Fabaceae	0	0	0	0	0	0	2	0.68	8.26	1.27	0.36	0.82
49	<i>Nothospondias staultii</i>	Anacardiaceae	1	0.53	8.7	1.4	1.63	1.51	2	0.52	9.84	1.27	0.21	0.74
50	<i>Ochroma lagopus</i>	Malvaceae	1	0.57	11.4	1.4	1.93	1.66	3	2.45	9.88	1.9	4.66	3.28
51	<i>Pentaclethra macrophylla</i>	Fabaceae	0	0	0	0	0	0	4	0.78	9.98	2.53	0.47	1.50
52	<i>Piptadeniastrum africanum</i>	Fabaceae	0	0	0	0	0	0	3	0.75	9.48	1.9	0.43	1.17
53	<i>Pseudospondia microcarpa</i>	Anacardiaceae	3	0.74	10.88	4.2	3.19	3.68	1	0.64	8.48	0.63	0.32	0.48
54	<i>Pseudospondias mombin</i>	Anacardiaceae	3	0.57	9.48	4.2	1.93	3.05	3	0.94	10.85	1.9	0.68	1.29
55	<i>Psydrax arnoldiana</i>	Rubiaceae	3	0.92	10.88	4.2	4.89	4.53	4	0.76	8.26	2.53	0.44	1.49
56	<i>Psydrax subcordata</i>	Rubiaceae	2	0.7	9.88	2.8	2.81	2.8	3	0.62	8.16	1.9	0.3	1.10
57	<i>Pterocarpus osun</i>	Leguminosae	2	0.57	9.98	2.8	1.93	2.36	2	0.64	8.15	1.27	0.32	0.80
58	<i>Pterygota macrocarpa</i>	Malvaceae	0	0	0	0	0	0	3	1.04	10.98	1.9	0.84	1.37
59	<i>Pycruanthus angolensis</i>	Myristicaceae	0	0	0	0	0	0	2	0.76	8.26	1.27	0.44	0.86
60	<i>Raphia hookeri</i>	Arecaceae	0	0	0	0	0	0	3	0.6	8.16	1.9	0.3	1.10
61	<i>Rauvolfia vomitoria</i>	Apocynaceae	0	0	0	0	0	0	1	1.5	8.96	0.63	1.75	1.19
62	<i>Rhodognaphalon brevicuspe</i>	Bombacaceae	0	0	0	0	0	0	3	1.19	9.98	1.9	1.10	1.50
63	<i>Riciodendron heudelotii</i>	Euphorbiaceae	0	0	0	0	0	0	1	0.98	12.88	0.63	0.74	0.69
64	<i>Rothimannia hispida</i>	Rubiaceae	0	0	0	0	0	0	2	0.62	9.28	1.27	0.30	0.79
65	<i>Senna siamea</i>	Fabaceae	0	0	0	0	0	0	3	0.61	8.34	1.9	0.29	1.10
66	<i>Spondias mombin</i>	Anacardiaceae	0	0	0	0	0	0	3	0.52	8.55	1.9	0.21	1.06
67	<i>Stercospermum acuminatissimu</i>	Bignoniaceae	0	0	0	0	0	0	2	0.78	9.44	1.27	0.47	0.87
68	<i>Sterculia rhinopetala</i>	Malvaceae	0	0	0	0	0	0	4	1.32	11.68	2.53	1.35	1.94

S/N	Species	Families	Ikogosi warm spring						Arinta waterfall					
			Freq	Dbh	Height	RD	RD <sub>0</sub>	IVI	Freq	Dbh	Height	RD	RD <sub>0</sub>	IVI
69	<i>Sterculia tragacantha</i>	Malvaceae	0	0	0	0	0	0	3	0.78	9.88	1.9	0.47	1.19
70	<i>Tabernamontana pachysiphon</i>	Apocynaceae	0	0	0	0	0	0	3	0.95	8.8	1.9	0.70	1.30
71	<i>Tectona grandis</i>	Verbenaceae	0	0	0	0	0	0	2	0.48	8.16	1.27	0.18	0.73
72	<i>Terminalia ivorensis</i>	Combretaceae	0	0	0	0	0	0	3	1.17	11.42	1.9	1.07	1.49
73	<i>Terminalia superba</i>	Combretaceae	0	0	0	0	0	0	2	0.64	8.64	1.27	0.32	0.80
74	<i>Theobroma cacao</i>	Malvaceae	0	0	0	0	0	0	3	0.72	9.52	1.9	0.41	1.16
75	<i>Triplochiton scleroxylon</i>	Malvaceae	0	0	0	0	0	0	1	0.7	9.36	0.63	0.43	0.53
76	<i>Uvariastrum pierreanum</i>	Annonaceae	0	0	0	0	0	0	4	3.57	11.88	2.53	9.90	6.22
77	<i>Voacanga Africana</i>	Apocynaceae	0	0	0	0	0	0	5	0.92	8.64	3.16	0.65	1.91
78	<i>Zanthoxylum gillettii</i>	Rutaceae	0	0	0	0	0	0	3	0.41	8.64	1.9	0.13	1.02
<b>Total</b>			<b>72</b>	<b>23.74</b>	<b>342.21</b>				<b>158</b>	<b>73.98</b>	<b>648.37</b>			

**Table 3. Contribution of family to tree species density in the study area**

SN	Families	Ikogosi warm spring	Arinta waterfall	Total Density
1	Anacardiaceae	4	5	9
2	Annonaceae	3	5	8
3	Apocynaceae	1	6	5
4	Arecaceae	1	1	2
5	Asparagaceae	1	0	1
6	Bignoniaceae	0	1	1
7	Burseraceae	1	0	1
8	Bombacaceae	0	1	1
9	Combretaceae	0	2	2
10	Euphorbiaceae	3	4	7
11	Fabaceae	4	8	12
12	Hypericaceae	1	0	1
13	Labiatae	0	1	1
14	Leguminosae	1	1	2
15	Loganiaceae	1	0	1
16	Malvaceae	4	11	15
17	Meliaceae	1	1	2
18	Moraceae	0	7	7
19	Myristicaceae	4	1	5
20	Rubiaceae	3	6	9
21	Rutaceae	0	1	1
22	Sapotaceae	2	2	4
24	Urticaceae	1	1	2
25	Verbenaceae	0	1	1
	<b>Total</b>	<b>36</b>	<b>66</b>	<b>102</b>

**Table 4. Diversity indices of tree species in the study areas**

Variables	Species		Family	
	Ikogosi	Arinta	Ikogosi	Arinta
Shannon_H	3.48	4.11	2.65	2.64
Evenness_e^H/S	0.90	0.92	0.83	0.70
Sorenson's coefficient		<b>0.5</b>		<b>0.71</b>

**Table 5. Number of species and characters of trees in different layers in the two sites**

Variables	Ikogosi warm spring	Arinta waterfall
Upper layer trees (≥ 22 m)	0	0
Mean DBH (cm)	0	0
Mean Height (m)	0	0
Middle layer trees (≥ 13 m ≤ 21 m)	1	6
Mean DBH (cm)	67	255.2
Mean Height (m)	12.72	12.57
lower stratum trees (≤ 12 m)	35	60
Mean DBH (cm)	65.91	97.78
Mean Height (m)	9.41	9.55

development has call for clearance of more land and consequent remover of trees in the build-up area. This assertion is in consonance with the findings of [25] on the negative impacts of recreation and tourism on plants communities in protected areas in Australia.

Biodiversity indices on plant species in the study area revealed that the general arrangement of plants across the two studied sites appears to be similar (Tables 2 and 3) with *Malacantha alnifolia*, *Voacanga africana* (5 each), *Anthocleista vogelii*, *Antiaris Africana* and *Delonix regia* (4 each) been the most frequently

encountered species in the study area. The dominant and the most important species in the study area is *Anthocleista vogelii* as revealed by the IVI (Table 2) while Malvaceae, Fabaceae, Anacardiaceae, Rubiaceae, Moraceae and Euphorbiaceae are the dominant family. This findings corroborated report of [26] that the tropical rainforest ecosystem of south west Nigeria is dominated by these sets of tree species and families. The preponderance of indigenous hardwood species at the two watersheds signifies little human interference. Similar observation has also been reported on tree species composition of Akure Strict Nature Reserve [27].

The high values of the Shannon indices in the two study sites revealed vegetation with very high tree species diversity and abundance. The Shannon indices of 3.48 and 4.11 obtained in the two watersheds are higher than the mean value of 3.34 obtained by [28] for sacred groves in south eastern Indian and closer to the average value (3.66) reported for some tropical rainforest sites in southern Nigeria by [29]. This result is an indication that the biological diversity of the watersheds is adequately conserved probably with local laws and taboos. Since one of the criteria for considering an ecosystem as a good ecotourism destination is its biodiversity richness, the observed floristic diversity of the forests at the two watersheds has great potentials for flora conservation and sustainable ecotourism development.

The slightly higher species evenness value observed in Arinta waterfall shows that the species are more evenly distributed than in Ikogosi warm spring. This result could be attributed to higher number of trees in the middle layer to form a continuous canopy. The horizontal and vertical structure of the forest at the watersheds as revealed by the diameter and height distribution indicates a forest structure that is immature and still expanding. This is evident in the floristic composition of the two watersheds which are within the middle and understory layer with small stem (Table 5). The predominantly middle and lower stratum vegetation of the two study sites especially the Arinta waterfalls could be an added advantage to a sustainable ecotourism sites if properly managed. This lower stratum with well-formed canopies could provide resting shade for visitors to the ecotourism centres. This assertion corroborates the submission of [30]

Observation from the study also revealed that some species are threatened and endangered, such species include *Anthonotha macrophylla*, *Aningeria robusta*, *Bridelia atroviridis*, *Cola milenii*, *Dracaena aborea*, *Harungana madagascariensis*, *Hollarhena floribunda*, *Ricinodendron heudelotii*, *Rauvolfia vomitoria*, *Triplochiton scleroxylon* and *Nauclea diderrichii* which were encountered once at the study sites. The implication of this is that by the virtue of their narrow range, they are vulnerable to extinction if proper conservation measures are not put in place.

## 5. CONCLUSION

Results from this study have revealed that the vegetation of the study area is a repository of many tree species with great potential for floral conservation and sustainable ecotourism development. Also the study has revealed that species diversity and abundance at the watersheds compared favourably with other similar forest ecosystem in the region. The study also revealed that the species composition at the two watersheds are within with middle and understory layers with small diameter indicates an immature forest structure that is still expanding. Observation from the study also revealed that some species are threatened and endangered. Consequent upon these results management strategies should be put in place to improve status of the watersheds. This could be done by involving the local communities in the planning and policy implementation meant protect the watershed from encroachment. Also conservation efforts should be stepped up for species with rarity index value to prevent them from going into extinction.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Abu JE, Adebisi LA. A review of Traditional Forest Uses. In: Popoola, (ed). Proceedings of a National Workshop Organized by FANCONSULT and Edo State Chapter of FAN held in Benin City, Edo State between 5<sup>th</sup> and 6<sup>th</sup> of September. 2002;42-50.
2. Koziell I, Saunders J. Living off Biodiversity: Exploring Livelihoods and

- Biodiversity in Natural Resources Management. IIED: London U.K. 2001;36.
3. Singh JS. The Biodiversity Crisis: A Multifaceted Review Current Science. 2002;82(6): 638-647
  4. UNEP. United nations environment programme: Word conservation monitoring center annual report. 2001;1-8.
  5. Hubbel SP. Neutral theory and evolution of ecological equivalence. Ecology. 2006; 87(6):1387-1398.
  6. Vetaas OR, Ferrer-Castan D. Patterns of woody plant species richness in Iberian Peninsula. Journal of Biogeography. 2008; 35(10):1863-1878.
  7. Bankole A. The Nigerian Tourism Sector: Economic Contribution, constraints and opportunities. Journal of Hospitality Financial Management. 2002;10(1):71-89.
  8. Pausas JG, Carreras J, Ferre'e A, Font X. Coarse scale plant species richness in relation to environmental heterogeneity. Journal of Vegetation Science. 2003;14: 661-668.
  9. Ijasan KC, Izobo MO. Assessing community engagement in tourism planning and development in Nigeria: A case study of Arinta waterfall tourist resort, Ipole Iloro Ekiti state. Transnational Journal of Science and Technnology. 2012;2(4):11-19.
  10. Gelormino E, Melis G, Marietta C, Costa G. From built environment to health inequalities: An explanatory framework based on evidence. Preventive Medicine Reports. 2015;2:737-745.
  11. Slinger-Friedman V. Ecotourism in Dominica: Studying the Potential for Economic Development, Environmental Protection and Cultural Conservation. Island Studies Journal. 2009;4(1):3-24.
  12. Stankov U, Stojanović V, Dragičević V, Arsenovic D. Ecotourism – An Alternative to Mass Tourism in Nature Park. Stara Planina. Journal of the Geographical Institute "Jovan Cvijie". 2011;61(1):43-59.
  13. Adekunle VAJ, Olagoke AO, Ogundare LF. Logging impact in tropical lowland humid forest on tree species diversity and environmental conservation. Applied Ecology and Environmental Research. 2013;11(3):491-511.
  14. Ogunjemite BG, Olaniyi OE, Akinwumi OO. Maintenance of ecological integrity of a warm spring site: The role of vegetation composition and cover of Ikogosi warm spring, Ekiti State, Nigeria. In Popoola, L., Ogunsanwo OY, Adewole NO (Eds.) Balancing environmental sustainability and livelihoods in an emerging economy. Proceedings of 36<sup>th</sup> annual conference of the Forestry Association of Nigeria, held at University of Uyo, Akwa Ibom State, Nigeria. 2013;553-562.
  15. Olaniyi OE, Ogunjemite BG. Ecotourism development in Ikogosi Warmspring, Ekiti-State, Nigeria: Implication on woody species composition and structure. Applied Tropical Agriculture. 2015;20(2):45-54.
  16. Kayode IB, Ayodele IA. Assessment of the Strength of Tourism Potentials of Ekiti State, Nigeria. Obeche Journal. 2012; 30(1):381-388. 0078-2912.
  17. Salami AT, Monitoring Nigerian forest with Nigeria Sat-1 and other satellites. In: Ayobami TS (Ed). Imperatives of space technology for sustainable forest management. Proceedings of an international stakeholders' workshop sponsored by National Space Research and Development Agency held in Abuja, Nigeria. 2006;26-61.
  18. Lawal A, Adekunle VAJ. A silvicultural approach to volume yield, biodiversity and soil fertility restoration of degraded natural forest in South west Nigeria. International Journal of Bio-diversity Science, Ecosystem Services and Management. 2013;9(3):201-214.
  19. Price PW. Insect ecology. 3<sup>rd</sup> Edition, Wiley, NY;1997.
  20. Kent M, Cooker P. Vegetation Description and Analysis: A practical Approach: Belhaven press, London. 1992;363.
  21. Brashears MB, Fajvan MA, Schuler TM. An assessment of canopy stratification and tree species diversity following clear cutting in Appalachian hardwoods. Forest Science. 2004;50(1):54-61.
  22. Longman KA, Jenik J. Tropical forest and its environment (2<sup>nd</sup> ed.). Tropical Ecology Series. John Wiley, New York. 1987;347.
  23. Oduwaiye EA, Oyeleye B, Oguntala AB. Species diversity and potential for forest regeneration in Okomu permanent sample plot: In Abu JE, Oni PI, Popoola L. (Eds.) Forestry and challenges of sustainable livelihood. Proceedings of the 28<sup>th</sup> annual conference of the Forestry Association of Nigeria, Akure. Ondo state, Nigeria. 2002; 264-272.
  24. Oladoye AO, Aduradola AM, Adedire MO, Agboola DA. Composition and stand

- structure of a regenerating tropical rainforest ecosystem in South-western Nigeria. *International Journal of Biodiversity and Conservation*. 2014; 6(11):765-776.
25. Pickering C, Hill W. Impacts of recreation and tourism on plants plant biodiversity and vegetation in protected areas in Australia. *Journal of Environmental Management*. 2007;85(4):791-800.
26. Adekunle VAJ, Olagoke AO, Ogundare LF. Rate of timber production in tropical rainforest ecosystem of southwest Nigeria and its implications on sustainable forest management. *Journal of Forestry Research*. 2010;21:225-230.
27. Adekunle VAJ, Olagoke AO, Ogundare LF. Logging impacts in tropical lowland humid forest on three species diversity and environmental conservation. *Applied Ecology and Environmental Research*. 2013;11(3): 491-511.
28. Rao BRP, Babu MVS, Reddy MS, Reddy AM, Rao VS, Sunitha S, Geneshaiah KN. Sacred groves in Southern Eastern Ghats, India: Are better managed than forest reserves. *Tropical Ecology*. 2011;52:79-90.
29. Adekunle VAJ, Adegoke AO. Diversity and biovolume of tree species in natural forest ecosystem in the bitumen-producing area of ondo state, Nigeria: A baseline study. *Biodiversity and Conservation*; 2007. DOI 101007/s10531-018-9279-y
30. Olaniyi OE, Ogunjemite BG. Ecotourism development in Ikogosi warm spring, Ekiti-State, Nigeria: Implication on woody species composition and structure. *Applied Tropical Agriculture*. 2015;20(2):45- 54.

© 2019 Olujobi et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<http://www.sdiarticle3.com/review-history/47239>